

On Energy Minimizers of the Diblock Copolymer Problem

Xiaofeng Ren *

Department of Mathematics and Statistics
Utah State University
Logan, UT 84322, USA

Juncheng Wei †

Department of Mathematics
The Chinese University of Hong Kong
Shatin, Hong Kong

February 19, 2003

Abstract

We view the free energy of a diblock copolymer system as a variational problem, in which the integrand of the functional contains an interesting nonlocal term, and a small parameter ϵ . We prove that as ϵ approaches 0, the energy minimizers develop a growing number, of the order $\epsilon^{-1/3}$, of periodic oscillations, explaining the micro-phase separation phenomenon.

1 Introduction

A di-block copolymer molecule is a linear-chain consisting of two sub-chains a and b grafted covalently to each other. The sub-chains a and b are made of different monomer units A and B , respectively. In polymer systems even a weak repulsion between unlike monomers A and B induces a strong repulsion between a and b . As a result the different sub-chains tend to segregate below some temperature T_c , but as they are chemically bonded, even a complete segregation of sub-chains a and b cannot lead to a macroscopic phase separation. Only a local micro-phase separation occurs: micro-domains rich in A and B are formed.

In [18] Ohta and Kawasaki introduced a free energy functional

$$\mathcal{F}(u) = \int_{\Omega} \left[\frac{\epsilon^2}{2} |\nabla u|^2 + W(u) + \frac{\sigma}{2} |(-\Delta)^{-1/2}(u - m)|^2 \right] dx.$$

*Supported in part by NSF grant DMS-9703727.

†Supported in part by an Earmarked Grant of RGC of Hong Kong.

The original formula in [18] is given for the whole space. The expression here on a bounded domain first appeared in Nishiura and Ohnishi [10].

The two unlike monomer units are represented by $u = -1$ and $u = 1$ respectively. The connectivity of the monomers in a chain leads to the long range interaction $\frac{\sigma}{2}|(-\Delta)^{-1/2}(u - m)|^2$ in the free energy. Here $-\Delta$ is viewed as a positive operator, and $(-\Delta)^{-1/2}$ is the square root of its inverse. The parameter σ is proportional to the inverse of the square root of the total chain length of the copolymer. $\frac{\epsilon^2}{2}|\nabla u|^2$ represents the inter-facial energy density at bonding points. The parameter ϵ is proportional to the thickness of interfaces between the two monomers. m stands for the mass ratio of the two monomer units.

When this free energy is minimized, the first term of the integrand prefers large blocks of monomers, thereby reducing the combined size of interfaces between the two monomers. The function W in the second term is a double well potential with two global minima at -1 and 1 , reflecting its preference for segregated monomers over mixtures. The third term is most interesting to us, which depends on u nonlocally, through a global operator $(-\Delta)^{-1/2}$. It favors rapid oscillation between the two monomer. When all these factors compete, the phenomenon known as micro-phase separation occurs.

The one dimensional case $\Omega = (0, 1)$ is particularly interesting because of the laminar structures observed in di-block copolymers. In an earlier paper [12] we studied the parameter range $\sigma \sim \epsilon$. Physically this means that the size of the sample is of the order $N^{2/3}l$ where N , the polymerization index, is the number of monomers in a chain molecule and l is the average distance between two adjacent monomers. We proved the existence of a family of local minima when ϵ is small, which are nearly periodic with the sizes of periods comparable to the size of domain $(0, 1)$.

In this paper we study a different parameter range $\sigma \sim 1$. Physically we are taking a larger sample of the size Nl . The admissible set is

$$X_m = \{u \in W^{1,2}(0, 1) : \int_0^1 u(x) dx = m\}, \quad m \in (-1, 1) \quad (1.1)$$

The constraint $\int_0^1 u = m$ reflects the total mass of one of the two micro-components. It must be in $(-1, 1)$ in order to have a mix of the two monomer units, ($u = -1$ and $u = 1$ respectively).

We re-state the functional

$$I_\epsilon(u) = \int_0^1 \left[\frac{\epsilon^2}{2}|u'|^2 + W(u) + \frac{1}{2}|(-D^2)^{-1/2}(u - m)|^2 \right] dx, \quad (1.2)$$

which we call the energy of u . The second order derivative operator

$$-D^2 : \{v \in W^{2,2} : v'(0) = v'(1) = 0, \int_0^1 v = 0\} \rightarrow \{w \in L^2 : \int_0^1 w = 0\}$$

is an isometry. Its inverse is positive from $\{w \in L^2 : \int_0^1 w = 0\}$ to its self. We denote the square root of this inverse by $(-D^2)^{-1/2}$. For every $u \in X_m$ we can

solve

$$-v'' = u - m, v'(0) = v'(1) = 0, \int_0^1 v = 0$$

for v . Such v is often denoted by $(-D^2)^{-1}(u - m)$. Then (1.2) becomes

$$I_\epsilon(u) = \int_0^1 \left[\frac{\epsilon^2}{2} |u'|^2 + W(u) + \frac{1}{2} |v'|^2 \right] dx. \quad (1.3)$$

Let u_ϵ be a global minimum of I_ϵ in X_m , i.e.

$$I_\epsilon(u_\epsilon) = \min_{u \in X_m} I_\epsilon(u). \quad (1.4)$$

The existence of u_ϵ is guaranteed by the usual variational argument. u_ϵ solves the Euler-Lagrange equation

$$-\epsilon^2 u'' + f(u) + (-D^2)^{-1}(u - m) = \lambda$$

where $f = W'$. The constant λ , the Lagrange multiplier, is unknown.

Defining $v_\epsilon = (-D^2)^{-1}(u_\epsilon - m)$ and λ_ϵ to be the Lagrange multiplier associated with u_ϵ , we rewrite the Euler-Lagrange equation for u_ϵ , v_ϵ and λ_ϵ as

$$\begin{cases} -\epsilon^2 u'' + f(u) + v = \lambda \\ -v'' = u - m \\ u'(0) = u'(1) = v'(0) = v'(1) = 0 \\ \int_0^1 u = m, \int_0^1 v = 0. \end{cases} \quad (1.5)$$

Note that without the nonlocal interaction term in (1.2) we have the more familiar functional

$$K_\epsilon(u) = \int_0^1 \left[\frac{\epsilon^2}{2} |u'|^2 + W(u) \right] dx. \quad (1.6)$$

Minimizers in X_m of K_ϵ are well known. When ϵ is small, K_ϵ has two global minima. One of them has a transition layer, whose width is of order ϵ , from -1 to 1 . The second is the reversal, i.e. the reflection with respect to the vertical line at $1/2$, of the first, (see Carr, Gurtin and Slemrod [1]).

The goal of this paper is to prove the following three theorems for the global minima of the nonlocal problem I_ϵ .

Theorem 1.1 *For small ϵ every global minimum u_ϵ is necessarily periodic, with exactly $N_\epsilon/2$ periods, where N_ϵ is the number of the transition layers of u_ϵ .*

Theorem 1.2 *For small ϵ I_ϵ has either two or four global minima. The case of two global minima is generic.*

Theorem 1.3 *The period of the global minima of I_ϵ has the asymptotic expansion $(\frac{96c_0\epsilon}{(1-m^2)^2})^{1/3} + O(\epsilon^{2/3})$, where c_0 is defined in (2.6).*

The proofs are rather straightforward, though some estimates in this paper look tedious. We obtain sharp lower and upper bounds for $I_\epsilon(u_\epsilon)$. The upper bound is deduced by a test function argument. The lower bound, which is harder to come by, comes after a careful study of u_ϵ .

With these bounds we study the length scale between adjacent transition layers of u_ϵ . A layer is characterized by a point x where $u_\epsilon(x)$ is not close to -1 or 1 . For technical reasons we set a value $\alpha \in (-1, 1)$, defined in (2.5), and say that x is an α -point if $u_\epsilon(x) = \alpha$. An α -point thus identifies a transition layer. We show that the distance between any two adjacent α -points of u_ϵ is comparable to $\epsilon^{1/3}$.

The proof of this fact is in Sections 6 and 7. We denote intervals separated by the α -points by p_i and q_i . On an interval p_i , u_ϵ is greater than α and on an interval q_i , u_ϵ is less than α . In Proposition 6.1 we show that $p_i = O(\epsilon^{1/3})$ and $q_i = O(\epsilon^{1/3})$. Then in Proposition 7.1 we improve the two estimates to $p_i \sim \epsilon^{1/3}$ and $q_i \sim \epsilon^{1/3}$.

Proposition 7.1 has the implication that the distance between any two adjacent zeros of v'_ϵ is also comparable to $\epsilon^{1/3}$. It allows us to localize I_ϵ to intervals separated by these zeros. After rescaling such intervals to $(0, 1)$ we obtain a problem similar to I_ϵ , but with a different parameter range. This new problem was the same as the one studied by the authors in [12]. The three theorems follow from some of its convexity properties.

The most important step in proving Proposition 6.1 is the establishment of a good lower bound for $I_\epsilon(u_\epsilon)$ in Section 5. This idea was used by Ni, Takagi and the second author in a series of papers, e.g. [7, 8, 9, 21], however in different settings. There the solutions are all spiky, instead of being periodic.

The special case that $m = 0$ and $W(-r) = W(r)$ was studied by Müller in [6]. He actually had a different looking functional

$$\tilde{I}_\epsilon(w) = \int_0^1 [\epsilon^2 |w''|^2 + \tilde{W}(w') + w^2] dx$$

in the admissible set $\{w \in W^{2,2}(0, 1) : w(0) = w(1) = 0\}$. Under the assumption $W(r) = W(-r)$, it was proved in [6] that global minima of \tilde{I}_ϵ are periodic.

\tilde{I}_ϵ itself has an interpretation in the elasticity theory. Imagine w as the displacement of an elastic bar under a loading device. w' is the strain field. The deformation of w gives rise to some elastic energy whose density is $\epsilon^2 |w''|^2 + W(w')$. Also assume that the bar is placed on an elastic foundation. The foundation interacts with the bar and contributes to some more energy with density w^2 . Adding these two terms we arrive at \tilde{I}_ϵ , the total energy of the system. See Truskinovsky and Zanzotto [19, 20] for more details.

To see how \tilde{I}_ϵ is related to I_ϵ , let u be an element in X_m and $v = (-D^2)^{-1}(u - m)$. Set $w = v'$. Then $w' = v'' = m - u$, $w'' = -u'$, and

$$I_\epsilon(u) = \frac{1}{2} \int_0^1 [\epsilon^2 |w''|^2 + 2W(m - w') + w^2] dx = \frac{1}{2} \tilde{I}_\epsilon(w),$$

if $\tilde{W}(r) = 2W(m - r)$. Since both \tilde{W} and W have two global minima at -1 and

1, m must be 0. What was proved in [6] translates to that when $W(r) = W(-r)$ and $m = 0$, the global minimizers of I_ϵ are periodic.

$W(r) = W(-r)$ may look like a technical restriction, but actually it, together with $m = 0$, imposes mathematically a symmetry within each period of a minimizer u_ϵ . If T is a period, then $u_\epsilon(x) = -u_\epsilon(T - x)$ for $x \in (0, T)$. The use of this symmetry is a key ingredient in [6]. In terms of application $m = 0$ requires that each of the two monomer units make exactly half of the volume, not a suitable condition for general copolymers.

We will prove the three theorems without assuming $W(r) = W(-r)$ or $m = 0$. Within each period, u_ϵ has no more symmetry. Instead u_ϵ is close to 1 on a portion of the period and close to -1 on another portion, generally of a different size, leaving the average of u_ϵ equal to m .

Our approach to the general case departs significantly from Müller's, when we analyze the important quantity $E(\epsilon, l)$, defined at the beginning of Section 10. l here is the distance between two adjacent zeros of v'_ϵ . In the symmetric case ($W(r) = W(-r)$, $m = 0$) E is convex with respect to l in a wide range of ϵ and l : $\epsilon \leq Cl/|\log l|$, as shown in [6]. This fact depends on a lower bound of eigenvalues of a linear problem (see Proposition 9.1 and the remark after its proof), when the symmetry condition is imposed. Without symmetry that linear problem has small eigenvalues. It turns out that the convexity of E is valid if we can show that ϵ and l lie in a narrower range: $C_1\epsilon^{1/3} \leq l \leq C_2\epsilon^{1/3}$. The remarks earlier after the statements of the three theorems explained how we prove this difficult estimate.

Other references on this subject include Ohnishi *et al* [11], Fife and Hillhorst [3], Choksi [2], Henry [4], Ren and Wei [13, 14, 15, 16, 17], and Muratov [5].

When estimating quantities, we adopt $O(\dots)$, $o(\dots)$, \sim convention. A term, say v_ϵ , satisfies $v_\epsilon = O(\epsilon^{1/3})$ if there exists a constant C independent of ϵ such that $|v_\epsilon(x)| \leq C\epsilon^{1/3}$ for all $x \in (0, 1)$. A term, say v_ϵ , satisfies $v_\epsilon = o(\epsilon^{1/3} \log \epsilon)$ if there exists a function $C(\epsilon)$, $C(\epsilon) \searrow 0$ as $\epsilon \searrow 0$, such that $|v_\epsilon(x)| \leq C(\epsilon)\epsilon^{1/3} \log \epsilon$ for all x . $O(\dots)$ and $o(\dots)$ also appear in inequalities. For instance, a term, say u_ϵ , satisfies $u_\epsilon \leq 1 + O(\epsilon^{1/3})$ if there exists $C > 0$ such that $u_\epsilon(x) - 1 \leq C\epsilon^{1/3}$ for all x . \sim indicates a comparable relation between two quantities. A term, say p_i , satisfies $p_i \sim \epsilon^{1/3}$ if there exist constants C_1 and C_2 such that $C_1\epsilon^{1/3} \leq p_i \leq C_2\epsilon^{1/3}$ for all i .

We require that all estimating quantities, like C , C_1 , C_2 , or $C(\cdot)$, depend on m and the overall shape of W only. Therefore all estimates involving O , o or \sim in this paper are *uniform* with respect to any variable/parameter that may appear, like x in $v_\epsilon(x)$ and i in p_i .

2 The local energy functional K_ϵ

The function W in the definition of I_ϵ is a balanced double well. More precisely

1. $W : (-\infty, \infty) \rightarrow [0, \infty)$ is C^5 ,
2. $W(r) = 0$ at $r = -1$ and $r = 1$, and $W(r) > 0$ at any other r ,

3. There exist a and b , $a > -1$, $a < b$, $b < 1$ such that $W''(r) > 0$ on $(-\infty, a) \cup (b, \infty)$ and $W''(r) < 0$ on (a, b) ,
4. W'' is bounded,
5. W' grows linearly, i.e. there exist C_1 and C_2 such that $C_1|r| \leq |W'(r)| \leq C_2|r|$ when r is large.

We have made these conditions consistent with the ones in the reference papers, like [1, 6]. The derivative of W is always denoted by f , and the local maximum of W between -1 and 1 by ω .

Next we list some well-known properties of the equation

$$-U'' + f(U) = 0. \quad (2.1)$$

It has the first integral

$$-|U'|^2 + 2W(U) = 2\gamma. \quad (2.2)$$

This first integral gives us a phase portrait of trajectories in the U vs. U' plane. The two equilibria $(-1, 0)$, $(1, 0)$ correspond to the two global minima of W at -1 and 1 . The third equilibrium $(\omega, 0)$, $\omega \in (-1, 1)$, comes from the local maximum ω of W . There are two heteroclinic orbits connecting $(-1, 0)$ to $(1, 0)$. They bound a family of periodic trajectories that in turn enclose $(\omega, 0)$. The remaining trajectories are unbounded.

One heteroclinic solution is denoted by H which solves

$$-H'' + f(H) = 0, \quad H(0) = \alpha, \quad H(\pm\infty) = \pm 1. \quad (2.3)$$

The constant α is a number between -1 and 1 defined later in (2.7) to identify transition layers. H has the first integral

$$-|H'|^2 + 2W(H) = 0. \quad (2.4)$$

Lemma 2.1 *1. There exists $C > 0$ such that as $t \rightarrow \pm\infty$, $H(t) = \pm 1 + O(e^{-C|t|})$, $H'(\pm t) = O(e^{-Ct})$, and $H''(\pm t) = O(e^{-Ct})$.*

2. Let G_s , $s > 0$, be the increasing solution of $-G_s'' + f(G_s) = 0$ with $G_s(0) = \alpha$ and $G_s'(s) = 0$. Then $\|G_s - H\|_{L^\infty(0,s)} = O(e^{-\nu s})$ for a constant $\nu > 0$. If G_s is the decreasing solution of the same equation and boundary conditions, then $\|G_s - H(\cdot)\|_{L^\infty(0,s)} = O(e^{-\nu s})$.

Proof. 1. From (2.4) we obtain

$$t = \int_0^t d\tau = \int_0^t \frac{dH(\tau)}{H'(\tau)} = \int_\alpha^{H(t)} \frac{dH}{\sqrt{2W(H)}} \sim -\log(1 - H(t)).$$

The convergence rates at ∞ then follow. The case of $t \rightarrow -\infty$ is similar.

2. The constant γ in (2.2) is $W(G_s(s))$ when $U = G_s$. The estimate in this part follows by comparing the time variable

$$t = \int_0^t d\tau = \int_0^{G_s(t)} \frac{dG_s}{\sqrt{2W(G_s) - 2W(G_s(s))}}$$

of G_s with that of H in part 1. \square

Lemma 2.2 1. Let G be a bounded solution of $-\Psi'' + f'(H)\Psi = 0$ on (a, ∞) , $(-\infty, a)$, or $(-\infty, \infty)$, where H is the heteroclinic solution defined in (2.3). Then there exists a constant c such that $\Psi = cH'$ and $H' \in W^{1,2}(-\infty, \infty)$.

2. There exists a constant $\iota > 0$ such that for every $\Phi \in W^{1,2}(-\infty, \infty)$ with $\int_{-\infty}^{\infty} \Phi H' dt = 0$, $\int_{-\infty}^{\infty} [|\Phi'|^2 + f'(H)\Phi^2] dt \geq \iota \int_{-\infty}^{\infty} \Phi^2 dt$.

Proof. 1. H' is obviously a solution of the linear equation. It is bounded and positive. Another linearly independent solution is $R(t) = H'(t) \int_0^t \frac{ds}{(H'(s))^2}$. Then there exist c and c^* such that $\Psi = cH' + c^*R$. However $R(\pm\infty) = \pm\infty$, while Ψ is bounded. So $c^* = 0$.

To see $H' \in W^{1,2}(-\infty, \infty)$ we return to the first integral (2.4), the equation (2.3), and the phase portrait, to compute

$$\int_{-\infty}^{\infty} (H'(t))^2 dt = \int_{-1}^1 \sqrt{2W(H)} dH, \quad \int_{-\infty}^{\infty} (H''(t))^2 dt = \int_{-1}^1 \frac{(f(H))^2}{\sqrt{2W(H)}} dH.$$

Both integrals on the right sides are convergent.

2. H is a global minimum of $\int_{-\infty}^{\infty} [\frac{1}{2}|G'|^2 + W(G)] dt$ in $\{G \in W_{loc}^{1,2}(-\infty, \infty) : G(\pm\infty) = \pm 1\}$. 0 is the principal eigenvalue of the second variation at H , corresponding to an eigen-function H' . The next eigenvalue gives rise to ι . \square

Let $\alpha \in (-1, 1)$ be the number so that

$$\frac{\int_{\alpha}^1 \sqrt{W(s)} ds}{1+m} = \frac{\int_{-1}^{\alpha} \sqrt{W(s)} ds}{1-m}. \quad (2.5)$$

Also define

$$c_{-1} = \sqrt{2} \int_{-1}^{\alpha} \sqrt{W(s)} ds, \quad c_1 = \sqrt{2} \int_{\alpha}^1 \sqrt{W(s)} ds, \quad c_0 = c_{-1} + c_1. \quad (2.6)$$

(2.5) implies that

$$\frac{c_1}{1+m} = \frac{c_{-1}}{1-m}. \quad (2.7)$$

The number α will be used to identify transition layers. If u_{ϵ} is a global minimum of I_{ϵ} in X_m , we say $x \in (0, 1)$ is a α -point of u_{ϵ} if $u_{\epsilon}(x) = \alpha$. Of course any number in $(-1, 1)$ can be used to identify transition layers of u_{ϵ} . The reason why we choose this particular value will come out in Section 6.

Finally we consider the functional K_{ϵ} in (1.6) on various admissible sets. Let

$$\begin{aligned} k(\epsilon) &= \min\{K_{\epsilon}(u) : u \in X_m\} \\ k_{-1}(\epsilon) &= \min\{K_{\epsilon}(u) : u \in W^{1,2}(0, 1), u(0) = u(1) = \alpha, u \leq \alpha\}, \\ k_1(\epsilon) &= \min\{K_{\epsilon}(u) : u \in W^{1,2}(0, 1), u(0) = u(1) = \alpha, u \geq \alpha\}, \\ k_{-1}^h(\epsilon) &= \min\{K_{\epsilon}(u) : u \in W^{1,2}(0, 1), u(0) = \alpha, u \leq \alpha\}, \\ k_1^h(\epsilon) &= \min\{K_{\epsilon}(u) : u \in W^{1,2}(0, 1), u(0) = \alpha, u \geq \alpha\}. \end{aligned} \quad (2.8)$$

Lemma 2.3 *There exists $\mu > 0$ for the following statements.*

1. $k(\epsilon) = c_0\epsilon + O(\epsilon^{-\mu/\epsilon})$.
2. $k_{-1}(\epsilon) = 2c_{-1}\epsilon + O(e^{-\mu/\epsilon})$.
3. $k_1(\epsilon) = 2c_1\epsilon + O(e^{-\mu/\epsilon})$.
4. $k_{-1}^h(\epsilon) = c_{-1}\epsilon + O(e^{-\mu/\epsilon})$.
5. $k_1^h(\epsilon) = c_1\epsilon + O(e^{-\mu/\epsilon})$.

Proof. Part 1 was proved in [1], Theorem 8.1. The proofs of 2-5 are standard and we only show a sketch for 5.

Recall H in (2.3). Use $H(\frac{x}{\epsilon}) \geq \alpha$ on $(0, 1)$ as a test function to compute $K_\epsilon(H(\frac{\cdot}{\epsilon}))$. Because of (2.3), we find

$$K_\epsilon(H(\frac{\cdot}{\epsilon})) = \sqrt{2} \int_0^1 \sqrt{W(H)} H'(\frac{x}{\epsilon}) dx = \epsilon \sqrt{2} \int_\alpha^{H(1/\epsilon)} \sqrt{W(H(t))} dt.$$

Due to the exponential convergence rate of $H(t) \rightarrow 1$ as $t \rightarrow \infty$, Lemma 2.1₁,

$$k_1^h(\epsilon) \leq \epsilon c_1 + O(\epsilon^{-C/\epsilon}). \quad (2.9)$$

Now we show that the inequality (2.9) is indeed an equality. Let w_ϵ be a global minimum of K_ϵ in the admissible set $\{u \in W^{1,2}(0, 1) : u(0) = \alpha, u \geq \alpha\}$, whose existence is guaranteed by the theory of obstacle problems. w_ϵ satisfies the variational inequality

$$\int_0^1 [\epsilon^2 w'_\epsilon(\phi' - w'_\epsilon) + f(w_\epsilon)(\phi - w_\epsilon)] dx \geq 0 \quad (2.10)$$

for every ϕ in the same admissible set.

The theory of variational inequalities asserts that $w_\epsilon \in W^{2,2}(0, 1)$. Let $S = \{x \in (0, 1) : w_\epsilon(x) = \alpha\}$, $U = (0, 1) \setminus S$. Then U is open and S relatively closed in $(0, 1)$. We show that $S = \emptyset$. Let $\bar{x} \in S$. Then $w_\epsilon(\bar{x}) = \alpha$ and $w'_\epsilon(\bar{x}) = 0$. It follows from (2.10) that

$$-\epsilon^2 w''_\epsilon + f(w_\epsilon) = 0 \quad (2.11)$$

on U . If we multiply the equation by w'_ϵ , then since $w'_\epsilon = 0$ on S , on the whole $(0, 1)$ there is a first integral

$$-\epsilon^2 |w'_\epsilon|^2 + 2W(w_\epsilon) = -\epsilon^2 |w'_\epsilon(\bar{x})|^2 + 2W(w_\epsilon(\bar{x})) = 2W(\alpha).$$

This implies that $W(w_\epsilon) \geq W(\alpha)$. Then $K_\epsilon(w_\epsilon) \geq W(\alpha) > 0$, which is inconsistent with (2.9) for small ϵ . This proves that no such \bar{x} exists and $S = \emptyset$. So w_ϵ solves (2.11) on $(0, 1)$.

At $x = 1$, (2.10) allows two possibilities. Either A: $w_\epsilon(1) > \alpha$ and $w'_\epsilon(1) = 0$, or B: $w_\epsilon(1) = \alpha$.

We first consider case A. Set $x = \epsilon t$, $U(t) = w_\epsilon(\epsilon t)$. We suppress the dependence of U on ϵ to keep notations simple. U satisfies (2.2). The constant γ there can be evaluated at $t = 1/\epsilon$ where $U'(1/\epsilon) = 0$. So $\gamma = W(U(1/\epsilon))$.

As $\epsilon \searrow 0$, $U'(0) \nearrow H'(0)$, $\gamma \searrow 0$ and the trajectory of U , which is a periodic orbit inside the two heteroclinic orbits, approaches that of H . It also follows that $U(1/\epsilon)$ tends to 1 from the left. Without ambiguity, for small ϵ denote this $U(1/\epsilon) = W^{-1}(\gamma)$.

Now we view γ , instead of ϵ , as the controlling parameter. (2.2) yields that the duration

$$\frac{1}{\epsilon} = \int_0^{1/\epsilon} dt = \int_\alpha^{W^{-1}(\gamma)} \frac{dU}{\sqrt{2(W(U) - \gamma)}} \sim \log \gamma,$$

and an estimate for the local energy

$$\begin{aligned} \epsilon^{-1} K_\epsilon(w_\epsilon) - c_1 &= \int_0^{1/\epsilon} \left[\frac{|U'|^2}{2} + W(U) \right] dt - c_1 \\ &= \int_\alpha^{W^{-1}(\gamma)} \frac{2W(U) - \gamma}{\sqrt{2(W(U) - \gamma)}} dU - c_1 \\ &\sim \gamma \log \gamma, \end{aligned}$$

as $\gamma \searrow 0$. So follows the estimate in 5. of this lemma.

Finally we rule out case B. If we again set $U = w_\epsilon(\epsilon t)$, then in the phase portrait this solution corresponds to a part of a periodic trajectory as well. However at $t = 1/\epsilon$, $(U(1/\epsilon), U'(1/\epsilon))$ is the mirror image of $(U(0), U'(0))$ about the horizontal axis. After a similar argument of phase plane analysis, we find $K_\epsilon(w_\epsilon) = 2c_1\epsilon + O(e^{-\mu/\epsilon})$, contradicting (2.9). \square

The constants μ in Lemma 2.3 and ν in Lemma 2.1 are henceforth fixed. They depend on W and m only.

Lemma 2.4 *Let w_ϵ be a global minimum of K_ϵ in X_m . Define*

$$w_1(x) = \begin{cases} -1, & x \in (0, \frac{1-m}{2}) \\ 1, & x \in (\frac{1-m}{2}, 1) \end{cases} \quad w_2(x) = \begin{cases} 1, & x \in (0, \frac{1+m}{2}) \\ -1, & x \in (\frac{1+m}{2}, 1) \end{cases}$$

Then either $\int_0^1 |w_\epsilon - w_1| dy = O(\epsilon \log \epsilon)$, or $\int_0^1 |w_\epsilon - w_2| dy = O(\epsilon \log \epsilon)$. For small ϵ w_ϵ is increasing in the first case and decreasing in the second case.

Proof. See Theorems 3.1 and 9.1 of [1]. \square

3 An upper bound of $I_\epsilon(u_\epsilon)$

Let us agree on the notation $\text{Ave}(w)$ for the mean of w , i.e. if w is defined on (a, b)

$$\text{Ave}(w) = \frac{\int_a^b w(x) dx}{b - a}. \quad (3.1)$$

Lemma 3.1 1. For every positive integer N

$$I_\epsilon(u_\epsilon) \leq c_0 \epsilon N + \frac{(1-m^2)^2}{24N^2} + O\left(-\frac{\epsilon}{N} \log(\epsilon N) + e^{-\frac{\mu}{\epsilon N}}\right).$$

2. When N is taken to be the integer closest to $\left(\frac{(1-m^2)^2}{12\epsilon c_0}\right)^{1/3}$,

$$I_\epsilon(u_\epsilon) \leq c_0^{2/3} (1-m^2)^{2/3} \left(\frac{9}{32}\right)^{1/3} \epsilon^{2/3} + O(\epsilon^{4/3} \log \epsilon).$$

Proof. Let N be a positive integer and $(0, 1)$ be equally divided by N . Set $l = \frac{1}{N}$. Minimize among $u \in W^{1,2}(0, l)$, subject to $\text{Ave}(u) = m$,

$$\int_0^l \left[\frac{\epsilon^2}{2} |u'|^2 + W(u) \right] dx$$

to find $u_{0,\epsilon}$. By rescaling $x = lz$, $\mathcal{U}_{0,\epsilon}(z) = u_{0,\epsilon}(lz)$ minimizes $K_{\frac{\epsilon}{l}}$ in X_m and

$$\begin{aligned} \int_0^l \left[\frac{\epsilon^2}{2} |u'_{0,\epsilon}|^2 + W(u_{0,\epsilon}) \right] dx &= l \int_0^1 \left[\frac{\epsilon^2}{2l^2} |\mathcal{U}'_{0,\epsilon}|^2 + W(\mathcal{U}_{0,\epsilon}) \right] dz \\ &= l K_{\frac{\epsilon}{l}}(\mathcal{U}_{0,\epsilon}) = lk \left(\frac{\epsilon}{l}\right). \end{aligned}$$

Extending $u_{0,\epsilon}$ to $(0, 1)$ by anti-symmetric reflection and using it as a test function for an upper bound of $I_\epsilon(u_\epsilon)$, we find

$$\int_0^1 \left[\frac{\epsilon^2}{2} |u'_{0,\epsilon}|^2 + W(u_{0,\epsilon}) \right] dx = Nlk \left(\frac{\epsilon}{l}\right) = k(\epsilon N) = c_0 \epsilon N + O\left(e^{-\frac{\mu}{\epsilon N}}\right), \quad (3.2)$$

where the last equation comes from Lemma 2.3.

To estimate the nonlocal part of $I_\epsilon(u_{0,\epsilon})$, let $v_{0,\epsilon}$ be the solution of $-v'' = u_{0,\epsilon} - m$, $v'(0) = v'(l) = 0$, $\text{Ave}(v) = 0$. Through anti-symmetric reflection $v_{0,\epsilon}$ is extended to $(0, 1)$ and $v_{0,\epsilon} = (-D^2)^{-1}(u_{0,\epsilon} - m)$.

Estimate $v_{0,\epsilon}$ by comparing it with v_0 which solves $-v'' = u_0 - m$, $v'(0) = v'(l) = 0$, $\text{Ave}(v) = 0$. Here u_0 is a step function with one jump from -1 to 1 , satisfying $\text{Ave}(u_0) = m$. Scale $(0, l)$ to $(0, 1)$. Let $\mathcal{U}_0(z) = u_0(lz)$, i.e.

$$\mathcal{U}_0(z) = \begin{cases} -1, & z \leq (1-m)/2 \\ 1, & z > (1-m)/2 \end{cases}. \quad (3.3)$$

Let $\mathcal{V}_0(z) = l^{-2} v_{0,\epsilon}(lz)$. Then $\mathcal{V}_0 = (-D^2)^{-1}(\mathcal{U}_0 - m)$.

We record the expression for \mathcal{V}_0 for later purposes.

$$\mathcal{V}_0 = \begin{cases} \frac{1+m}{2} \left[z^2 - \left(\frac{1-m}{2}\right)^2 \right] - \frac{(1-m^2)m}{6}, & z \in \left[0, \frac{1-m}{2}\right] \\ -\frac{1-m}{2} \left[(1-z)^2 - \left(\frac{1+m}{2}\right)^2 \right] - \frac{(1-m^2)m}{6}, & z \in \left[\frac{1-m}{2}, 1\right] \end{cases}. \quad (3.4)$$

Recall $\mathcal{U}_{0,\epsilon}(z) = u_{0,\epsilon}(lz)$. Define $\mathcal{V}_{0,\epsilon}(z) = l^{-2}v_{0,\epsilon}(lz)$. It is clear that $\mathcal{V}_{0,\epsilon} = (-D^2)^{-1}(\mathcal{U}_{0,\epsilon} - m)$. Therefore $\|v_{0,\epsilon}\|_\infty = l^2\|\mathcal{V}_{0,\epsilon}\|_\infty = O(l^2)$ and $\|v_0\|_\infty = l^2\|\mathcal{V}_0\|_\infty = O(l^2)$.

Apply Lemma 2.4 to $\mathcal{U}_{0,\epsilon}$, a minimum of $K_{\frac{\epsilon}{l}}$, to obtain

$$\int_0^1 |\mathcal{U}_{0,\epsilon} - \mathcal{U}_0| dz = O\left(\frac{\epsilon}{l} \log\left(\frac{\epsilon}{l}\right)\right),$$

which yields

$$\int_0^l |u_{0,\epsilon} - u_0| dx = l \int_0^1 |\mathcal{U}_{0,\epsilon} - \mathcal{U}_0| dz = lO\left(\frac{\epsilon}{l} \log\left(\frac{\epsilon}{l}\right)\right) = O\left(\epsilon \log\left(\frac{\epsilon}{l}\right)\right).$$

Then by multiplying the equation $-D^2w = u_{0,\epsilon} - u_0$ that $v_{0,\epsilon} - v_0$ satisfies by $v_{0,\epsilon} + v_0$ and integrating by parts, we find

$$\begin{aligned} \int_0^l (|v'_{0,\epsilon}|^2 - |v'_0|^2) dx &= \int_0^l (v'_{0,\epsilon} - v'_0)(v'_{0,\epsilon} + v'_0) dx \\ &= \int_0^l (u_{0,\epsilon} - u_0)(v_{0,\epsilon} + v_0) dx \\ &= O\left(\epsilon \log\left(\frac{\epsilon}{l}\right)\right) \|v_{0,\epsilon} + v_0\|_\infty \\ &= O\left(\epsilon l^2 \log\left(\frac{\epsilon}{l}\right)\right). \end{aligned}$$

On the interval $(0, 1)$,

$$\int_0^1 (|v'_{0,\epsilon}|^2 - |v'_0|^2) dx = O\left(\epsilon l \log\left(\frac{\epsilon}{l}\right)\right) = O\left(\frac{\epsilon}{N} \log(\epsilon N)\right).$$

$\int_0^1 \frac{1}{2} |v'_0|^2 dx$ can be evaluated (using (3.4), or see formulae (3.7) and (3.8) of [12]) and it turns out

$$\int_0^1 \frac{1}{2} |v'_0|^2 dx = \frac{(1 - m^2)^2}{24N^2}.$$

Then the nonlocal part of $I_\epsilon(u_{0,\epsilon})$ is bounded by $\frac{(1 - m^2)^2}{24N^2} + O\left(\frac{\epsilon}{N} \log(\epsilon N)\right)$. Combining this with (3.2), we obtain the first part of the lemma.

This estimate hints that the number of α -points of u_ϵ is of order $\epsilon^{-1/3}$. When N is taken to be the integer closest to $\left(\frac{(1 - m^2)^2}{12\epsilon c_0}\right)^{1/3}$, the optimal integer that minimize the right side of Lemma 3.1₁, we derive 2. \square

4 Some implications of the upper bound

Proposition 4.1 1. $\|v_\epsilon\|_\infty = O(\epsilon^{1/3})$,

2. $\lambda_\epsilon = O(\epsilon^{1/3})$,

$$3. \quad -1 + O(\epsilon^{1/3}) \leq u_\epsilon \leq 1 + O(\epsilon^{1/3}).$$

Proof. Lemma 3.1₂ implies $\int_0^1 |v'_\epsilon|^2 \leq C\epsilon^{2/3}$. And since $\int_0^1 v_\epsilon = 0$, we find $\|v_\epsilon\|_\infty = O(\epsilon^{1/3})$. Also by the same lemma $\int_0^1 W(u_\epsilon) \leq C\epsilon^{2/3}$. Integrating (1.5₁) we find

$$\begin{aligned} |\lambda_\epsilon| &= \left| \int_0^1 f(u_\epsilon) dx \right| \leq \int_0^1 |f(u_\epsilon)| dx \leq C \int_0^1 W^{1/2}(u_\epsilon) dx \\ &\leq C \left(\int_0^1 W(u_\epsilon) dx \right)^{1/2} \leq C\epsilon^{1/3}. \end{aligned}$$

The equation (1.5₁) yields $-\epsilon^2 u''_\epsilon + f(u_\epsilon) = O(\epsilon^{1/3})$. Let x_ϵ be a global maximum of u_ϵ . Then $u''_\epsilon(x_\epsilon) \leq 0$, whether or not x_ϵ is on the boundary, since $u'_\epsilon(0) = u'_\epsilon(1) = 0$. So $f(u_\epsilon(x_\epsilon)) \leq O(\epsilon^{1/3})$, which implies $u_\epsilon(x_\epsilon) \leq 1 + O(\epsilon^{1/3})$. The lower bound for u_ϵ follows after a similar argument. \square

It is often necessary to inspect u_ϵ in a scale comparable to ϵ . Let $x_\epsilon \in (0, 1)$ be an arbitrary point. Introduce t and U_ϵ so that $\epsilon t + x_\epsilon = x$ and $U_\epsilon(t) = u_\epsilon(x)$. According to Proposition 4.1_{1,2}, U_ϵ satisfies

$$-U''_\epsilon + f(U_\epsilon) = O(\epsilon^{1/3}) \tag{4.1}$$

on the expanding interval $(-\frac{x_\epsilon}{\epsilon}, \frac{1-x_\epsilon}{\epsilon})$. Since Proposition 4.1₃ implies $|U_\epsilon| \leq 1 + O(\epsilon^{1/3})$, the regularity theory of second order differential equations asserts that along any sequence U_{ϵ_n} of U_ϵ with $\epsilon_n \rightarrow 0$ there exists a subsequence that converges locally (at least) in C^1 to a function G which satisfies

$$-G'' + f(G) = 0, \quad -1 \leq G \leq 1, \tag{4.2}$$

on the whole interval $(-\infty, \infty)$, or a half interval (a, ∞) or $(-\infty, b)$.

Observing the phase portrait of this equation, we conclude that G must be A: a heteroclinic solution, i.e. a translate or a reversed translate of H defined in (2.3), B: the constant solution -1 or the constant solution 1 , C: the constant solution ω (the local maximum of W between -1 and 1), or D: a periodic solution whose trajectory is bounded by the two heteroclinic orbits in the phase portrait.

Lemma 4.2 *Case C or case D does not occur.*

Proof. We prove this by contradiction. Suppose that G is the non-stable constant ω or a periodic solution. We will construct a function whose energy is lower than that of u_ϵ , hence contradicting the fact that u_ϵ is a minimizer. To make notations manageable, any sequence or further subsequences of u_ϵ will still be denoted by u_ϵ instead of u_{ϵ_n} .

Take $\theta > 3$ to be a large number to be determined later. Always make θ be a integer multiple of the period of G if G is periodic. Without the loss of generality we assume $\limsup x_\epsilon \leq 1/2$. Let ξ be a smooth function defined on

$(-\infty, \infty)$ so that $\xi(t) = 0$ if $t \leq 0$, $\xi(t) = 1$ if $t \geq 1$, $|\xi(t)| \leq 1$ for all t . For each $r \in (1, \theta - 2)$ define

$$U_{\epsilon, r}(t) = \begin{cases} U_{\epsilon}(t), & t \notin (0, \theta) \\ (U_{\epsilon}(t) + 1)(1 - \xi(t)) - 1, & 0 \leq t \leq r \\ 2\xi(t - r) - 1, & r \leq t \leq r + 1 \\ (U_{\epsilon}(t) - 1)\xi(t - \theta + 1) + 1, & r + 1 \leq t \leq \theta \end{cases}. \quad (4.3)$$

We have replaced U_{ϵ} in the interval $(0, \theta)$ by a function which is -1 on $(1, r)$ and 1 on $(r + 1, \theta - 1)$. Similarly set

$$F_r(t) = \begin{cases} G(t), & t \notin (0, \theta) \\ (G(t) + 1)(1 - \xi(t)) - 1, & 0 \leq t \leq r \\ 2\xi(t - r) - 1, & r \leq t \leq r + 1 \\ (G(t) - 1)\xi(t - \theta + 1) + 1, & r + 1 \leq t \leq \theta \end{cases}, \quad (4.4)$$

Since $U_{\epsilon} \rightarrow G$ in $C^1[0, \theta]$, $U_{\epsilon, r} \rightarrow F_r$ in $C^1[0, \theta]$. We need to choose r properly to have $\int_0^{\theta} U_{\epsilon, t} = \int_0^{\theta} F_r$, so later the function that we will construct to have lower energy will be in the admissible set X_m .

Since θ is taken to be a multiple of the period of G if G is periodic, we see that $\frac{1}{\theta} \int_0^{\theta} G(t) dt \in (-1, 1)$ is independent of θ . Take $\eta > 0$ to be small so that $\frac{1}{\theta} \int_0^{\theta} G(t) dt \pm \eta \in (-1, 1)$. First set

$$r = r_1 = \frac{1 - \frac{1}{\theta} \int_0^{\theta} G(t) dt + \eta \theta}{2}.$$

Clearly $r_1 \in (1, \theta - 2)$ when θ is large. As $\theta \rightarrow \infty$, by the definition of F_r , (4.4),

$$\frac{1}{\theta} \int_0^{\theta} F_{r_1}(t) dt \rightarrow \frac{1}{\theta} \int_0^{\theta} G(t) dt - \eta.$$

Then set

$$r = r_2 = \frac{1 - \frac{1}{\theta} \int_0^{\theta} G(t) dt - \eta \theta}{2},$$

which is also in $(1, \theta - 2)$ when θ is large. As $\theta \rightarrow \infty$,

$$\frac{1}{\theta} \int_0^{\theta} F_{r_2}(t) dt \rightarrow \frac{1}{\theta} \int_0^{\theta} G(t) dt + \eta.$$

Therefore if we choose θ large enough

$$\frac{1}{\theta} \int_0^{\theta} F_{r_1}(t) dt < \frac{1}{\theta} \int_0^{\theta} G(t) dt < \frac{1}{\theta} \int_0^{\theta} F_{r_2}(t) dt.$$

After this large θ is chosen, we take small ϵ so that

$$\frac{1}{\theta} \int_0^{\theta} U_{\epsilon, r_1}(t) dt < \frac{1}{\theta} \int_0^{\theta} U_{\epsilon}(t) dt < \frac{1}{\theta} \int_0^{\theta} U_{\epsilon, r_2}(t) dt.$$

With both θ and ϵ chosen we set $r \in (r_1, r_2)$ so that $\int_0^\theta U_{\epsilon,r}(t) dt = \int_0^\theta U_\epsilon(t) dt$.

Back to the x -coordinate, we define $u_{\epsilon,r}(x) = U_{\epsilon,r}(t)$ which is in the admissible set X_m . We now proceed to compare the energy of u_ϵ and $u_{\epsilon,r}$, starting with the local part. As $\epsilon \searrow 0$,

$$\int_0^\theta |U'_{\epsilon,r}|^2 dt \rightarrow \int_0^\theta |F'_r|^2 dt = \int_0^1 |F'_r|^2 dt + \int_r^{r+1} |F'_r|^2 dt + \int_{\theta-1}^\theta |F'_r|^2 dt$$

which is bounded from above by a number independent of θ and r . The same is true for

$$\int_0^\theta W(U_{\epsilon,r}) dt \rightarrow \int_0^\theta W(F_r) dt.$$

So there exists $C > 0$ independent of θ and r such that

$$\int_0^\theta \left[\frac{|F'_r|^2}{2} + W(F_r) \right] dt \leq C.$$

Then for small ϵ

$$\int_{x_\epsilon}^{x_\epsilon + \epsilon\theta} \left[\frac{\epsilon^2}{2} |u'_{\epsilon,r}|^2 + W(u_{\epsilon,r}) \right] dx = \epsilon \int_0^\theta \left[\frac{|U'_{\epsilon,r}|^2}{2} + W(U_{\epsilon,r}) \right] dt \leq 2\epsilon C. \quad (4.5)$$

On the other hand since G , periodic or unstable constant, lies strictly away from -1 and 1 , there exists $c > 0$, independent of θ , such that $\int_0^\theta W(G(t)) dt \geq c\theta$. Therefore

$$\int_0^\theta \left[\frac{|G'|^2}{2} + W(G) \right] dt \geq c\theta.$$

Then for small ϵ

$$\int_{x_\epsilon}^{x_\epsilon + \epsilon\theta} \left[\frac{\epsilon^2}{2} |u'_\epsilon|^2 + W(u_\epsilon) \right] dx = \epsilon \int_0^\theta \left[\frac{|U'_\epsilon|^2}{2} + W(U_\epsilon) \right] dt \geq \frac{\epsilon\theta c}{2}. \quad (4.6)$$

We see that the local energy is reduced if θ is large.

To compare the nonlocal energy we work with the x -coordinate. Define $v_{\epsilon,r} = (-D^2)^{-1}(u_{\epsilon,r} - m)$. $v'_{\epsilon,r}$ and v'_ϵ agree outside $(x_\epsilon, x_\epsilon + \epsilon\theta)$. Clearly $v'_{\epsilon,r} = O(1)$ and $v'_\epsilon = O(1)$ because $u_{\epsilon,r}$ and u_ϵ are of order $O(1)$. Since $-(v'_{\epsilon,r} - v'_\epsilon)' = u_{\epsilon,r} - u_\epsilon$ on $(x_\epsilon, x_\epsilon + \epsilon\theta)$, $v'_{\epsilon,r} - v'_\epsilon = O(\epsilon\theta)$ there. Then

$$\begin{aligned} \int_{x_\epsilon}^{x_\epsilon + \epsilon\theta} |v'_{\epsilon,r}|^2 dx - \int_{x_\epsilon}^{x_\epsilon + \epsilon\theta} |v'_\epsilon|^2 dx &= \int_{x_\epsilon}^{x_\epsilon + \epsilon\theta} (v'_{\epsilon,r} - v'_\epsilon)(v'_{\epsilon,r} + v'_\epsilon) dx \\ &= \int_{x_\epsilon}^{x_\epsilon + \epsilon\theta} O(\epsilon\theta) \cdot O(1) dx = O(\epsilon^2\theta^2). \end{aligned}$$

Combining this with (4.5) and (4.6) we deduce

$$I_\epsilon(u_{\epsilon,r}) - I_\epsilon(u_\epsilon) \leq 2\epsilon C - \frac{\epsilon\theta c}{2} + O(\epsilon^2\theta^2).$$

Just like in the construction of $U_{\epsilon,r}$, we first choose θ large then ϵ small, so $I_\epsilon(u_{\epsilon,r}) < I_\epsilon(u_\epsilon)$. \square

We first use this lemma to study α -points of u_ϵ . Recall from Section 2 that x is an α -point if $u_\epsilon(x) = \alpha$.

Proposition 4.3 *When ϵ is small, $u'_\epsilon(x_\epsilon) \neq 0$ at every α -point x_ϵ .*

Proof. From Lemma 4.2 $u_\epsilon(\epsilon t + x_\epsilon) \rightarrow G$ locally in C^1 , where G is heteroclinic or ± 1 . Since $u_\epsilon(x_\epsilon) = \alpha$, $G(0) = \alpha$. Then $G(t) = H(t)$ or $G(t) = H(-t)$ (H is defined in (2.3)). Then $\epsilon u'_\epsilon(x_\epsilon) \rightarrow \pm H'(0) \neq 0$. \square

The proof actually says more: $u'_\epsilon(x_\epsilon) \rightarrow \pm\infty$. Proposition 4.3 implies that the α -points of u_ϵ are *non-degenerate*, meaning that every time the graph of u_ϵ touches the horizontal level α , it crosses it. The next application of Lemma 4.2 shows that α -points do not appear in any neighborhood of the boundary of $(0, 1)$ whose size is of order ϵ .

Proposition 4.4 *If x_ϵ is an α -point of u_ϵ , $\frac{\epsilon}{x_\epsilon} = o(1)$ and $\frac{\epsilon}{1-x_\epsilon} = o(1)$.*

Proof. Of course one of $\frac{\epsilon}{x_\epsilon} = o(1)$ and $\frac{\epsilon}{1-x_\epsilon} = o(1)$ must hold. Suppose the former is true and the latter is false. Then we can assume $\frac{1-x_\epsilon}{\epsilon} \rightarrow b \geq 0$. Let $U_\epsilon(t) = u_\epsilon(\epsilon t + x_\epsilon)$. Again by Lemma 4.2 $U_\epsilon(t)$ converges to $H(t)$ or $H(-t)$ locally in C^1 . However $0 = U'_\epsilon(\frac{1-x_\epsilon}{\epsilon}) \rightarrow \pm H'(b) \neq 0$. A contradiction. \square

These two propositions imply that the number of α -points is finite for each small ϵ . Let us denote them by $x_1, x_2, \dots, x_{N_\epsilon}$, in increasing order. We suppress the dependence of the x_i 's on ϵ to simplify notation. Throughout the rest of the paper we assume without the loss of generality that $u_\epsilon > \alpha$ on $(0, x_1)$ and N_ϵ is even. We set $M_\epsilon = N_\epsilon/2$. Let

$$\begin{cases} p_1 = x_1, & p_2 = x_3 - x_2, & \dots & p_{M_\epsilon+1} = 1 - x_{N_\epsilon} \\ q_1 = x_2 - x_1, & q_2 = x_4 - x_3, & \dots, & q_{M_\epsilon} = x_{N_\epsilon} - x_{N_\epsilon-1} \end{cases}. \quad (4.7)$$

When no confusion exists we call the interval whose length is p_i the interval p_i and the interval whose length is q_i the interval q_i . Because of the non-degeneracy of the x_i 's, $u_\epsilon > \alpha$ on every interval p_i and $u_\epsilon < \alpha$ on every interval q_i . The last interval $(x_{N_\epsilon}, 1)$ is $p_{M_\epsilon+1}$. Again the p_i 's and q_i 's depend on ϵ . With this setting α -point x_{2i-2} is followed by the interval p_i , which is followed by x_{2i-1} , which is followed by the interval q_i .

Proposition 4.5 *$\frac{\epsilon}{p_i} = o(1)$ and $\frac{\epsilon}{q_i} = o(1)$.*

Proof. The cases of p_1 and $p_{M_\epsilon+1}$ are already covered by Proposition 4.4. Suppose this proposition is false. There exist adjacent α -points x_ϵ and x_ϵ^* such that $\frac{x_\epsilon^* - x_\epsilon}{\epsilon} \rightarrow d \geq 0$. Again the convergence is really along a sequence ϵ_n of ϵ , but we stay with ϵ . Without the loss of generality we assume $u_\epsilon > \alpha$ on $(x_\epsilon, x_\epsilon^*)$. Let $U_\epsilon(t) = u_\epsilon(\epsilon t + x_\epsilon)$.

If $d = 0$, then there exists $t_\epsilon \in (0, \frac{x_\epsilon^* - x_\epsilon}{\epsilon})$ such that $U'_\epsilon(t_\epsilon) = 0$. As $\epsilon \searrow 0$, we have $\frac{x_\epsilon^* - x_\epsilon}{\epsilon} \rightarrow 0$ and $t_\epsilon \rightarrow 0$. Also by Lemma 4.2 and the facts that $U_\epsilon(0) = \alpha$ and $U_\epsilon > \alpha$ on $(0, \frac{x_\epsilon^* - x_\epsilon}{\epsilon})$, $U_\epsilon(t) \rightarrow H(t)$ locally in C^1 . Then $0 = U'_\epsilon(t_\epsilon) \rightarrow H'(0) \neq 0$. A contradiction.

If $d > 0$, then again $U_\epsilon(t) \rightarrow H(t)$. So $\alpha = U_\epsilon(\frac{x_\epsilon^* - x_\epsilon}{\epsilon}) \rightarrow H(d)$. But $H(d) = \alpha$ is impossible since $H(0) = \alpha$ and H is strictly increasing. \square

Lemma 4.6 1. For $i = 2, 3, \dots, M_\epsilon + 1$,

$$\begin{aligned} \|u_\epsilon(\epsilon t + x_{2i-2}) - H(t)\|_{C^2[0, p_i/(2\epsilon)]} &= O(\epsilon^{1/3}) + O(e^{-\nu p_i/(2\epsilon)}), \\ \|u_\epsilon(\epsilon t + x_{2i-2}) - H(t)\|_{C^2[-q_{i-1}/(2\epsilon), 0]} &= O(\epsilon^{1/3}) + O(e^{-\nu q_{i-1}/(2\epsilon)}). \end{aligned}$$

2. For $i = 1, 2, \dots, M_\epsilon$,

$$\begin{aligned} \|u_\epsilon(\epsilon t + x_{2i-1}) - H(-t)\|_{C^2[0, q_i/(2\epsilon)]} &= O(\epsilon^{1/3}) + O(e^{-\nu q_i/(2\epsilon)}), \\ \|u_\epsilon(\epsilon t + x_{2i-1}) - H(-t)\|_{C^2[-p_i/(2\epsilon), 0]} &= O(\epsilon^{1/3}) + O(e^{-\nu p_i/(2\epsilon)}). \end{aligned}$$

In this lemma if an estimate is on the end interval p_1 or $p_{M_\epsilon+1}$, then the (2ϵ) 's on both sides of the estimate should read ϵ .

Proof. We only prove the first estimate of Lemma 4.6₁, since the other three are similar. There are two different cases. When $i = 1, 2, \dots, M_\epsilon$, u_ϵ is estimated on a p_i interval with two α -points x_{2i-2} and x_{2i-1} as the boundary. When $i = M_\epsilon + 1$, u_ϵ is estimated on $(x_{2M_\epsilon}, 1)$, an end interval. In order to study the two cases in a unified way, in this proof we extend the domain of u_ϵ and v_ϵ to $(0, 1 + p_{M_\epsilon+1})$ by setting $u_\epsilon(x) = u_\epsilon(2 - x)$ and $v_\epsilon(x) = v_\epsilon(2 - x)$ for $x \in (1, 1 + p_{M_\epsilon+1})$. u_ϵ and v_ϵ still solve (1.5) on $(0, 1 + p_{M_\epsilon+1})$ and $u_\epsilon(1 + p_{M_\epsilon+1}) = \alpha$. Let $x = \epsilon t + x_{2i-2}$ and $U_\epsilon(t) = u_\epsilon(\epsilon t + x_{2i-2})$. The proof consists of four steps.

Step 1: $\|U_\epsilon - H\|_{L^\infty(0, p_i/(2\epsilon))} = o(1)$. As $\epsilon \searrow 0$ by Proposition 4.5 $p_i/(2\epsilon) \rightarrow \infty$, and by Lemma 4.2 $U_\epsilon \rightarrow H$ locally in C^1 . If this convergence is not in $L^\infty(0, p_i/(2\epsilon))$, there exists $h_\epsilon \in (0, p_i/(2\epsilon))$ such that $|U_\epsilon(h_\epsilon) - H(h_\epsilon)| \not\rightarrow 0$ and $h_\epsilon \rightarrow \infty$. Thus $U_\epsilon(h_\epsilon)$ stays away from 1. Shift $U_\epsilon(t)$ to $U_\epsilon(t + h_\epsilon)$. Let G be such that $U_\epsilon(t + h_\epsilon) \rightarrow G$ locally in C^1 and $-G'' + f(G) = 0$. Then G is either 1 or heteroclinic by Lemma 4.2. If $G = 1$, $U_\epsilon(h_\epsilon) \rightarrow 1$. A contradiction. If G is heteroclinic, $G(\zeta) < \alpha$ at some ζ . Then $U_\epsilon(\zeta + h_\epsilon) < \alpha$ when ϵ is small. This is impossible since when $t = \zeta + h_\epsilon$, $x = \epsilon(\zeta + h_\epsilon) + x_{2i-2} \in (x_{2i-2}, x_{2i-1})$ where $u_\epsilon > \alpha$.

Step 2: $\|U_\epsilon - H\|_{L^\infty(0, p_i/(2\epsilon))} = O(\epsilon^{1/3}) + O(e^{-\nu p_i/(2\epsilon)})$. Let $G_{p_i/(2\epsilon)}$ be the increasing solution of $-G'' + f(G) = 0$ with the boundary conditions $G_{p_i/(2\epsilon)}(0) = \alpha$ and $G'_{p_i/(2\epsilon)}(p_i/\epsilon) = 0$. Note that $G_{p_i/(2\epsilon)}$ is part of a periodic trajectory in the phase plane and $G_{p_i/(2\epsilon)}(p_i/\epsilon) = \alpha$. We first show that $\|U_\epsilon - G_{p_i/(2\epsilon)}\|_{L^\infty(0, p_i/\epsilon)} = O(\epsilon^{1/3})$.

On the contrary suppose that $\|U_\epsilon - G_{p_i/(2\epsilon)}\|_{L^\infty(0, p_i/\epsilon)} \epsilon^{-1/3} \rightarrow \infty$. Let

$$\Psi_\epsilon = \frac{U_\epsilon - G_{p_i/(2\epsilon)}}{\|U_\epsilon - G_{p_i/(2\epsilon)}\|_{L^\infty(0, p_i/\epsilon)}}.$$

By Proposition 4.1_{1,2}, U_ϵ satisfies $-U_\epsilon'' + f(U_\epsilon) = O(\epsilon^{1/3})$. So Ψ_ϵ satisfies $-\Psi_\epsilon'' + f'(\dots)\Psi_\epsilon = o(1)$, $\Psi_\epsilon(0) = \Psi_\epsilon(p_i/\epsilon) = 0$, where f' is evaluated at a number between U_ϵ and $G_{p_i/(2\epsilon)}$, whose exact value is not important for us. Without the loss of generality we assume that the maximum of $|\Psi_\epsilon|$ is achieved at $h_\epsilon \in [0, p_i/\epsilon]$ which is a global maximum, i.e. $\Psi_\epsilon(h_\epsilon) = 1$. There are three possibilities for the location of h_ϵ . A: there exist $\eta > 0$ such that $h_\epsilon < \eta$ for all ϵ . B: there exists $\eta > 0$ such that $h_\epsilon > p_i/\epsilon - \eta$ for all ϵ . C: Neither of the above.

If case A occurs, by the fact that $G_{p_i/(2\epsilon)} \rightarrow H$ in $L^\infty(0, p_i/(2\epsilon))$ as $\epsilon \rightarrow 0$, Lemma 2.1₂, and Step 1, we find $\Psi_\epsilon \rightarrow \Psi$ locally in C^1 where Ψ satisfies $-\Psi'' + f'(H)\Psi = 0$ on $(0, \infty)$. Since $|\Psi| \leq 1$ Lemma 2.2₁ asserts $\Psi = cH'$ for some c . Also $\Psi(0) = \lim \Psi_\epsilon(0) = 0$. Since $H'(0) \neq 0$, $c = 0$ and $\Psi = 0$. This is clearly inconsistent with $\Psi_\epsilon(h_\epsilon) = 1$ and $h_\epsilon < \eta$ for all small ϵ .

Case B can be ruled out in the same manner. When case C occurs we assume $h_\epsilon \in (0, p_i/\epsilon)$, $h_\epsilon \rightarrow \infty$ and $p_i/\epsilon - h_\epsilon \rightarrow \infty$. By Step 1, or a similar assertion $\|U_\epsilon - H(p_i/\epsilon - \cdot)\|_{L^\infty(p_i/(2\epsilon), p_i/\epsilon)} = o(1)$, we find that in the equation for Ψ_ϵ , $-\Psi_\epsilon''(h_\epsilon) \geq 0$ (since h_ϵ is a maximum) and $f'(\dots)\Psi_\epsilon(h_\epsilon) \rightarrow f'(1) > 0$. Thus the equation can not be satisfied at h_ϵ when ϵ is small. So we have proved that $\|U_\epsilon - G_{p_i/(2\epsilon)}\|_{L^\infty(0, p_i/\epsilon)} = O(\epsilon^{1/3})$. Lemma 2.1₂ then completes Step 2.

Step 3: $\|U_\epsilon'' - H''\|_{L^\infty(0, p_i/(2\epsilon))} = O(\epsilon^{1/3}) + O(e^{-\nu p_i/\epsilon})$. From Steps 1, 2 and the equations (4.1) and (2.3) that U_ϵ and H satisfy respectively,

$$(U_\epsilon'' - H'') = f'(\dots)(U_\epsilon - H) + O(\epsilon^{1/3}) = O(\epsilon^{1/3}) + O(e^{-\nu p_i/\epsilon}).$$

Step 4: $\|U_\epsilon' - H'\|_{L^\infty(0, p_i/(2\epsilon))} = O(\epsilon^{1/3}) + O(e^{-\nu p_i/\epsilon})$. Let $S_\epsilon = U_\epsilon - H$. Then $S = O(\epsilon^{1/3}) + O(e^{-\nu p_i/\epsilon})$ and $S'' = O(\epsilon^{1/3}) + O(e^{-\nu p_i/\epsilon})$ by Steps 2 and 3. Assume without the loss of generality $h, h+1 \in (0, p_i/(2\epsilon))$. (Otherwise consider $h, h-1$). Then

$$\begin{aligned} O(\epsilon^{1/3}) + O(e^{-\nu p_i/\epsilon}) &= S_\epsilon(h+1) = S_\epsilon(h) + S_\epsilon'(h) + \frac{1}{2}S_\epsilon''(\dots) \\ &= S_\epsilon'(h) + O(\epsilon^{1/3}) + O(e^{-\nu p_i/\epsilon}), \end{aligned}$$

Hence $S_\epsilon'(h) = O(\epsilon^{1/3}) + O(e^{-\nu p_i/\epsilon})$. \square

5 A lower bound of $I_\epsilon(u_\epsilon)$

A scaling in Lemma 2.3_{2,3,5} yields a lower bound for the local part of $I_\epsilon(u_\epsilon)$.

Lemma 5.1 *On a p_i or q_i interval the local part of $I_\epsilon(u_\epsilon)$ has the lower bound*

$$\begin{aligned} \int_{p_i} \left[\frac{\epsilon^2}{2} |u_\epsilon'|^2 + W(u_\epsilon) \right] dx &\geq \begin{cases} c_1\epsilon + p_i O(e^{-\mu p_i/\epsilon}) & i = 1, M_\epsilon + 1 \\ 2c_1\epsilon + p_i O(e^{-\mu p_i/\epsilon}) & i \neq 1, M_\epsilon + 1 \end{cases} \\ \int_{q_i} \left[\frac{\epsilon^2}{2} |u_\epsilon'|^2 + W(u_\epsilon) \right] dx &\geq 2c_{-1}\epsilon + q_i O(e^{-\mu q_i/\epsilon}). \end{aligned}$$

More difficult to find are the lower bounds for the nonlocal part of $I_\epsilon(u_\epsilon)$.

Lemma 5.2 *On a p_i or q_i interval the nonlocal part of $I_\epsilon(u_\epsilon)$ has the lower bound*

$$\begin{aligned}
\frac{1}{2} \int_{p_i} |v'_\epsilon|^2 &\geq \frac{(1-m)^2}{6} p_i^3 + p_i^3 O(\epsilon^{1/3}) + p_i^2 O(\epsilon), \quad i = 1, M_\epsilon + 1 \\
\frac{1}{2} \int_{p_i} |v'_\epsilon|^2 &\geq \frac{(1-m)^2}{24} p_i^3 + \frac{p_i}{2} [v'_\epsilon(x_{2i-2}) - \frac{(1-m)p_i}{2} + p_i O(\epsilon^{1/3}) + O(\epsilon)]^2 \\
&\quad + p_i^3 O(\epsilon^{1/3}) + p_i^2 O(\epsilon), \quad i \neq 1, M_\epsilon + 1 \\
\frac{1}{2} \int_{q_i} |v'_\epsilon|^2 &\geq \frac{(1+m)^2}{24} q_i^3 + \frac{q_i}{2} [v'_\epsilon(x_{2i-1}) + \frac{(1+m)q_i}{2} + q_i O(\epsilon^{1/3}) + O(\epsilon)]^2 \\
&\quad + q_i^3 O(\epsilon^{1/3}) + q_i^2 O(\epsilon), \quad i = 1, 2, \dots, M_\epsilon.
\end{aligned}$$

Proof. On $(0, x_1)$, with the help of Lemma 4.6, we have

$$\begin{aligned}
v'_\epsilon(x) &= v'_\epsilon(0) - \int_0^x (u_\epsilon - m) dy \\
&= - \int_0^x (1-m) dy + \int_0^x [H(\frac{p_1 - y}{\epsilon}) - u_\epsilon] dy + \int_0^x [1 - H(\frac{p_1 - y}{\epsilon})] dy \\
&= -(1-m)x + p_1 O(\epsilon^{1/3}) + p_1 O(e^{-\nu p_1/\epsilon}) + O(\epsilon) \\
&= -(1-m)x + p_1 O(\epsilon^{1/3}) + O(\epsilon),
\end{aligned}$$

where the reduction to the last line follows from the estimate $p_1 O(e^{-\nu p_1/\epsilon}) = \epsilon O(\frac{p_1}{\epsilon} e^{-\nu p_1/\epsilon}) = o(\epsilon)$. This leads to

$$\frac{1}{2} \int_{p_1} |v'_\epsilon|^2 dx = \frac{(1-m)^2 p_1^3}{6} + p_1^3 O(\epsilon^{1/3}) + p_1^2 O(\epsilon).$$

On interval (x_1, x_2)

$$\begin{aligned}
v'_\epsilon(x) &= v'_\epsilon(x_1) - \int_{x_1}^x (u_\epsilon - m) dy \\
&= v'_\epsilon(x_1) + (1+m)(x - x_1) + q_1 O(\epsilon^{1/3}) + O(\epsilon), \quad (5.1)
\end{aligned}$$

which implies

$$\begin{aligned}
&\frac{1}{2} \int_{q_1} |v'_\epsilon|^2 dx \\
&= \frac{1}{2} \int_{q_1} [v'_\epsilon(x_1) + (1+m)(x - x_1) + q_1 O(\epsilon^{1/3}) + O(\epsilon)]^2 dx \\
&= \frac{1}{2} \int_{q_1} [v'_\epsilon(x_1) + (1+m)(x - x_1)]^2 dx
\end{aligned}$$

$$\begin{aligned}
& +v'_\epsilon(x_1)[(q_1)^2O(\epsilon^{1/3}) + q_1O(\epsilon)] + (q_1)^3O(\epsilon^{1/3}) + (q_1)^2O(\epsilon) \\
= & \frac{[v'_\epsilon(x_1) + (1+m)(x-x_1)]^3}{6(1+m)} \Big|_{x=x_1}^{x=x_2} \\
& +v'_\epsilon(x_1)[(q_1)^2O(\epsilon^{1/3}) + q_1O(\epsilon)] + (q_1)^3O(\epsilon^{1/3}) + (q_1)^2O(\epsilon) \\
= & \frac{1}{6(1+m)} [2(\frac{(1+m)q_1}{2})^3 + 3(1+m)q_1(v'_\epsilon(x_1) + \frac{(1+m)q_1}{2})^2] \\
& +v'_\epsilon(x_1)[(q_1)^2O(\epsilon^{1/3}) + q_1O(\epsilon)] + (q_1)^3O(\epsilon^{1/3}) + (q_1)^2O(\epsilon) \\
= & \frac{(1+m)^2(q_1)^3}{24} + \frac{q_1}{2}[v'_\epsilon(x_1) + \frac{(1+m)q_1}{2} + q_1O(\epsilon^{1/3}) + O(\epsilon)]^2 \\
& + (q_1)^3O(\epsilon^{1/3}) + (q_1)^2O(\epsilon).
\end{aligned}$$

Continue this argument until we reach the q_{M_ϵ} interval $(x_{N_\epsilon-1}, x_{N_\epsilon})$. Finally on $(x_{N_\epsilon}, 1)$ we use an estimate similar to the one on $(0, x_1)$, i.e. write

$$v'_\epsilon(x) = v'_\epsilon(1) - \int_1^x (u_\epsilon - m) dy = (1-m)(1-x) + p_{M_\epsilon+1}O(\epsilon^{1/3}) + O(\epsilon),$$

to derive

$$\frac{1}{2} \int_{p_{M_\epsilon+1}} |v'_\epsilon|^2 dx = \frac{(1-m)^2 p_{M_\epsilon+1}^3}{6} + p_{M_\epsilon+1}^3 O(\epsilon^{1/3}) + p_{M_\epsilon+1}^2 O(\epsilon). \quad \square$$

Two remarks are in order. First the two square terms in the lemma involving $v'_\epsilon(x_{2i-1,2})$ will be only used once, though critically, in the proof of Proposition 7.2. In the other applications they will simply be dropped.

Second we have presented this lemma arguing first with $(0, x_1)$ and then proceeding to the right. As a consequence $v'_\epsilon(x_{N_\epsilon})$ does not appear in the estimates. Naturally there is another version of this lemma where we start with $(x_{N_\epsilon}, 1)$ and proceed backwards. Then the second and the third inequalities become

$$\begin{aligned}
\frac{1}{2} \int_{p_i} |v'_\epsilon|^2 & \geq \frac{(1-m)^2}{24} p_i^3 + \frac{p_i}{2} [v'_\epsilon(x_{2i+1}) + \frac{(1-m)p_i}{2} + p_i O(\epsilon^{1/3}) + O(\epsilon)]^2 \\
& + p_i^3 O(\epsilon^{1/3}) + p_i^2 O(\epsilon), \quad i \neq 1, M_\epsilon + 1 \tag{5.2} \\
\frac{1}{2} \int_{q_i} |v'_\epsilon|^2 & \geq \frac{(1+m)^2}{24} q_i^3 + \frac{q_i}{2} [v'_\epsilon(x_{2i}) - \frac{(1+m)q_i}{2} + q_i O(\epsilon^{1/3}) + O(\epsilon)]^2 \\
& + q_i^3 O(\epsilon^{1/3}) + q_i^2 O(\epsilon).
\end{aligned}$$

In this version $v'_\epsilon(x_1)$ does not appear.

Following Lemma 5.2 is a very rough upper bound for p_i and q_i .

Proposition 5.3 $p_i = O(\epsilon^{2/9})$ and $q_i = O(\epsilon^{2/9})$.

Proof. Let us consider the case of p_i , $i \neq 1, M_\epsilon + 1$. The other two cases can be handled similarly. According to Lemma 5.2

$$I_\epsilon(u_\epsilon) \geq \frac{(1-m)^2}{24} p_i^3 + p_i^3 O(\epsilon^{1/3}) + p_i^2 O(\epsilon).$$

Because of Proposition 4.5, the last two terms on the right side can be written as $p_i^3 o(1)$, which is small compared to the first term on the right side. Also because of the upper bound, Lemma 3.1₂, for $I_\epsilon(\epsilon)$, something of order $O(\epsilon^{2/3})$, we find that $p_i^3 = O(\epsilon^{2/3})$ and $q_i^3 = O(\epsilon^{2/3})$. \square

Sum over i in Lemmas 5.1 and 5.2 to obtain our first lower bound of $I_\epsilon(u_\epsilon)$.

Lemma 5.4

$$\begin{aligned} I_\epsilon(u_\epsilon) \geq & c_0 \epsilon N_\epsilon + \sum_{i=1}^{M_\epsilon+1} p_i O(e^{-\mu p_i/\epsilon}) + \sum_{i=1}^{M_\epsilon} q_i O(e^{-\mu q_i/\epsilon}) \\ & + \frac{(1-m)^2}{24} [4p_1^3 + \sum_{i=2}^{M_\epsilon} p_i^3 + 4p_{M_\epsilon+1}^3] + \frac{(1+m)^2}{24} \sum_{i=1}^{M_\epsilon} q_i^3 \\ & + [\sum_{i=1}^{M_\epsilon+1} p_i^3 + \sum_{i=1}^{M_\epsilon} q_i^3] O(\epsilon^{1/3}) + [\sum_{i=1}^{M_\epsilon+1} p_i^2 + \sum_{i=1}^{M_\epsilon} q_i^2] O(\epsilon). \end{aligned}$$

An important consequence of Lemma 5.4 is that $1/N_\epsilon \sim \epsilon^{1/3}$. We need a simple technical lemma first.

Lemma 5.5 1. In the set $\{(p_1, p_2, p_3, \dots, p_{M+1}) : p_i > 0, i = 1, 2, \dots, M+1, p_1 + p_2 + \dots + p_{M+1} = d, d > 0\}$, $4p_1^3 + \sum_{i=2}^M p_i^3 + 4p_{M+1}^3$ is minimized when $2p_1, p_2, \dots, p_M$, and $2p_{M+1}$ are all equal to $p = \frac{d}{M}$. Moreover

$$\begin{aligned} 4p_1^3 + \sum_{i=2}^M p_i^3 + 4p_{M+1}^3 & \geq 4\left(\frac{p}{2}\right)^3 + \sum_{i=2}^M p^3 + 4\left(\frac{p}{2}\right)^3 \\ & + 4p\left(p_1 - \frac{p}{2}\right)^2 + \sum_{i=2}^M 2p(p_i - p)^2 + 4p\left(p_{M+1} - \frac{p}{2}\right)^2. \end{aligned}$$

2. In the set $\{(q_1, q_2, q_3, \dots, q_M) : q_i > 0, i = 1, 2, \dots, M, q_1 + q_2 + \dots + q_M = d, d > 0\}$, $\sum_{i=1}^M q_i^3$ is minimized when q_1, q_2, \dots, q_M are all equal to $q = \frac{d}{M}$. Moreover

$$\sum_{i=1}^M q_i^3 \geq \sum_{i=1}^M q^3 + \sum_{i=1}^M 2q(q_i - q)^2.$$

Proof. We only treat case 1. Note

$$\begin{aligned} p_i^3 & = p^3 + 3p^2(p_i - p) + 2p(p_i - p)^2 + p_i(p_i - p)^2 \\ & \geq p^3 + 3p^2(p_i - p) + 2p(p_i - p)^2. \end{aligned}$$

when $i \neq 1$ or $M+1$. And when $i = 1$ or $M+1$,

$$\begin{aligned} 4p_1^3 & = 4\left(\frac{p}{2}\right)^3 + 3p^2\left(p_1 - \frac{p}{2}\right) + 4p\left(p_1 - \frac{p}{2}\right)^2 + 4p_1\left(p_1 - \frac{p}{2}\right)^2 \\ & \geq 4\left(\frac{p}{2}\right)^3 + 3p^2\left(p_1 - \frac{p}{2}\right) + 4p\left(p_1 - \frac{p}{2}\right)^2 \end{aligned}$$

The lemma then follows after we sum over i . \square

We also need the facts that

$$\sum_{i=1}^{M_\epsilon+1} p_i = \frac{1+m}{2} + O(\epsilon^{1/3} + \epsilon N_\epsilon), \quad \sum_{i=1}^{M_\epsilon} q_i = \frac{1-m}{2} + O(\epsilon^{1/3} + \epsilon N_\epsilon) \quad (5.3)$$

To see (5.3) we note that

$$m = \int_0^1 u_\epsilon dx = \sum_{i=1}^{M_\epsilon+1} \int_{p_i} u_\epsilon dx + \sum_{i=1}^{M_\epsilon} \int_{q_i} u_\epsilon dx$$

Every p_i or q_i interval is further divided in the middle, except the end intervals. Then, for example, with $U_\epsilon(t) = u_\epsilon(\epsilon t + x_{2i-2})$,

$$\begin{aligned} \int_{x_{2i-2}}^{x_{2i-2}+p_i/2} u_\epsilon dx &= \epsilon \int_0^{p_i/(2\epsilon)} U_\epsilon dt \\ &= \epsilon \int_0^{p_i/(2\epsilon)} (U_\epsilon - H) dt + \epsilon \int_0^{p_i/(2\epsilon)} (H - 1) dt + \frac{p_i}{2}. \end{aligned}$$

The first term of the last line is of order

$$p_i O(\epsilon^{1/3}) + p_i O(e^{-\nu p_i/\epsilon}) = p_i O(\epsilon^{1/3}) + \epsilon O\left(\frac{p_i}{\epsilon} e^{-\nu p_i/\epsilon}\right) = p_i O(\epsilon^{1/3}) + o(\epsilon)$$

by Lemma 4.6. The second term is of order $O(\epsilon)$, because $|H - 1|$ is integrable on $(0, \infty)$. Summing over all the p_i and q_i intervals, we deduce

$$\sum_{i=1}^{M_\epsilon+1} p_i - \sum_{i=1}^{M_\epsilon} q_i = m + O(\epsilon^{1/3}) + O(\epsilon N_\epsilon).$$

On the other hand

$$\sum_{i=1}^{M_\epsilon+1} p_i + \sum_{i=1}^{M_\epsilon} q_i = 1.$$

(5.3) follows after we solve these two equations.

Proposition 5.6 $\frac{1}{N_\epsilon} \sim \epsilon^{1/3}$.

Proof. We only need a weaker version of Lemma 5.4. Note that

$$\sum_{i=1}^{M_\epsilon} p_i O(e^{-\mu p_i/\epsilon}) = \epsilon \sum_{i=1}^{M_\epsilon} \frac{p_i}{\epsilon} O(e^{-\mu p_i/\epsilon}) = N_\epsilon o(\epsilon),$$

since $\frac{p_i}{\epsilon} O(e^{-\mu p_i/\epsilon}) = o(1)$. By Proposition 4.5

$$p_i^3 O(\epsilon^{1/3}) + p_i^2 O(\epsilon) = p_i^3 (O(\epsilon^{1/3}) + \frac{\epsilon}{p_i} O(1)) = p_i^3 o(1).$$

Then by Lemma 5.4

$$\begin{aligned}
I_\epsilon(u_\epsilon) &\geq c_0\epsilon N_\epsilon + N_\epsilon o(\epsilon) + \frac{(1-m)^2}{24} [4p_1^3 + \sum_{i=2}^{M_\epsilon} p_i^3 + 4p_{M_\epsilon+1}^3] \\
&\quad + \frac{(1+m)^2}{24} \sum_{i=1}^{M_\epsilon} q_i^3 + [\sum_{i=1}^{M_\epsilon+1} p_i^3 + \sum_{i=1}^{M_\epsilon} q_i^3] o(1) \\
&= c_0\epsilon N_\epsilon + N_\epsilon o(\epsilon) + (\frac{(1-m)^2}{24} + o(1)) [4p_1^3 + \sum_{i=1}^{M_\epsilon} p_i^3 + 4p_{M_\epsilon+1}^3] \\
&\quad + (\frac{(1+m)^2}{24} + o(1)) \sum_{i=1}^{M_\epsilon} q_i^3. \tag{5.4}
\end{aligned}$$

According to Lemma 5.5 and (5.3) $4p_1^3 + (p_2)^3 + \dots + 4p_{M_\epsilon+1}^3$ achieves minimum if all $2p_1, p_2, \dots, 2p_{M_\epsilon+1}$ happen to be

$$p = \frac{1}{M_\epsilon} \sum_{i=1}^{M_\epsilon+1} p_i = \frac{\frac{1+m}{2} + O(\epsilon^{1/3} + \epsilon N_\epsilon)}{M_\epsilon} = \frac{\frac{1+m}{2} + o(1)}{M_\epsilon}. \tag{5.5}$$

Therefore

$$4p_1^3 + \sum_{i=2}^{M_\epsilon} p_i^3 + 4p_{M_\epsilon+1}^3 \geq M_\epsilon^{-2} (\frac{1+m}{2} + o(1))^3. \tag{5.6}$$

After applying the same argument to q_i , we deduce from (5.4)

$$\begin{aligned}
I_\epsilon(u_\epsilon) &\geq c_0\epsilon N_\epsilon + N_\epsilon o(\epsilon) + \frac{(1-m)^2}{24} (\frac{1+m}{2})^3 M_\epsilon^{-2} \\
&\quad + \frac{(1+m)^2}{24} (\frac{1-m}{2})^3 M_\epsilon^{-2} + M_\epsilon^{-2} o(1) \\
&= c_0\epsilon N_\epsilon + N_\epsilon o(\epsilon) + \frac{(1-m^2)^2}{24N_\epsilon^2} + N_\epsilon^{-2} o(1). \tag{5.7}
\end{aligned}$$

Recall the upper bound, Lemma 3.1₂, for $I_\epsilon(u_\epsilon)$. We find

$$c_0\epsilon N_\epsilon + N_\epsilon o(\epsilon) + \frac{(1-m^2)^2}{24N_\epsilon^2} + N_\epsilon^{-2} o(1) = O(\epsilon^{2/3}).$$

Therefore

$$N_\epsilon = O(\epsilon^{-1/3}), \quad N_\epsilon^{-2} = O(\epsilon^{2/3}),$$

which completes the proof. \square

6 The first estimation of p_i and q_i

The crude lower and upper bounds for p_i and q_i in Propositions 4.5 and 5.3 are improved in this and the next sections. The upper bound is lowered to $O(\epsilon^{1/3})$

first. To prove this we have to treat *long* p_i and q_i intervals and possible *short* p_i and q_i intervals differently. Let c_2 be a positive number large enough so that when $p_i \geq -c_2\epsilon \log \epsilon$,

$$p_i e^{-\mu p_i/\epsilon} = O(\epsilon^{13/9}), \quad e^{-\nu p_i/\epsilon} = o(\epsilon^{1/3}). \quad (6.1)$$

When p_i (or q_i) is not an end interval, we say p_i (or q_i) is long if $p_i \geq -c_2\epsilon \log \epsilon$ (or $q_i \geq -c_2\epsilon \log \epsilon$). When p_i (or q_i) is an end interval, we say p_i (or q_i) is long if $p_i \geq -\frac{c_2}{2}\epsilon \log \epsilon$ (or $q_i \geq -\frac{c_2}{2}\epsilon \log \epsilon$). Otherwise we say p_i (or q_i) is short. Let P_L and P_S be the numbers of long and short p_i -intervals respectively and Q_L and Q_S be the numbers of long and short q_i -intervals respectively. Here we count an end interval as $1/2$, so P_L, P_S, Q_L, Q_S are integers or half integers.

In the next section we will show that short intervals do not exist, (see (7.6)).

Proposition 6.1 $p_i = O(\epsilon^{1/3})$ and $q_i = O(\epsilon^{1/3})$.

Proof. On a short p_i or q_i -interval we ignore the nonlocal part of the energy and use Lemma 5.1 to have

$$\int_{p_i} [\frac{\epsilon^2}{2}|u'_\epsilon|^2 + W(u_\epsilon) + \frac{1}{2}|v'_\epsilon|^2] dx \geq c_1\epsilon, \quad \int_{q_i} [\frac{\epsilon^2}{2}|u'_\epsilon|^2 + W(u_\epsilon) + \frac{1}{2}|v'_\epsilon|^2] dx \geq c_{-1}\epsilon. \quad (6.2)$$

Here we have again used the fact

$$2c_1\epsilon + p_i O(e^{-\mu p_i/\epsilon}) = 2c_1\epsilon + \epsilon O(\frac{p_i}{\epsilon} e^{-\mu p_i/\epsilon}) = 2c_1\epsilon + o(\epsilon) \geq c_1\epsilon$$

when ϵ is small. If an end interval, p_1 or $p_{M_\epsilon+1}$, happens to be short, replace $c_1\epsilon$ by $\frac{c_1}{2}\epsilon$ in (6.2).

On a long interval we note by Proposition 5.3 that $O(\epsilon)p_i^2$ which appears in Lemma 5.2 is $O(\epsilon^{13/9})$. Then by Lemmas 5.1, 5.2 and the definition (6.1) of long intervals,

$$\begin{aligned} & \int_{p_i} [\frac{\epsilon^2}{2}|u'_\epsilon|^2 + W(u_\epsilon) + \frac{1}{2}|v'_\epsilon|^2] dx \\ & \geq 2c_1\epsilon + O(e^{-\mu p_i/\epsilon}) + [\frac{(1-m)^2}{24} + O(\epsilon^{1/3})]p_i^3 + O(\epsilon)p_i^2 \\ & = 2c_1\epsilon + [\frac{(1-m)^2}{24} + O(\epsilon^{1/3})]p_i^3 + O(\epsilon^{13/9}), \end{aligned} \quad (6.3)$$

$$\begin{aligned} & \int_{q_i} [\frac{\epsilon^2}{2}|u'_\epsilon|^2 + W(u_\epsilon) + \frac{1}{2}|v'_\epsilon|^2] dx \\ & \geq 2c_{-1}\epsilon + [\frac{(1+m)^2}{24} + O(\epsilon^{1/3})]q_i^3 + O(\epsilon^{13/9}). \end{aligned} \quad (6.4)$$

If p_i happens to be the end interval p_1 or $p_{M_\epsilon+1}$, (6.3) is replaced by

$$c_1\epsilon + [\frac{(1-m)^2}{6} + O(\epsilon^{1/3})]p_i^3 + O(\epsilon^{13/9}). \quad (6.5)$$

Sum (6.2) through (6.4) over i :

$$\begin{aligned}
I_\epsilon(u_\epsilon) &\geq c_1\epsilon P_S + c_{-1}\epsilon Q_S + O(\epsilon^{10/9}) \\
&\quad + \sum_{i:long} \{2c_1\epsilon + [\frac{(1-m)^2}{24} + O(\epsilon^{1/3})]p_i^3\} \\
&\quad + \sum_{i:long} \{2c_{-1}\epsilon + [\frac{(1+m)^2}{24} + O(\epsilon^{1/3})]q_i^3\} \tag{6.6}
\end{aligned}$$

where $O(\epsilon^{10/9})$ follows from $(P_L + Q_L)O(\epsilon^{13/9}) = N_\epsilon O(\epsilon^{13/9}) = O(\epsilon^{10/9})$ by Proposition 5.6. Again if i in the first sum of the last inequality happens to be 1 or $M_\epsilon + 1$, the quantity in the sum should read the first two terms of (6.5).

Note that with Proposition 5.6, (5.3) is simplified to

$$\sum_{i=1}^{M_\epsilon+1} p_i = \frac{1+m}{2} + O(\epsilon^{1/3}), \quad \sum_{i=1}^{M_\epsilon} q_i = \frac{1-m}{2} + O(\epsilon^{1/3}) \tag{6.7}$$

Therefore

$$\sum_{i:long} p_i = \frac{1+m}{2} + O(\epsilon^{1/3}), \quad \sum_{i:long} q_i = \frac{1-m}{2} + O(\epsilon^{1/3}), \tag{6.8}$$

since

$$\sum_{i:long} p_i = \sum_{i=1}^{M_\epsilon+1} p_i - \sum_{i:short} p_i = \frac{1+m}{2} + O(\epsilon^{1/3}) + N_\epsilon O(-\epsilon \log \epsilon).$$

We again use Lemma 5.5 to deduce, using the same convention when an end interval is involved,

$$\begin{aligned}
I_\epsilon(u_\epsilon) &\geq c_1\epsilon P_S + c_{-1}\epsilon Q_S + O(\epsilon^{10/9}) \\
&\quad + \sum_{i:long} \{2c_1\epsilon + [\frac{(1-m)^2}{24} + O(\epsilon^{1/3})]p_i^3\} \\
&\quad + \sum_{i:long} \{2c_{-1}\epsilon + [\frac{(1+m)^2}{24} + O(\epsilon^{1/3})]q_i^3\} \\
&\geq c_1\epsilon P_S + c_{-1}\epsilon Q_S + O(\epsilon^{10/9}) + 2c_1\epsilon P_L + 2c_{-1}\epsilon Q_L \\
&\quad + (P_L)^{-2} [\frac{(1-m)^2}{24} + O(\epsilon^{1/3})] (\frac{1+m}{2} + O(\epsilon^{1/3}))^3 \\
&\quad + (Q_L)^{-2} [\frac{(1+m)^2}{24} + O(\epsilon^{1/3})] (\frac{1-m}{2} + O(\epsilon^{1/3}))^3 \tag{6.9}
\end{aligned}$$

where the last step follows from (6.8) and

$$p = (P_L)^{-1} \sum_{i:long} p_i, \quad q = (Q_L)^{-1} \sum_{i:long} q_i$$

The upper bound of $I_\epsilon(u_\epsilon)$, Lemma 3.1₂, then implies

$$2c_1\epsilon P_L = O(\epsilon^{2/3}), \quad (P_L)^{-2} \left[\frac{(1-m)^2}{24} + O(\epsilon^{1/3}) \right] \left(\frac{1-m}{2} + O(\epsilon^{1/3}) \right)^3 = O(\epsilon^{2/3}).$$

Therefore, after applying a similar argument to q_i , we find

$$P_L \sim \epsilon^{-1/3}, \quad Q_L \sim \epsilon^{-1/3}. \quad (6.10)$$

(6.10) in turn simplifies (6.9) to

$$\begin{aligned} I_\epsilon(u_\epsilon) &\geq c_1\epsilon P_S + c_{-1}\epsilon Q_S + O(\epsilon) \\ &\quad + 2c_1\epsilon P_L + (P_L)^{-2} \frac{(1-m)^2}{24} \left(\frac{1+m}{2} \right)^3 \\ &\quad + 2c_{-1}\epsilon Q_L + (Q_L)^{-2} \frac{(1+m)^2}{24} \left(\frac{1-m}{2} \right)^3 \end{aligned} \quad (6.11)$$

Now the mysterious definition (2.5) of α comes to play. Relation (2.7) implies that the last two lines in (6.11) are *proportional*. They are simultaneously minimized if P_L and Q_L happen to be the integer or half integer that minimizes them. Denote this integer or half integer by R_ϵ . As in Lemma 3.1₂ $R_\epsilon \sim \epsilon^{-1/3}$. Then we deduce from (6.11), replacing both P_L and Q_L by R_ϵ ,

$$I_\epsilon(u_\epsilon) \geq c_1\epsilon P_S + c_{-1}\epsilon Q_S + O(\epsilon) + 2c_0\epsilon R_\epsilon + \frac{(1-m^2)^2}{96R_\epsilon^2}. \quad (6.12)$$

Now use $N = 2R_\epsilon$ in Lemma 3.1₁ to obtain an upper bound

$$I_\epsilon(u_\epsilon) \leq 2c_0\epsilon R_\epsilon + \frac{(1-m^2)^2}{96R_\epsilon^2} + O(\epsilon^{4/3} \log \epsilon),$$

which, combined with (6.12), gives $c_1\epsilon P_S + c_{-1}\epsilon Q_S = O(\epsilon)$. Therefore

$$P_S = O(1), \quad Q_S = O(1). \quad (6.13)$$

We now revisit (6.6) with the full power of Lemma 5.5. Because we know from (6.10)

$$p \sim \epsilon^{1/3}, \quad q \sim \epsilon^{1/3} \quad (6.14)$$

and also because of (6.10) and (6.13), using them to handle the error terms we find that (6.6) yields

$$\begin{aligned} I_\epsilon(u_\epsilon) &\geq O(\epsilon) + \sum_{i:long} \left\{ 2c_1\epsilon + \left[\frac{(1-m)^2}{24} + O(\epsilon^{1/3}) \right] p_i^3 \right\} \\ &\quad + \sum_{i:long} \left\{ 2c_1\epsilon + \left[\frac{(1+m)^2}{24} + O(\epsilon^{1/3}) \right] q_i^3 \right\} \\ &\geq O(\epsilon) + \sum_{i:long} \left\{ 2c_1\epsilon + \left[\frac{(1-m)^2}{24} + O(\epsilon^{1/3}) \right] [p^3 + 2p(p_i - p)^2] \right\} \end{aligned}$$

$$\begin{aligned}
& + \sum_{i:long} \{2c_1\epsilon + [\frac{(1+m)^2}{24} + O(\epsilon^{1/3})][q^3 + 2q(q_i - q)^2]\} \\
= & O(\epsilon) + 2c_1\epsilon P_L + (P_L)^{-2} \frac{(1-m)^2}{24} (\frac{1+m}{2})^3 \\
& + 2c_{-1}\epsilon Q_L + (Q_L)^{-2} \frac{(1+m)^2}{24} (\frac{1-m}{2})^3 \\
& + 2 \sum_{i:long} [\frac{(1+m)^2}{24} + O(\epsilon^{1/3})] p(p_i - p)^2 \\
& + 2 \sum_{i:long} [\frac{(1-m)^2}{24} + O(\epsilon^{1/3})] q(q_i - q)^2.
\end{aligned}$$

If p_i is an end interval p_1 or $p_{M_\epsilon+1}$, then in the second last line $p(p_i - p)$ should read $2p(p_i - \frac{p}{2})$. We again replace P_L and Q_L by R_ϵ , introduced before (6.12), to have a quantity less than or equal to $I_\epsilon(u_\epsilon)$. Also take $N = 2R_\epsilon$ in Lemma 3.1₁ to bound $I_\epsilon(u_\epsilon)$ from above. Combining these two bounds, as in the argument before (6.13), we obtain

$$O(\epsilon) + \sum_{i:long} \frac{(1+m)^2}{24} p(p_i - p)^2 + \sum_{i:long} \frac{(1-m)^2}{24} q(q_i - q)^2 \leq O(\epsilon^{4/3} \log \epsilon),$$

i.e.

$$\sum_{i:long} \frac{(1+m)^2}{24} p(p_i - p)^2 + \sum_{i:long} \frac{(1-m)^2}{24} q(q_i - q)^2 = O(\epsilon), \quad (6.15)$$

which implies $p(p_i - p)^2 = O(\epsilon)$, $q(q_i - q)^2 = O(\epsilon)$. The proposition follows since $p \sim \epsilon^{1/3}$ and $q \sim \epsilon^{1/3}$, (6.14). \square

With the help of Proposition 6.1 Lemma 4.6 is sharpened to

Lemma 6.2 1. For $i = 2, 3, \dots, M_\epsilon + 1$, if the q_{i-1} interval before x_{2i-2} and the p_i interval after x_{2i-2} are both long then

$$\begin{aligned}
\|u_\epsilon(\epsilon t + x_{2i-2}) - H(t)\|_{C^2[0, p_i/(2\epsilon)]} &= o(\epsilon^{1/3}), \\
\|u_\epsilon(\epsilon t + x_{2i-2}) - H(t)\|_{C^2[-q_{i-1}/(2\epsilon), 0]} &= o(\epsilon^{1/3}).
\end{aligned}$$

2. For $i = 1, 2, \dots, M_\epsilon$, if the p_i interval before x_{2i-1} and the q_i interval after x_{2i-1} are both long then

$$\begin{aligned}
\|u_\epsilon(\epsilon t + x_{2i-1}) - H(-t)\|_{C^2[0, q_i/(2\epsilon)]} &= o(\epsilon^{1/3}), \\
\|u_\epsilon(\epsilon t + x_{2i-1}) - H(-t)\|_{C^2[-p_i/(2\epsilon), 0]} &= o(\epsilon^{1/3}).
\end{aligned}$$

Proof. It follows from $\int_0^1 |v'_\epsilon|^2 = O(\epsilon^{2/3})$ and $v''_\epsilon = O(1)$ that $v'_\epsilon = o(1)$ on $(0, 1)$. Let x_{2i-2} be an α -point between two long intervals q_{i-1} and p_i . For

every $x \in (a, b) := (x_{2i-2} - \frac{q_i-1}{2}, x_{2i-2} + \frac{p_i}{2})$ by Proposition 6.1

$$v_\epsilon(x) = v_\epsilon(x_{2i-2}) + \int_{x_{2i-2}}^x v'_\epsilon dx = v_\epsilon(x_{2i-2}) + o(1) \cdot O(\epsilon^{1/3}) = v_\epsilon(x_{2i-2}) + o(\epsilon^{1/3}). \quad (6.16)$$

Let $u_\epsilon = w_\epsilon + \phi_\epsilon$ where $w_\epsilon = H(\frac{x-x_{2i-2}}{\epsilon})$. Lemma 4.6 and the definition (6.1) of long intervals imply that

$$\|\phi_\epsilon(\epsilon t + x_{2i-2})\|_{C^2[a,b]} = O(\epsilon^{1/3}) + O(e^{-\nu p_i/\epsilon}) + O(e^{-\nu q_i-1/\epsilon}) = O(\epsilon^{1/3}). \quad (6.17)$$

Rewrite (1.5₁) as

$$-\epsilon^2(w''_\epsilon + \phi''_\epsilon) + f(w_\epsilon) + f'(w_\epsilon)\phi_\epsilon + \frac{1}{2}f''(\dots)\phi_\epsilon^2 + v_\epsilon(x_{2i-2}) - \lambda_\epsilon + o(\epsilon^{1/3}) = 0,$$

which is simplified to

$$-\epsilon^2\phi''_\epsilon + f'(w_\epsilon)\phi_\epsilon + v_\epsilon(x_{2i-2}) - \lambda_\epsilon + o(\epsilon^{1/3}) = 0$$

if we use (6.17) for ϕ_ϵ in the f'' term. Multiply this equation by w'_ϵ and integrate over (a, b)

$$\int_a^b [-\epsilon^2\phi''_\epsilon w'_\epsilon + f'(w_\epsilon)\phi'_\epsilon w'_\epsilon] + \int_a^b [v_\epsilon(x_{2i-2}) - \lambda_\epsilon + o(\epsilon^{1/3})]w'_\epsilon = 0.$$

Then integrate by parts

$$(-\epsilon^2\phi'_\epsilon w'_\epsilon + \epsilon^2\phi_\epsilon w''_\epsilon)|_{x=a}^{x=b} + [v_\epsilon(x_{2i-2}) - \lambda_\epsilon + o(\epsilon^{1/3})](2 + o(1)) = 0. \quad (6.18)$$

Use (6.17) again to deduce

$$\begin{aligned} & [v_\epsilon(x_i) - \lambda_\epsilon + o(\epsilon^{1/3})](2 + o(1)) \\ &= -[-O(\epsilon^2\epsilon^{1/3}\epsilon^{-1}\epsilon^{-1})H'(\frac{x-x_{2i-2}}{\epsilon}) + O(\epsilon^2\epsilon^{1/3}\epsilon^{-2})H''(\frac{x-x_{2i-2}}{\epsilon})]_{x=a}^{x=b} \\ &= O(\epsilon^{1/3}) \cdot o(1) + O(\epsilon^{1/3}) \cdot o(1) \\ &= o(\epsilon^{1/3}). \end{aligned}$$

Therefore $v_\epsilon(x_i) - \lambda_\epsilon = o(\epsilon^{1/3})$. Combining this with (6.16), we deduce that on $(0, 1)$, $v_\epsilon - \lambda_\epsilon = o(\epsilon^{1/3})$ and u_ϵ satisfies $-\epsilon^2 u''_\epsilon + f(u_\epsilon) = o(\epsilon^{1/3})$.

Now we follow the proof of Lemma 4.6, with all the $O(\epsilon^{1/3})$, $O(e^{-\nu p_i/\epsilon})$ and $O(e^{-\nu q_i/\epsilon})$ terms replaced by $o(\epsilon^{1/3})$, to complete the proof of this lemma. \square

This upgrade to Lemma 4.6 gives us a much needed improvement of the lower bound in Lemma 5.2.

Lemma 6.3 *On a long p_i (q_i respectively) interval which is not adjacent (to the left or right) to a short interval, the nonlocal part of $I_\epsilon(u_\epsilon)$ has the lower*

bound

$$\begin{aligned}
\frac{1}{2} \int_{p_i} |v'_\epsilon|^2 &\geq \frac{(1-m)^2}{6} p_i^3 + o(\epsilon^{4/3}), \quad i = 1, M_\epsilon + 1 \\
\frac{1}{2} \int_{p_i} |v'_\epsilon|^2 &\geq \frac{(1-m)^2}{24} p_i^3 + \frac{p_i}{2} [v'_\epsilon(x_{2i-2}) - \frac{(1-m)p_i}{2} + o(\epsilon^{2/3})]^2 + o(\epsilon^{4/3}), \\
&\quad i \neq 1, M_\epsilon + 1 \\
\frac{1}{2} \int_{q_i} |v'_\epsilon|^2 &\geq \frac{(1+m)^2}{24} q_i^3 + \frac{q_i}{2} [v'_\epsilon(x_{2i-1}) + \frac{(1+m)q_i}{2} + o(\epsilon^{2/3})]^2 + o(\epsilon^{4/3}).
\end{aligned}$$

Proof. We follow the proof of Lemma 5.2 with all the $O(\epsilon^{1/3})$'s replaced by $o(\epsilon^{1/3})$, while in the mean time using Proposition 6.1 to simplify error terms. \square

As pointed out in the second remark following the proof of Lemma 5.2, there is another version of Lemma 6.3 analogous to (5.2)

$$\begin{aligned}
\frac{1}{2} \int_{p_i} |v'_\epsilon|^2 &\geq \frac{(1-m)^2}{24} p_i^3 + \frac{p_i}{2} [v'_\epsilon(x_{2i+1}) + \frac{(1-m)p_i}{2} + o(\epsilon^{2/3})]^2 + o(\epsilon^{4/3}), \\
&\quad i \neq 1, M_\epsilon + 1 \tag{6.19} \\
\frac{1}{2} \int_{q_i} |v'_\epsilon|^2 &\geq \frac{(1+m)^2}{24} q_i^3 + \frac{q_i}{2} [v'_\epsilon(x_{2i}) - \frac{(1+m)q_i}{2} + o(\epsilon^{2/3})]^2 + o(\epsilon^{4/3}).
\end{aligned}$$

7 The second estimation of p_i and q_i

The goal of this section is to improve Proposition 6.1 to $p_i \sim \epsilon^{1/3}$ and $q_i \sim \epsilon^{1/3}$. In particular we need to show that there are no short intervals. When dealing with the end intervals, this section adopts the same convention as in the last section.

We now redo the proof of Proposition 6.1 with this new lower bound, Lemma 6.3, to improve the proposition to

Proposition 7.1 $p_i \sim \epsilon^{1/3}$ and $q_i \sim \epsilon^{1/3}$.

Proof. Following the argument in the proof of Proposition 6.1 leading to (6.6), using Lemma 6.3 instead of Lemma 5.2.

More specifically on a short interval we use the same estimates (6.2). On a long interval there are two possibilities: it is adjacent to a short interval; or it is not adjacent to a short interval. In the first case, we retain the estimates (6.3) and (6.4), which are simplified by Proposition 6.1 to

$$\begin{aligned}
\int_{p_i} [\frac{\epsilon^2}{2} |u'_\epsilon|^2 + W(u_\epsilon) + \frac{1}{2} |v'_\epsilon|^2] dx &\geq 2c_1 \epsilon + \frac{(1-m)^2}{24} p_i^3 + O(\epsilon^{4/3}), \tag{7.1} \\
\int_{q_i} [\frac{\epsilon^2}{2} |u'_\epsilon|^2 + W(u_\epsilon) + \frac{1}{2} |v'_\epsilon|^2] dx &\geq 2c_{-1} \epsilon + \frac{(1+m)^2}{24} q_i^3 + O(\epsilon^{4/3}).
\end{aligned}$$

In the second case we apply Lemma 6.3 to have

$$\begin{aligned} \int_{p_i} [\frac{\epsilon^2}{2}|u'_\epsilon|^2 + W(u_\epsilon) + \frac{1}{2}|v'_\epsilon|^2] dx &\geq 2c_1\epsilon + \frac{(1-m)^2}{24}p_i^3 + o(\epsilon^{4/3}), \quad (7.2) \\ \int_{q_i} [\frac{\epsilon^2}{2}|u'_\epsilon|^2 + W(u_\epsilon) + \frac{1}{2}|v'_\epsilon|^2] dx &\geq 2c_{-1}\epsilon + \frac{(1+m)^2}{24}q_i^3 + o(\epsilon^{4/3}). \end{aligned}$$

As we sum over (6.2), (7.1) and (7.2) we note that there are at most $O(1)$ many terms from (7.1) because of (6.13), and $P_L \sim \epsilon^{-1/3}$ (and $Q_L \sim \epsilon^{-1/3}$ by (6.10)) many terms from (7.2). Therefore

$$\begin{aligned} I_\epsilon(u_\epsilon) &\geq c_1\epsilon P_S + c_{-1}\epsilon Q_S + o(\epsilon) + \sum_{i:long} [2c_1\epsilon + \frac{(1-m)^2}{24}p_i^3] \\ &\quad + \sum_{i:long} [2c_{-1}\epsilon + \frac{(1+m)^2}{24}q_i^3]. \end{aligned} \quad (7.3)$$

Formula (6.8) needs to be improved as well. Because of (6.13) and the definition of short intervals

$$m = \int_0^1 u_\epsilon dx = \sum_{i:long} \int_{p_i} u_\epsilon dx + \sum_{i:long} \int_{q_i} u_\epsilon dx + O(\epsilon \log \epsilon).$$

Again every long p_i or q_i interval is further divided in the middle, except the end intervals. For example, with $U_\epsilon(t) = u_\epsilon(\epsilon t + x_{2i-2})$,

$$\begin{aligned} \int_{x_{2i-2}}^{x_{2i-2} + \frac{p_i}{2}} u_\epsilon dx &= \epsilon \int_0^{p_i/(2\epsilon)} U_\epsilon dt \\ &= \epsilon \int_0^{p_i/(2\epsilon)} (U_\epsilon - H) dt + \epsilon \int_0^{p_i/(2\epsilon)} (H - 1) dt + \frac{p_i}{2}. \end{aligned}$$

Now if one of the intervals before or after x_{2i-2} is short, we use the same estimate as in the proof of Proposition 6.1, i.e.

$$\int_{x_{2i-2}}^{x_{2i-2} + \frac{p_i}{2}} u_\epsilon dx = \frac{p_i}{2} + p_i O(\epsilon^{1/3}) + O(\epsilon) = \frac{p_i}{2} + O(\epsilon^{2/3}).$$

There are at most $O(1)$ many such x_{2i-2} 's. If neither of the intervals before or after x_{2i-2} is short, we use Lemma 6.2 to find

$$\int_{x_{2i-2}}^{x_{2i-2} + \frac{p_i}{2}} u_\epsilon dx = p_i o(\epsilon^{1/3}) + O(\epsilon) + \frac{p_i}{2} = \frac{p_i}{2} + o(\epsilon^{2/3}).$$

There are $P_L \sim \epsilon^{-1/3}$ many such x_{2i-2} 's.

Now we sum over all long intervals to find

$$\sum_{i:long} p_i - \sum_{i:long} q_i = m + o(\epsilon^{1/3}).$$

On the other hand by (6.13)

$$\sum_{i:long} p_i + \sum_{i:long} q_i = 1 - \sum_{i:short} p_i - \sum_{i:short} q_i = 1 - O(1) \cdot O(\epsilon \log \epsilon) = 1 + o(\epsilon^{1/3}).$$

The last two equations imply

$$\sum_{i:long} p_i = \frac{1+m}{2} + o(\epsilon^{1/3}), \quad \sum_{i:long} q_i = \frac{1-m}{2} + o(\epsilon^{1/3}). \quad (7.4)$$

Again we set

$$p = (P_L)^{-1} \sum_{i:long} p_i, \quad q = (Q_L)^{-1} \sum_{i:long} q_i$$

and continue from (7.3) with the help of Lemma 5.5 and (7.4)

$$\begin{aligned} I_\epsilon(u_\epsilon) &\geq c_1 \epsilon P_S + c_{-1} \epsilon Q_S + o(\epsilon) \\ &\quad + \sum_{i:long} [2c_1 \epsilon + \frac{(1-m)^2}{24} p^3] + \sum_{i:long} [2c_{-1} \epsilon + \frac{(1+m)^2}{24} q^3] \\ &\geq c_1 \epsilon P_S + c_{-1} \epsilon Q_S + o(\epsilon) + 2c_1 \epsilon P_L + 2c_{-1} \epsilon Q_L \\ &\quad + (P_L)^{-2} \frac{(1-m)^2}{24} \left(\frac{1+m}{2} + o(\epsilon^{1/3})\right)^3 \\ &\quad + (Q_L)^{-2} \frac{(1+m)^2}{24} \left(\frac{1-m}{2} + o(\epsilon^{1/3})\right)^3 \\ &= c_1 \epsilon P_S + c_{-1} \epsilon Q_S + o(\epsilon) + 2c_1 \epsilon P_L + 2c_{-1} \epsilon Q_L + \\ &\quad (P_L)^{-2} \frac{(1-m)^2}{24} \left(\frac{1+m}{2}\right)^3 + (Q_L)^{-2} \frac{(1+m)^2}{24} \left(\frac{1-m}{2}\right)^3, \end{aligned} \quad (7.5)$$

where the simplification of error terms to the last two lines uses the estimate (6.10) of P_L and Q_L .

The last quantity is further reduced after we replace P_L and Q_L both by R_ϵ , introduced before (6.12). Also take $N = 2R_\epsilon$ in Lemma 3.1₁ to have an upper bound. Combine these two bounds to deduce

$$\begin{aligned} &2c_0 \epsilon R_\epsilon + \frac{(1-m^2)^2}{96R_\epsilon^2} + O(\epsilon^{4/3} \log \epsilon) \\ &\geq c_1 \epsilon P_S + c_{-1} \epsilon Q_S + o(\epsilon) + 2c_0 \epsilon R_\epsilon + \frac{(1-m^2)^2}{96R_\epsilon^2}, \end{aligned}$$

which leads to $c_1 \epsilon P_S + c_{-1} \epsilon Q_S = o(\epsilon)$. Hence

$$P_S = Q_S = 0. \quad (7.6)$$

There are no short intervals and $P_L = Q_L = M_\epsilon = N_\epsilon/2 \sim \epsilon^{-1/3}$.

Revisit (7.3) to deduce, using (7.4), (7.6) and Lemma 5.5,

$$I_\epsilon(u_\epsilon) \geq \sum_{i=1}^{M_\epsilon+1} [2c_1 \epsilon + \frac{(1-m)^2}{24} p_i^3] + \sum_{i=1}^{M_\epsilon} [2c_{-1} \epsilon + \frac{(1+m)^2}{24} q_i^3] + o(\epsilon)$$

$$\begin{aligned}
&\geq \sum_{i=1}^{M_\epsilon+1} [2c_1\epsilon + \frac{(1-m)^2}{24}p^3] + \sum_{i=1}^{M_\epsilon} [2c_{-1}\epsilon + \frac{(1+m)^2}{24}q^3] + o(\epsilon) \\
&\quad + \frac{(1-m)^2}{24} \sum_{i=1}^{M_\epsilon+1} p(p_i - p)^2 + \frac{(1+m)^2}{24} \sum_{i=1}^{M_\epsilon} q(q_i - q)^2 \\
&= o(\epsilon) + 2c_1\epsilon M_\epsilon + \frac{(1-m)^2}{24} \left(\frac{1+m}{2}\right)^3 M_\epsilon^{-2} \\
&\quad + 2c_{-1}\epsilon M_\epsilon + \frac{(1+m)^2}{24} \left(\frac{1-m}{2}\right)^3 M_\epsilon^{-2} \\
&\quad + \frac{(1-m)^2}{24} \sum_{i=1}^{M_\epsilon+1} p(p_i - p)^2 + \frac{(1+m)^2}{24} \sum_{i=1}^{M_\epsilon} q(q_i - q)^2 \\
&= o(\epsilon) + c_0\epsilon N_\epsilon + \frac{(1-m^2)^2}{24N_\epsilon^2} \\
&\quad + \frac{(1-m)^2}{24} \sum_{i=1}^{M_\epsilon+1} p(p_i - p)^2 + \frac{(1+m)^2}{24} \sum_{i=1}^{M_\epsilon} q(q_i - q)^2. \quad (7.7)
\end{aligned}$$

Use N_ϵ for N in Lemma 3.1₁, and deduce, as in (6.15),

$$\frac{(1-m)^2}{24} \sum_{i=1}^{M_\epsilon+1} p(p_i - p)^2 + \frac{(1+m)^2}{24} \sum_{i=1}^{M_\epsilon} q(q_i - q)^2 = o(\epsilon). \quad (7.8)$$

This implies, since $p \sim \epsilon^{1/3}$ and $q \sim \epsilon^{1/3}$ by (6.14), that $p_i - p = o(\epsilon^{1/3})$, $q_i - q = o(\epsilon^{1/3})$. Therefore $p_i \sim \epsilon^{1/3}$ and $q_i \sim \epsilon^{1/3}$. \square

We turn our attention to the zeros of v'_ϵ from the α -points of u_ϵ .

Proposition 7.2 *Let $x_1, x_2, \dots, x_{N_\epsilon}$ be the α -points of u_ϵ . Then v'_ϵ has exactly $N_\epsilon - 1$ zeros, denoted by $y_1, y_2, \dots, y_{N_\epsilon-1}$, in $(0, 1)$, distributed between the α -points of u_ϵ , i.e.*

$$0 < x_1 < y_1 < x_2 < y_2 < \dots < x_{N_\epsilon-1} < y_{N_\epsilon-1} < x_{N_\epsilon} < 1,$$

with the property $y_i = \frac{x_i + x_{i+1}}{2} + o(\epsilon^{1/3})$. In particular $y_{i+1} - y_i \sim \epsilon^{1/3}$.

Proof. We first claim that for $i = 1, 2, \dots, M_\epsilon$

$$v'_\epsilon(x_{2i-1}) = -\frac{(1+m)q_i}{2} + o(\epsilon^{1/3}), \quad v'_\epsilon(x_{2i-2}) = \frac{(1-m)p_i}{2} + o(\epsilon^{1/3}). \quad (7.9)$$

The careful reader may have noticed that $v'_\epsilon(x_{N_\epsilon})$ is not covered here. We will fix this problem later. We assemble a lower bound for $I_\epsilon(u_\epsilon)$ one last time, using Lemmas 5.1, 6.3, (7.6) and Proposition 7.1,

$$I_\epsilon(u_\epsilon) \geq c_0\epsilon N_\epsilon + \frac{(1-m)^2}{24} [4p_1^3 + \sum_{i=2}^{M_\epsilon} p_i^3 + 4p_{M_\epsilon+1}^3] + \frac{(1+m)^2}{24} \sum_{i=1}^{M_\epsilon} q_i^3$$

$$\begin{aligned}
& + \sum_{i=2}^{M_\epsilon} \frac{p_i}{2} [v'_\epsilon(x_{2i-2}) - \frac{(1-m)p_i}{2} + o(\epsilon^{2/3})]^2 \\
& + \sum_{i=1}^{M_\epsilon} \frac{q_i}{2} [v'_\epsilon(x_{2i-1}) + \frac{(1+m)q_i}{2} + o(\epsilon^{2/3})]^2 + o(\epsilon) \\
\geq & c_0 \epsilon N_\epsilon + \frac{(1-m^2)^2}{24N_\epsilon^2} + o(\epsilon) \\
& + \sum_{i=2}^{M_\epsilon} \frac{p_i}{2} [v'_\epsilon(x_{2i-2}) - \frac{(1-m)p_i}{2} + o(\epsilon^{2/3})]^2 \\
& + \sum_{i=1}^{M_\epsilon} \frac{q_i}{2} [v'_\epsilon(x_{2i-1}) + \frac{(1+m)q_i}{2} + o(\epsilon^{2/3})]^2 \tag{7.10}
\end{aligned}$$

where the last inequality follows from Lemma 5.5, (7.4) as in (7.7). Note that this is the only place where the full power of Lemma 6.3 is realized. We match this lower bound with the upper bound, Lemma 3.1₁ setting $N = N_\epsilon$. Then

$$\begin{aligned}
& \sum_{i=2}^{M_\epsilon} \frac{p_i}{2} [v'_\epsilon(x_{2i-2}) - \frac{(1-m)p_i}{2} + o(\epsilon^{2/3})]^2 + \\
& \sum_{i=1}^{M_\epsilon} \frac{q_i}{2} [v'_\epsilon(x_{2i-1}) + \frac{(1+m)q_i}{2} + o(\epsilon^{2/3})]^2 = o(\epsilon).
\end{aligned}$$

Since $p_i, q_i \sim \epsilon^{1/3}$, (Proposition 7.1), we obtain (7.9).

We now fix the problem about $v'_\epsilon(x_{N_\epsilon})$ in this claim. Simply repeat the same argument with (6.19), the other version of Lemma 6.3 mentioned after its proof. Then we find that for $i = 1, 2, \dots, M_\epsilon$

$$v'_\epsilon(x_{2i-1}) = -\frac{(1-m)p_i}{2} + o(\epsilon^{1/3}), \quad v'_\epsilon(x_{2i}) = \frac{(1+m)q_i}{2} + o(\epsilon^{1/3}). \tag{7.11}$$

We take up the example of x_1 and x_2 between which we will find y_1 . Other cases can be handled similarly. Estimate $v'_\epsilon(x_1)$ by (7.9) and $v'_\epsilon(x_2)$ by (7.11)

$$v'_\epsilon(x_1) = -\frac{(1+m)q_1}{2} + o(\epsilon^{1/3}), \quad v'_\epsilon(x_2) = \frac{(1+m)q_1}{2} + o(\epsilon^{1/3}). \tag{7.12}$$

We make a note here that estimating $v'_\epsilon(x_2)$ by (7.9) will give $(1+m)q_1 = (1-m)p_2 + o(\epsilon^{1/3})$. By (7.12) there exists $y_1 \in (x_1, x_2)$ such that $v'_\epsilon(y_1) = 0$, since $q_1 \sim \epsilon^{1/3}$ by Proposition 7.1.

Next we estimate $y_1 - x_1$. For this purpose we use (5.1) to find

$$0 = v'_\epsilon(y_1) = v'_\epsilon(x_1) + (1+m)(y_1 - x_1) + O(\epsilon^{2/3}),$$

which implies, with the help of (7.12),

$$y_1 - x_1 = -\frac{v'_\epsilon(x_1)}{1+m} + O(\epsilon^{2/3}) = \frac{q_1}{2} + o(\epsilon^{1/3}).$$

Finally we see that y_1 , which must be in an $o(\epsilon^{1/3})$ neighborhood of $\frac{x_1+x_2}{2}$, is unique. For by Lemma 6.2 in this neighborhood $v_\epsilon'' \sim 1 + m$, so v_ϵ' is strictly increasing there. \square

8 The one layer local minima of $J_{\epsilon,l}$

Let $l_i = y_i - y_{i-1}$, $i = 1, \dots, N_\epsilon$, where $y_0 = 0$, $y_{N_\epsilon} = 1$. Between two zero points of v_ϵ' we integrate the equation $-v_\epsilon'' = u_\epsilon - m$ and find $\frac{1}{l_i} \int_{y_{i-1}}^{y_i} u_\epsilon dx = m$. This allows us to localize the energy of u_ϵ on (y_{i-1}, y_i) . If we set $l_i z + y_{i-1} = x$, $\mathcal{U}_{\epsilon,i}(z) = u_\epsilon(x)$, and $\mathcal{V}_{\epsilon,i}(z) = l_i^{-2} v_\epsilon(x)$, then $\int_0^1 \mathcal{U}_{\epsilon,i} dz = m$, $-\mathcal{V}_{\epsilon,i}'' = \mathcal{U}_{\epsilon,i} - m$, $\mathcal{V}_{\epsilon,i}'(0) = \mathcal{V}_{\epsilon,i}'(1) = 0$. More importantly

$$\begin{aligned} I_\epsilon(u_\epsilon) &= \sum_{i=1}^{N_\epsilon} \int_{y_{i-1}}^{y_i} \left[\frac{\epsilon^2}{2} |u_\epsilon'|^2 + W(u_\epsilon) + \frac{1}{2} |v_\epsilon'|^2 \right] dx \\ &= \sum_{i=1}^{N_\epsilon} l_i \int_0^1 \left[\frac{\epsilon^2}{2l_i^2} |\mathcal{U}_{\epsilon,i}'|^2 + W(\mathcal{U}_{\epsilon,i}) + \frac{l_i^2}{2} |\mathcal{V}_{\epsilon,i}'|^2 \right] dz \\ &= \sum_{i=1}^{N_\epsilon} l_i J_{\epsilon,l_i}(\mathcal{U}_{\epsilon,i}), \end{aligned} \quad (8.1)$$

if we define a new variational problem:

$$J_{\epsilon,l}(\mathcal{U}) = \int_0^1 \left[\frac{\epsilon^2}{2l^2} |\mathcal{U}'|^2 + W(\mathcal{U}) + \frac{l^2}{2} |(-D^2)^{-1/2}(\mathcal{U} - m)|^2 \right] dz, \quad \mathcal{U} \in X_m. \quad (8.2)$$

This problem has two parameters, ϵ and l . Because of Proposition 7.2, we only need to consider the range of ϵ and l that satisfies $l \sim \epsilon^{1/3}$, i.e. we assume that there exist C_1 and C_2 such that

$$\epsilon \rightarrow 0, \quad C_1 \epsilon^{1/3} \leq l \leq C_2 \epsilon^{1/3}. \quad (8.3)$$

It is sometimes more convenient to use a different pair of parameters, ε and d , where

$$\varepsilon = \frac{\epsilon}{l} \sim \epsilon^{2/3} \rightarrow 0, \quad d = \frac{l^3}{\epsilon} \sim 1. \quad (8.4)$$

With respect to these new parameters $J_{\epsilon,l}$ in (8.2) takes the form

$$J_{\varepsilon,d}(\mathcal{U}) = \int_0^1 \left[\frac{\varepsilon^2}{2} |\mathcal{U}'|^2 + W(\mathcal{U}) + \frac{\varepsilon d}{2} |(-D^2)^{-1/2}(\mathcal{U} - m)|^2 \right] dz. \quad (8.5)$$

The Euler-Lagrange equation of this functional is

$$\begin{cases} -\varepsilon^2 \mathcal{U}'' + f(\mathcal{U}) + \varepsilon d (-D^2)^{-1}(\mathcal{U} - m) = \lambda \\ \mathcal{U}'(0) = \mathcal{U}'(1) = 0, \quad \int_0^1 \mathcal{U} dz = m. \end{cases} \quad (8.6)$$

It was proved in Theorem 1.1 [12] that $J_{\varepsilon,d}$ has a number of local minima. We focus on the ones with one transition layer. The theorem asserts that there exists $\delta > 0$, independent of ε and d , such that in the ball

$$B_\delta = \{\mathcal{U} \in L^2(0,1) : \|\mathcal{U} - \mathcal{U}_0\|_2 < \delta\}$$

there is \mathcal{U}_ε with

$$J_{\varepsilon,d}(\mathcal{U}_\varepsilon) = \inf\{J_{\varepsilon,d}(\mathcal{U}) : \mathcal{U} \in B_\delta\},$$

for all ε and d in the range (8.4). Here $\mathcal{U}_0 \in X_m$ is the same function as in (3.3).

Note that in its notation the local minimum \mathcal{U}_ε 's dependence on d is suppressed. Also proved in Theorem 1.1 [12] was that

$$\lim_{\varepsilon \rightarrow 0} \|\mathcal{U}_\varepsilon - \mathcal{U}_0\|_{L^2(0,1)} = 0, \quad (8.7)$$

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} J_{\varepsilon,d}(\mathcal{U}_\varepsilon) = c_0 + \int_0^1 \frac{d}{2} |(-D^2)^{-1/2}(\mathcal{U}_0 - m)|^2 dz. \quad (8.8)$$

The reversal $\mathcal{U}_\varepsilon^R$ of \mathcal{U}_ε , i.e. $\mathcal{U}_\varepsilon^R(z) = \mathcal{U}_\varepsilon(1-z)$, is a local minimum of $J_{\varepsilon,d}$ in

$$B_\delta^R = \{\mathcal{U} \in L^2(0,1) : \|\mathcal{U} - \mathcal{U}_0^R\| < \delta\}$$

where \mathcal{U}_0^R is the reversal of \mathcal{U}_0 . $\mathcal{U}_\varepsilon^R$ has properties similar to (8.7) and (8.8). δ is sufficiently small so that $B_\delta \cap B_\delta^R = \emptyset$.

Given \mathcal{U}_ε let $\mathcal{V}_\varepsilon = (-D^2)^{-1}(\mathcal{U}_\varepsilon - m)$, and λ_ε the Lagrange multiplier of (8.6) associated with \mathcal{U}_ε . Following the argument of Proposition 4.1, with the help of (8.8), we find

$$\begin{cases} \|\mathcal{V}_\varepsilon\|_{L^\infty(0,1)} = O(1), \\ \lambda_\varepsilon = O(\varepsilon^{1/2}), \\ -1 + O(\varepsilon^{1/2}) \leq \mathcal{U}_\varepsilon \leq 1 + O(\varepsilon^{1/2}). \end{cases} \quad (8.9)$$

As in the earlier sections, we often study \mathcal{U}_ε on a smaller scale. Let $z_\varepsilon \in (0,1)$. Introduce $U_\varepsilon(t) = \mathcal{U}_\varepsilon(\varepsilon t + z_\varepsilon)$. Then (8.6₁) and (8.9_{1,2,3}) imply that $-U_\varepsilon'' + f(U_\varepsilon) = O(\varepsilon^{1/2})$ and $U_\varepsilon \rightarrow G$ locally in C^1 (at least) where G is a solution of $-G'' + f(G) = 0$. Similar to Lemma 4.2 we have that G is heteroclinic or ± 1 . For if this is not true, then $G = \omega$, the local maximum of W , or is periodic. In either case for $\theta > 0$

$$\liminf_{\varepsilon \rightarrow 0} \varepsilon^{-1} J_{\varepsilon,d}(\mathcal{U}_\varepsilon) \geq \liminf_{\varepsilon \rightarrow 0} \varepsilon^{-1} \int_{z_\varepsilon - \varepsilon\theta}^{z_\varepsilon + \varepsilon\theta} W(\mathcal{U}_\varepsilon) dz = \int_{-\theta}^{\theta} W(G) dt.$$

The last quantity can be made arbitrarily large if we choose θ large. This contradicts (8.8).

In this section we need not use α to characterize transition layers. But for the sake of consistency we continue to do so. Following the same arguments in Propositions 4.3, 4.4 and 4.5, we obtain

Proposition 8.1 1. At every α -point z_ε , $\mathcal{U}_\varepsilon'(z_\varepsilon) \neq 0$.

2. If z_ε is an α -point, $\frac{\varepsilon}{z_\varepsilon} = o(1)$ and $\frac{\varepsilon}{1-z_\varepsilon} = o(1)$.

3. If z_ε and z_ε^* are two α -points, $\frac{\varepsilon}{|z_\varepsilon - z_\varepsilon^*|} = o(1)$.

Proposition 8.2 *When ε is small, \mathcal{U}_ε has a unique α -point, denoted by z_ε . As $\varepsilon \searrow 0$, $z_\varepsilon \rightarrow \frac{1-m}{2}$ and $\|\mathcal{U}_\varepsilon - H(\frac{\cdot - z_\varepsilon}{\varepsilon})\|_{L^\infty(0,1)} \rightarrow 0$.*

Proof. To see the existence of an α -point, note that $\int_0^1 \mathcal{U}_\varepsilon = m$ implies that there exists z'_ε where $\mathcal{U}_\varepsilon(z'_\varepsilon) = m$. Similar to the location of α -points (Proposition 8.1₂), $\frac{\varepsilon}{z'_\varepsilon} = o(1)$ and $\frac{\varepsilon}{1-z'_\varepsilon} = o(1)$. $\mathcal{U}_\varepsilon(\varepsilon t + z'_\varepsilon)$ converges in C^1 to a heteroclinic solution of $-G'' + f(G) = 0$ with $G(0) = m$ by the remarks following (8.9). Then $\mathcal{U}_\varepsilon(z_\varepsilon) = \alpha$ at a point z_ε , such that $|z_\varepsilon - z'_\varepsilon| = O(\varepsilon)$.

To see the uniqueness of z_ε , suppose in contrary there are two α -points, z_ε and z_ε^* , of \mathcal{U}_ε . Without the loss of generality assume $\mathcal{U}'_\varepsilon(z_\varepsilon) > 0$ and $\mathcal{U}'_\varepsilon(z_\varepsilon^*) < 0$ by Proposition 8.1₁. Then by Proposition 8.1_{2,3} and the remarks after (8.9) for every $\theta > 0$, as $\varepsilon \searrow 0$,

$$\begin{aligned} & \varepsilon^{-1} \int_0^1 \left[\frac{\varepsilon^2}{2} |\mathcal{U}'_\varepsilon|^2 + W(\mathcal{U}_\varepsilon) \right] dz \\ & \geq \int_{-\theta}^\theta \left[\frac{1}{2} \left| \frac{d}{dt} \mathcal{U}_\varepsilon(\varepsilon t + z_\varepsilon) \right|^2 + W(\mathcal{U}_\varepsilon(\varepsilon t + z_\varepsilon)) \right] dt \\ & \quad + \int_{-\theta}^\theta \left[\frac{1}{2} \left| \frac{d}{dt} \mathcal{U}_\varepsilon(\varepsilon t + z_\varepsilon^*) \right|^2 + W(\mathcal{U}_\varepsilon(\varepsilon t + z_\varepsilon^*)) \right] dt \\ & \rightarrow \int_{-\theta}^\theta \left[\frac{1}{2} |H'(t)|^2 + W(H(t)) \right] dt + \int_{-\theta}^\theta \left[\frac{1}{2} |H'(-t)|^2 + W(H(-t)) \right] dt \\ & \geq \frac{3c_0}{2} \end{aligned}$$

if we choose θ large enough. On the other hand

$$\int_0^1 \frac{d}{2} |(-D^2)^{-1/2} (\mathcal{U}_\varepsilon - m)|^2 dz \rightarrow \int_0^1 \frac{d}{2} |(-D^2)^{-1/2} (\mathcal{U}_0 - m)|^2 dz,$$

because of (8.7) and the continuity of the nonlocal part of $J_{\varepsilon,d}$ under the L^2 norm. Therefore

$$\liminf_{\varepsilon \rightarrow 0} \varepsilon^{-1} J_{\varepsilon,d}(\mathcal{U}_\varepsilon) \geq \frac{3c_0}{2} + \int_0^1 \frac{d}{2} |(-D^2)^{-1/2} (\mathcal{U}_0 - m)|^2 dz,$$

contradicting (8.8).

$\mathcal{U}_\varepsilon(\varepsilon t + z_\varepsilon)$ converges locally in C^1 to $H(t)$ or $H(-t)$. We show that the first case implies the conclusions of this proposition, and the second case does not occur. Assume that $H(t)$ is the local limit. If $\|\mathcal{U}_\varepsilon - H(\frac{\cdot - z_\varepsilon}{\varepsilon})\|_\infty = o(1)$ is false, there exists $h_\varepsilon \in (0,1)$ such that $\frac{|z_\varepsilon - h_\varepsilon|}{\varepsilon} \rightarrow \infty$ and $|\mathcal{U}_\varepsilon(h_\varepsilon) - H(\frac{h_\varepsilon - z_\varepsilon}{\varepsilon})|$ stays

away from 0. Thus $|\mathcal{U}_\varepsilon(h_\varepsilon)|$ stays away from 1. Now consider $\mathcal{U}_\varepsilon(\varepsilon t + h_\varepsilon)$, which converges locally in C^1 to a heteroclinic solution of $-G'' + f(G) = 0$. Because the derivative of the heteroclinic solution is never zero and $\mathcal{U}'_\varepsilon(0) = \mathcal{U}'_\varepsilon(1) = 0$, $\frac{h_\varepsilon}{\varepsilon} \rightarrow \infty$ and $\frac{1-h_\varepsilon}{\varepsilon} \rightarrow \infty$. There exists $t_\varepsilon = O(1)$ such that $\varepsilon t_\varepsilon + h_\varepsilon \in (0, 1)$ and $\mathcal{U}_\varepsilon(\varepsilon t_\varepsilon + h_\varepsilon) = \alpha$. But $\frac{|\varepsilon t_\varepsilon + h_\varepsilon - z_\varepsilon|}{\varepsilon} \rightarrow \infty$. So we have found two α -points z_ε and $\varepsilon t_\varepsilon + h_\varepsilon$, contradicting the uniqueness of z_ε . Finally $\|\mathcal{U}_\varepsilon - H(\frac{\cdot - z_\varepsilon}{\varepsilon})\|_\infty = o(1)$ and $\int_0^1 \mathcal{U}_\varepsilon dz = m$ yield that $z_\varepsilon \rightarrow \frac{1-m}{2}$.

If $\mathcal{U}_\varepsilon(\varepsilon t + z_\varepsilon)$ converges locally in C^1 to $H(-t)$, the same argument leads to $\|\mathcal{U}_\varepsilon - H(\frac{z_\varepsilon - \cdot}{\varepsilon})\|_\infty = o(1)$ and $z_\varepsilon \rightarrow \frac{1+m}{2}$. Therefore $\mathcal{U}_\varepsilon \in B_\delta^R$ for small ε , contradicting $B_\delta \cap B_\delta^R = \emptyset$. \square

We define

$$\phi_0(z) = \begin{cases} -\frac{d[\mathcal{V}_0(z) - \mathcal{V}_0(\frac{1-m}{2})]}{f'(-1)}, & 0 < z \leq \frac{1-m}{2} \\ -\frac{d[\mathcal{V}_0(z) - \mathcal{V}_0(\frac{1-m}{2})]}{f'(1)}, & \frac{1-m}{2} < z < 1 \end{cases} \quad (8.10)$$

where $\mathcal{V}_0 = (-D^2)^{-1}(\mathcal{U}_0 - m)$ (see (3.4)). This function's derivative has a jump discontinuity at $\frac{1-m}{2}$, unless $f'(-1) = f'(1)$.

Proposition 8.3

$$\begin{aligned} \mathcal{U}_\varepsilon(z) &= H\left(\frac{z - z_\varepsilon}{\varepsilon}\right) + \phi_0(z)\varepsilon + O(\varepsilon^2), \\ z_\varepsilon &= \frac{1-m}{2} + c_3\varepsilon + O(\varepsilon^2), \end{aligned}$$

where $c_3 = \frac{1}{2}(\int_{-\infty}^0 (H+1) dt + \int_0^\infty (H-1) dt + \int_0^1 \phi_0 dz)$.

Proof. The first several steps are similar to those in the proof of Lemma 4.6. Anticipating an asymptotic expansion, we write

$$\mathcal{U}_\varepsilon(z) = H\left(\frac{z - z_\varepsilon}{\varepsilon}\right) + \phi_\varepsilon(z)\varepsilon.$$

By (8.9_{1,2}) $\phi_\varepsilon\varepsilon$ satisfies $-\varepsilon^2(\phi_\varepsilon\varepsilon)'' + f'(\dots)(\phi_\varepsilon\varepsilon) = O(\varepsilon^{1/2})$. Arguing as in Step 2 of the proof of Lemma 4.6 on the intervals $(0, z_\varepsilon)$ and $(z_\varepsilon, 1)$ separately, with the help of Proposition 8.2 which asserts $\phi_\varepsilon\varepsilon = o(1)$, we deduce

$$\phi_\varepsilon\varepsilon = O(\varepsilon^{1/2}). \quad (8.11)$$

Then argue as in Steps 3 and 4 of the same lemma to obtain

$$(\phi_\varepsilon\varepsilon)' = \varepsilon^{-1}O(\varepsilon^{1/2}). \quad (8.12)$$

Because of (8.11), rewrite the equation for $\phi_\varepsilon\varepsilon$ as

$$-\varepsilon^2(\phi_\varepsilon\varepsilon)'' + f'(H)(\phi_\varepsilon\varepsilon) + O(\varepsilon) = \lambda_\varepsilon.$$

Multiply this by $\varepsilon^{-1}H'(\frac{z-z_\varepsilon}{\varepsilon})$ and integrate by parts (as in the proof of Lemma 6.2):

$$[-\varepsilon(\phi_\varepsilon)'H'(\frac{z-z_\varepsilon}{\varepsilon}) + (\phi_\varepsilon\varepsilon)H''(\frac{z-z_\varepsilon}{\varepsilon})]_0^1 = [\lambda_\varepsilon - O(\varepsilon)] \int_{-z_\varepsilon/\varepsilon}^{(1-z_\varepsilon)/\varepsilon} H'(t) dt.$$

The exponential decay rates of H' and H'' , (8.11), and (8.12) improve (8.9₂) to

$$\lambda_\varepsilon = O(\varepsilon). \quad (8.13)$$

This estimate implies that ϕ_ε satisfies $-\varepsilon^2\phi_\varepsilon'' + f'(\dots)\phi_\varepsilon = O(1)$. The argument before (8.11) and (8.12) gives

$$\phi_\varepsilon = O(1), \quad \phi_\varepsilon' = \varepsilon^{-1}O(1), \quad (8.14)$$

improving (8.11) and (8.12).

At this point we make a preliminary estimate of z_ε . From (8.14) we see $\mathcal{U}_\varepsilon(z) = H(\frac{z-z_\varepsilon}{\varepsilon}) + O(\varepsilon)$. Integrating this over $(0, 1)$ yields

$$\begin{aligned} m &= \int_0^1 H(\frac{z-z_\varepsilon}{\varepsilon}) dz + O(\varepsilon) \\ &= \varepsilon \int_{-z_\varepsilon/\varepsilon}^{(1-z_\varepsilon)/\varepsilon} H(t) dt + O(\varepsilon) \\ &= \varepsilon \left[\int_{-z_\varepsilon/\varepsilon}^0 (H(t) + 1) + \int_0^{(1-z_\varepsilon)/\varepsilon} (H(t) - 1) + \frac{1-2z_\varepsilon}{\varepsilon} \right] + O(\varepsilon) \\ &= 1 - 2z_\varepsilon + O(\varepsilon). \end{aligned}$$

Therefore

$$z_\varepsilon = \frac{1-m}{2} + O(\varepsilon). \quad (8.15)$$

By (8.14) we write the equation that ϕ_ε satisfies as

$$-\varepsilon^2\phi_\varepsilon'' + f'(H)\phi_\varepsilon + O(\varepsilon) + d\mathcal{V}_\varepsilon = \frac{\lambda_\varepsilon}{\varepsilon}.$$

Again multiply it by $\varepsilon^{-1}H'(\frac{z-z_\varepsilon}{\varepsilon})$ and integrate by parts:

$$\begin{aligned} &[-\varepsilon\phi_\varepsilon'H'(\frac{z-z_\varepsilon}{\varepsilon}) + \phi_\varepsilon\varepsilon H''(\frac{z-z_\varepsilon}{\varepsilon})]_0^1 \\ &= \int_0^1 \varepsilon^{-1} \left[\frac{\lambda_\varepsilon}{\varepsilon} - d\mathcal{V}_\varepsilon + O(\varepsilon) \right] H'(\frac{z-z_\varepsilon}{\varepsilon}) dz. \\ &= \int_{-z_\varepsilon/\varepsilon}^{(1-z_\varepsilon)/\varepsilon} \left[\frac{\lambda_\varepsilon}{\varepsilon} - d\mathcal{V}_\varepsilon(z_\varepsilon) + O(1)\varepsilon t + O(\varepsilon) \right] H'(t) dt \\ &= 2 \left[\frac{\lambda_\varepsilon}{\varepsilon} - d\mathcal{V}_\varepsilon(z_\varepsilon) \right] + O(\varepsilon). \end{aligned}$$

We have used the fact $\mathcal{V}'_\varepsilon = O(1)$, which follows from (8.9₃) and the regularity theory for $(-D^2)^{-1}$, to reach the third line. The exponential decay rates of H' and H'' in line one imply that

$$\lambda_\varepsilon = \varepsilon d\mathcal{V}_\varepsilon(z_\varepsilon) + O(\varepsilon^2), \quad (8.16)$$

upgrading (8.13).

With (8.16) we obtain $-\varepsilon^2\phi''_\varepsilon + f'(H)\phi_\varepsilon + d(\mathcal{V}_\varepsilon - \mathcal{V}_\varepsilon(z_\varepsilon)) = O(\varepsilon)$. On $(0, z_\varepsilon)$ set

$$\phi_\varepsilon(z) = -\frac{d(\mathcal{V}_\varepsilon(z) - \mathcal{V}_\varepsilon(z_\varepsilon))}{f'(-1)} + \psi_\varepsilon.$$

Then ψ_ε satisfies

$$-\varepsilon^2\psi''_\varepsilon + f'(H)\psi_\varepsilon + \frac{f'(-1) - f'(H)}{f'(-1)}d[\mathcal{V}_\varepsilon - \mathcal{V}_\varepsilon(z_\varepsilon)] = O(\varepsilon),$$

with the boundary conditions $\psi'_\varepsilon(0) = -\varepsilon^{-2}H'(-z_\varepsilon/\varepsilon) = \varepsilon^{-2}O(e^{-C/\varepsilon})$ and $\psi_\varepsilon(z_\varepsilon) = 0$. Note that because of $\mathcal{V}''_\varepsilon = O(1)$ by (8.9₃), and the exponential convergence rate of H to -1 at $-\infty$,

$$|[f'(-1) - f'(H(\frac{z - z_\varepsilon}{\varepsilon}))](\mathcal{V}_\varepsilon(z) - \mathcal{V}_\varepsilon(z_\varepsilon))| \leq C|(H(t) + 1)\varepsilon t| \leq C\varepsilon.$$

So the equation for ψ_ε is further simplified to $-\varepsilon^2\psi''_\varepsilon + f'(H)\psi_\varepsilon = O(\varepsilon)$. Then argue as in (8.11) to conclude that $\psi_\varepsilon = O(\varepsilon)$.

In summary we have shown, after a similar argument on $(z_\varepsilon, 1)$,

$$\mathcal{U}_\varepsilon(z) = \begin{cases} H(\frac{z - z_\varepsilon}{\varepsilon}) - \frac{\varepsilon d(\mathcal{V}_\varepsilon(z) - \mathcal{V}_\varepsilon(z_\varepsilon))}{f'(-1)} + O(\varepsilon^2), & z \in (0, z_\varepsilon) \\ \alpha, & z = z_\varepsilon \\ H(\frac{z - z_\varepsilon}{\varepsilon}) - \frac{\varepsilon d(\mathcal{V}_\varepsilon(z) - \mathcal{V}_\varepsilon(z_\varepsilon))}{f'(1)} + O(\varepsilon^2), & z \in (z_\varepsilon, 1) \end{cases}.$$

To complete the proof of the first estimate of this proposition, we compare \mathcal{V}_ε to \mathcal{V}_0 . Let $\mathcal{Z} = \mathcal{V}_\varepsilon - \mathcal{V}_0 = (-D^2)^{-1}(\mathcal{U}_\varepsilon - \mathcal{U}_0)$. According to (8.14) and (8.15),

$$\begin{aligned} \|\mathcal{U}_\varepsilon - \mathcal{U}_0\|_1 &= \int_0^1 |H(\frac{z - z_\varepsilon}{\varepsilon}) - \mathcal{U}_0| dz + O(\varepsilon) \\ &= \varepsilon \int_{-(1-m)/(2\varepsilon)}^{(1+m)/(2\varepsilon)} |H(t + O(1)) - \mathcal{U}_0(\varepsilon t + \frac{1-m}{2})| dt + O(\varepsilon) \\ &= O(\varepsilon). \end{aligned}$$

Integrating the linear differential equation for \mathcal{Z} we see

$$\mathcal{V}_\varepsilon = \mathcal{V}_0 + O(\varepsilon), \quad \mathcal{V}'_\varepsilon = \mathcal{V}'_0 + O(\varepsilon). \quad (8.17)$$

The first estimate of this proposition then follows.

To establish the second estimate, integrate the first estimate over $(0, 1)$:

$$\begin{aligned}
& m + O(\varepsilon^2) \\
&= \int_0^1 H\left(\frac{z - z_\varepsilon}{\varepsilon}\right) dz + \varepsilon \int_0^1 \phi_0 dz \\
&= \varepsilon \left[\int_{-z_\varepsilon/\varepsilon}^0 (H(t) + 1) + \int_0^{(1-z_\varepsilon)/\varepsilon} (H(t) - 1) + \frac{1 - 2z_\varepsilon}{\varepsilon} + \int_0^1 \phi_0 dz \right] \\
&= 1 - 2z_\varepsilon + \varepsilon \left[\int_{-\infty}^0 (H + 1) dt + \int_0^{\infty} (H - 1) dt + \int_0^1 \phi_0 dz + O(e^{-C/\varepsilon}) \right]. \quad \square
\end{aligned}$$

The next result will be very handy later.

Lemma 8.4 *Let $F \in C^2(-\infty, \infty)$ be such that $F(\pm 1) = 0$. Then*

$$\int_0^1 F(\mathcal{U}_\varepsilon) dz = \varepsilon \int_{-\infty}^{\infty} F(H) dt + \varepsilon \int_0^1 F'(\pm 1) \phi_0 dz + O(\varepsilon^2),$$

where ± 1 is -1 on $(0, (1 - m)/2)$ and 1 on $((1 - m)/2, 1)$.

Proof. According to Proposition 8.3

$$\begin{aligned}
\int_0^1 F(\mathcal{U}_\varepsilon) dz &= \int_0^1 F\left(H\left(\frac{z - z_\varepsilon}{\varepsilon}\right) + \varepsilon \phi_0 + O(\varepsilon^2)\right) dz \\
&= \varepsilon \int_{-z_\varepsilon/\varepsilon}^{(1-z_\varepsilon)/\varepsilon} F(H(t)) dt + \varepsilon \int_0^1 F'\left(H\left(\frac{z - z_\varepsilon}{\varepsilon}\right)\right) \phi_0 dz + O(\varepsilon^2) \\
&= \varepsilon \int_{-\infty}^{\infty} F(H) dt + \varepsilon \int_0^1 F'(\pm 1) \phi_0 dz + O(\varepsilon^2). \quad \square
\end{aligned}$$

9 The second variation of $J_{\varepsilon, d}$

We now study the second variation of J_ε at \mathcal{U}_ε and give a bound on the principle eigenvalue of the linearized operator of (8.6).

Proposition 9.1 *There is $c_4 > 0$ such that for all $\varphi \in W^{1,2}(0, 1)$, $\text{Ave}(\varphi) = 0$,*

$$\int_0^1 [\varepsilon^2 |\varphi'|^2 + f'(\mathcal{U}_\varepsilon) \varphi^2 + \varepsilon d |(-D^2)^{-1/2} \varphi|^2] dz \geq c_4 \varepsilon \int_0^1 \varphi^2 dz.$$

Proof. All we need to prove is that if Λ is an eigenvalue of the following eigenvalue problem

$$\begin{cases} -\varepsilon^2 \psi'' + f'(\mathcal{U}_\varepsilon) \psi + \varepsilon d (-D^2)^{-1} \psi = \eta + \Lambda \psi \\ \psi'(0) = \psi'(1) = 0, \text{Ave}(\psi) = 0, \psi \not\equiv 0 \end{cases}, \quad (9.1)$$

then $\Lambda \geq c_4\varepsilon$ for some constant $c_4 > 0$. Since \mathcal{U}_ε minimizes J_ε locally, Λ must be ≥ 0 . Suppose the assertion of this proposition is false. Then $\Lambda = o(\varepsilon)$.

We normalize the eigen-function ψ so that $\|\psi\|_2 = 1$. Let H_ε be a modification of H so that $H_\varepsilon(t) = -1$ if $t \leq -z_\varepsilon/(2\varepsilon)$ and $H_\varepsilon(t) = 1$ if $t \geq (1 - z_\varepsilon)/(2\varepsilon)$. Moreover $H_\varepsilon = H + O(e^{-C/\varepsilon})$, $H'_\varepsilon = H' + O(e^{-C/\varepsilon})$, and $H''_\varepsilon = H'' + O(e^{-C/\varepsilon})$. Then let $h_\varepsilon(z) = \frac{1}{\varepsilon}H'_\varepsilon(\frac{z-z_\varepsilon}{\varepsilon})$. This h_ε has compact support. It follows from Lemma 2.2₂ that for all $\varphi \in W^{1,2}(0,1)$ with $\int_0^1 \varphi h_\varepsilon = 0$,

$$\int_0^1 [\varepsilon^2|\varphi'|^2 + f'(H)\varphi^2] dz \geq c_5 \int_0^1 \varphi^2 dz. \quad (9.2)$$

We decompose $\psi = ch_\varepsilon + \psi^\perp$, $\int_0^1 h_\varepsilon \psi^\perp dz = 0$. Let $A = (-D^2)^{-1}(h_\varepsilon - \text{Ave}(h_\varepsilon))$ and $B = (-D^2)^{-1}(\psi^\perp - \text{Ave}(\psi^\perp))$. Note that

$$A = O(1), \quad B = \|\psi^\perp\|_2 O(1), \quad (9.3)$$

and $(-D^2)^{-1}\psi = cA + B$. By integrating the decomposition of ψ we find

$$\int_0^1 \psi^\perp = -2c. \quad (9.4)$$

By integrating (9.1) we observe

$$\eta = \int_0^1 f'(\mathcal{U}_\varepsilon)\psi dz = \int_0^1 [f'(H) + O(\varepsilon)](ch_\varepsilon + \psi^\perp) dz.$$

After estimating the right side, we deduce

$$|\eta| = |c|O(\varepsilon) + \|\psi^\perp\|_2 O(1). \quad (9.5)$$

The equation for ψ^\perp is

$$\begin{aligned} & -\varepsilon^2(\psi^\perp)'' + f'(\mathcal{U}_\varepsilon)\psi^\perp + \varepsilon d(cA + B) + c[f'(\mathcal{U}_\varepsilon) - f'(H)]h_\varepsilon + O(e^{-C/\varepsilon}) \\ & = \eta + \Lambda(ch + \psi^\perp). \end{aligned}$$

Multiply this by ψ^\perp and integrate by parts to obtain

$$\begin{aligned} & \int_0^1 [\varepsilon^2|(\psi^\perp)'|^2 + f'(\mathcal{U}_\varepsilon)|\psi^\perp|^2 + \varepsilon d|B'|^2 - \Lambda|\psi^\perp|^2] dz \\ & = \int_0^1 [-c(f'(\mathcal{U}_\varepsilon) - f'(H))h_\varepsilon\psi^\perp - c\varepsilon dA\psi^\perp + \eta\psi^\perp + \psi^\perp O(e^{-C/\varepsilon})] dz. \end{aligned}$$

By (9.2) and the assumption on Λ we find that the first line is $\geq c_6 \int_0^1 |\psi^\perp|^2 dz$. To estimate second line we note, with the help of (9.3),

$$\left| \int_0^1 -c(f'(\mathcal{U}_\varepsilon) - f'(H))h_\varepsilon\psi^\perp dz \right| = |c| \cdot \|\psi^\perp\|_2 O(\varepsilon^{3/2})$$

$$\begin{aligned}
\left| \int_0^1 -c\varepsilon dA\psi^\perp dz \right| &= |c| \cdot \|\psi^\perp\|_2 O(\varepsilon) \\
\left| \int_0^1 \eta\psi^\perp dz \right| &= 2|\eta| \cdot |c| \\
\left| \int_0^1 O(e^{-C/\varepsilon})\psi^\perp dz \right| &= \|\psi^\perp\|_2 O(e^{-C/\varepsilon}).
\end{aligned}$$

The first one here is less obvious. Note that

$$\begin{aligned}
&\|(f'(\mathcal{U}_\varepsilon) - f'(H))h_\varepsilon\|_2^2 \\
&\leq C \int_0^1 (\varepsilon\phi_0 + O(\varepsilon^2))^2 h_\varepsilon^2 dz \\
&= C\varepsilon^3 \int_{-z_\varepsilon/\varepsilon}^{(1-z_\varepsilon)/\varepsilon} \left[\frac{\phi_0(\varepsilon t + z_\varepsilon) - \phi_0((1-m)/2)}{\varepsilon} + O(1) \right]^2 H'_\varepsilon(t)^2 dt \\
&\leq C\varepsilon^3 \int_{-z_\varepsilon/\varepsilon}^{(1-z_\varepsilon)/\varepsilon} (|t| + O(1))^2 H'_\varepsilon(t)^2 dt \\
&= O(\varepsilon^3)
\end{aligned} \tag{9.6}$$

by Proposition 8.3, $\phi_0((1-m)/2) = 0$ and the Lipschitz continuity of ϕ_0 .

Therefore the last integral identity implies

$$c_6 \|\psi^\perp\|_2^2 \leq |c| \cdot \|\psi^\perp\|_2 O(\varepsilon) + 2|\eta| \cdot |c| + \|\psi^\perp\|_2 O(e^{-C/\varepsilon}),$$

which, combined with (9.5), leads to

$$\|\psi^\perp\|_2 \leq |c|O(1) + O(e^{-C/\varepsilon}), \tag{9.7}$$

$$|\eta| \leq |c|O(1) + O(e^{-C/\varepsilon}). \tag{9.8}$$

It also leads to, with the help of $2|c| = \left| \int_0^1 \psi^\perp dz \right| \leq \|\psi^\perp\|_2$,

$$|c| \leq |\eta|O(1) + O(e^{-C/\varepsilon}). \tag{9.9}$$

Next we multiply the equation for ψ^\perp by h_ε and integrate by parts to obtain

$$\int_0^1 [\psi^\perp O(e^{-C/\varepsilon}) + \varepsilon d(cA+B)h_\varepsilon + (f'(\mathcal{U}_\varepsilon) - f'(H))h_\varepsilon\psi] dz = \int_0^1 (\eta h_\varepsilon + \Lambda c h_\varepsilon^2) dz.$$

We estimate each term, using (9.7):

$$\begin{aligned}
\left| \int_0^1 \psi^\perp O(e^{-C/\varepsilon}) dz \right| &= |c|O(e^{-C/\varepsilon}) + O(e^{-C/\varepsilon}) \\
\left| \int_0^1 \varepsilon dcAh_\varepsilon dz \right| &= |c|O(\varepsilon) \\
\left| \int_0^1 \varepsilon dBh dz \right| &= |c|O(\varepsilon) + O(e^{-C/\varepsilon})
\end{aligned}$$

$$\begin{aligned}
\left| \int_0^1 (f'(\mathcal{U}_\varepsilon) - f'(H))h_\varepsilon \psi \, dz \right| &= |c|O(\varepsilon) + O(e^{-C/\varepsilon}) \\
\int_0^1 \eta h_\varepsilon \, dz &= -2\eta \\
\int_0^1 \Lambda c h^2 &= |c|o(1).
\end{aligned}$$

All of the above are easy with the possible exception of the fourth estimate. One writes $\psi = ch_\varepsilon + \psi^\perp$, so

$$\begin{aligned}
\left| \int_0^1 (f'(\mathcal{U}_\varepsilon) - f'(H))h_\varepsilon \psi^\perp \, dz \right| &\leq \|(f'(\mathcal{U}_\varepsilon) - f'(H))h_\varepsilon\|_2 \|\psi^\perp\|_2 \\
&= |c|O(\varepsilon^{3/2}) + O(e^{-C/\varepsilon}),
\end{aligned}$$

by (9.6). And arguing as in (9.6) we find

$$\left| \int_0^1 (f'(\mathcal{U}_\varepsilon) - f'(H))h_\varepsilon^2 \, dz \right| = O(\varepsilon).$$

The last integral identity then implies $\eta = |c|o(1) + O(e^{-C/\varepsilon})$. Because of (9.7), (9.8) and (9.9) we have $|c| = O(e^{-C/\varepsilon})$ and $\|\psi^\perp\|_2 = O(e^{-C/\varepsilon})$. So $\|\psi\|_2 = O(e^{-C/\varepsilon})$, contradicting $\|\psi\|_2 = 1$. \square

The related result in [6] has no ε after c_4 . The reason is that the extra condition that $\varphi(\frac{1}{2}) = 0$ was assumed. Here without this condition we have small eigenvalues. A consequence of this proposition is that \mathcal{U}_ε is unique.

Proposition 9.2 *For small ε , if $\mathcal{U}_\varepsilon, \mathcal{U}_\varepsilon^* \in B_\delta$ satisfying $J_{\varepsilon,d}(\mathcal{U}_\varepsilon) = J_{\varepsilon,d}(\mathcal{U}_\varepsilon^*) = \inf\{J_{\varepsilon,d}(\mathcal{U}) : \mathcal{U} \in B_\delta\}$, then $\mathcal{U}_\varepsilon = \mathcal{U}_\varepsilon^*$. The same is also true in B_δ^R .*

Proof. Let \mathcal{U}_ε and $\mathcal{U}_\varepsilon^*$ be as in the statement of the proposition. We first show that $\mathcal{U}_\varepsilon - \mathcal{U}_\varepsilon^* = O(\varepsilon^2)$, and then use Proposition 9.1 to conclude that they are identical.

The first estimate of Proposition 8.3 asserts that

$$\mathcal{U}_\varepsilon^* - \mathcal{U}_\varepsilon = H\left(\frac{z - z_\varepsilon^*}{\varepsilon}\right) - H\left(\frac{z - z_\varepsilon}{\varepsilon}\right) + O(\varepsilon^2),$$

where z_ε (z_ε^* respectively) is the α -point of \mathcal{U}_ε ($\mathcal{U}_\varepsilon^*$ respectively). The second estimate of Proposition 8.3 says $z_\varepsilon^* - z_\varepsilon = O(\varepsilon^2)$. Therefore

$$\begin{aligned}
\mathcal{U}_\varepsilon^* - \mathcal{U}_\varepsilon &= \frac{z_\varepsilon - z_\varepsilon^*}{\varepsilon} H'\left(\frac{z - z_\varepsilon}{\varepsilon}\right) + O(\varepsilon^2) \\
&= \kappa(\varepsilon) H'\left(\frac{z - z_\varepsilon}{\varepsilon}\right) \varepsilon + O(\varepsilon^2),
\end{aligned} \tag{9.10}$$

where $\kappa(\varepsilon) = \frac{z_\varepsilon - z_\varepsilon^*}{\varepsilon^2} = O(1)$.

We next show that $\kappa(\varepsilon) = O(\varepsilon^{1/2})$. Let $\mathcal{W}_\varepsilon = \mathcal{U}_\varepsilon^* - \mathcal{U}_\varepsilon$, $\mathcal{Z}_\varepsilon = \mathcal{V}_\varepsilon^* - \mathcal{V}_\varepsilon$, where $\mathcal{V}_\varepsilon = (-D^2)^{-1}(\mathcal{U}_\varepsilon - m)$ and $\mathcal{V}_\varepsilon^* = (-D^2)^{-1}(\mathcal{U}_\varepsilon^* - m)$. \mathcal{W}_ε satisfies

$$-\varepsilon^2 \mathcal{W}_\varepsilon'' + f'(\mathcal{U}_\varepsilon) \mathcal{W}_\varepsilon + \varepsilon d \mathcal{Z}_\varepsilon + [f(\mathcal{U}_\varepsilon^*) - f(\mathcal{U}_\varepsilon) - f'(\mathcal{U}_\varepsilon) \mathcal{W}_\varepsilon] = \lambda_\varepsilon^* - \lambda_\varepsilon.$$

Multiply by \mathcal{W}_ε and integrate by parts to deduce, with the help of (9.10):

$$\begin{aligned} & \int_0^1 [\varepsilon^2 |\mathcal{W}_\varepsilon'|^2 + f'(\mathcal{U}_\varepsilon) \mathcal{W}_\varepsilon^2 + \varepsilon d |\mathcal{Z}_\varepsilon'|^2] dz \\ &= - \int_0^1 [f(\mathcal{U}_\varepsilon^*) - f(\mathcal{U}_\varepsilon) - f'(\mathcal{U}_\varepsilon) \mathcal{W}_\varepsilon] \mathcal{W}_\varepsilon dz \\ &= - \frac{1}{2} \int_0^1 f''(H(\frac{z - z_\varepsilon}{\varepsilon}) + O(\varepsilon)) (\kappa(\varepsilon) H'(\frac{z - z_\varepsilon}{\varepsilon}) \varepsilon + O(\varepsilon^2))^3 dz \\ &= - \frac{\varepsilon^4}{2} \int_{-z_\varepsilon/\varepsilon}^{(1-z_\varepsilon)/\varepsilon} f''(H(t) + O(\varepsilon)) (\kappa(\varepsilon) H'(t) + O(\varepsilon))^3 dt \\ &= - \frac{\kappa(\varepsilon) \varepsilon^4}{2} \int_{-\infty}^{\infty} f''(H) (H')^3 dt + O(\varepsilon^5) \\ &= O(\varepsilon^5), \end{aligned} \tag{9.11}$$

since by (2.4) and integration by parts

$$\int_{-\infty}^{\infty} f''(H) (H')^3 dt = \int_{-1}^1 2f''(H) W(H) dH = -(f(H))^2 \Big|_{H=-1}^{H=1} = 0.$$

Note that when the Taylor expansion is used in line three of (9.11), by Proposition 8.3 and (9.10) both \mathcal{U}_ε and $\mathcal{U}_\varepsilon^*$ are $H(\frac{z - z_\varepsilon}{\varepsilon}) + O(\varepsilon)$. So we put it in $f''(\dots)$.

Combining (9.11) with Proposition 9.1 we obtain $\int_0^1 |\mathcal{W}_\varepsilon|^2 dz = O(\varepsilon^4)$. But on the other hand (9.10) implies

$$\begin{aligned} \int_0^1 |\mathcal{W}_\varepsilon|^2 dz &= \int_0^1 |\kappa(\varepsilon) H'(\frac{z - z_\varepsilon}{\varepsilon}) \varepsilon + O(\varepsilon^2)|^2 dz \\ &= \varepsilon^3 \int_{-z_\varepsilon/\varepsilon}^{(1-z_\varepsilon)/\varepsilon} |\kappa(\varepsilon) H'(t) + O(\varepsilon)|^2 dt \\ &= \kappa^2(\varepsilon) \varepsilon^3 \int_{-\infty}^{\infty} |H'|^2 dt + O(\varepsilon^4). \end{aligned}$$

Therefore $\kappa(\varepsilon) = O(\varepsilon^{1/2})$. And hence $\mathcal{W}_\varepsilon = O(\varepsilon^{3/2})$ by (9.10).

Finally we revisit the first two lines of (9.11), which imply

$$\begin{aligned} & \int_0^1 [\varepsilon^2 |\mathcal{W}_\varepsilon'|^2 + f'(\mathcal{U}_\varepsilon) \mathcal{W}_\varepsilon^2 + \varepsilon d |\mathcal{Z}_\varepsilon'|^2] dz \leq C \int_0^1 |\mathcal{W}_\varepsilon|^3 dz \\ & \leq C \|\mathcal{W}_\varepsilon\|_{L^\infty(0,1)} \int_0^1 \mathcal{W}_\varepsilon^2 dz \leq C \varepsilon^{3/2} \int_0^1 \mathcal{W}_\varepsilon^2 dz. \end{aligned}$$

Proposition 9.1 yields $c_4\varepsilon \int_0^1 \mathcal{W}_\varepsilon^2 dz \leq C\varepsilon^{3/2} \int_0^1 \mathcal{W}_\varepsilon^2 dz$. Thus $\mathcal{W}_\varepsilon = 0$. \square

We return to the parameters ε and l . Rename \mathcal{U}_ε , the unique minimum in B_δ , $\mathcal{U}_{\varepsilon,l}$. The non-degeneracy implied by Proposition 9.1 allows us to apply the implicit function theorem to conclude that $\mathcal{U}_{\varepsilon,l}$ is differentiable in l under the $W^{1,2}$ norm. Let $\mathcal{W}_{\varepsilon,l} = \frac{\partial \mathcal{U}_{\varepsilon,l}}{\partial l}$.

Proposition 9.3

$$\mathcal{W}_{\varepsilon,l} = H' \left(\frac{l(z-z_\varepsilon)}{\varepsilon} \right) \frac{z-z_\varepsilon}{\varepsilon} - \text{Ave} \left(H' \left(\frac{l(z-z_\varepsilon)}{\varepsilon} \right) \frac{z-z_\varepsilon}{\varepsilon} \right) + \phi,$$

with $\|\phi\|_2 = O(1/l)$. And

$$\phi = cH' \left(\frac{l(z-z_\varepsilon)}{\varepsilon} \right) \frac{l}{\varepsilon} + \phi^\perp,$$

with $\int_0^1 H' \left(\frac{l(z-z_\varepsilon)}{\varepsilon} \right) \phi^\perp dz = 0$, $c = O(1)$ and $\|\phi^\perp\|_2 = O(l)$.

Proof. Differentiate (8.6) with respect to l to obtain the equation

$$-\left(\frac{\varepsilon}{l}\right)^2 \mathcal{W}_{\varepsilon,l}'' + f'(\mathcal{U}_{\varepsilon,l}) \mathcal{W}_{\varepsilon,l} + l^2 (-D^2)^{-1} \mathcal{W}_{\varepsilon,l} + \frac{2}{l} f(\mathcal{U}_{\varepsilon,l}) + 4l \mathcal{V}_{\varepsilon,l} - \frac{2\lambda_\varepsilon}{l} = \lambda_l, \quad (9.12)$$

for $\mathcal{W}_{\varepsilon,l}$, where λ_l is the derivative of λ_ε with respect to l .

As in the proof of Proposition 9.1 we replace H by H_ε . Define $g_\varepsilon = H'_\varepsilon \left(\frac{l(z-z_\varepsilon)}{\varepsilon} \right) \frac{z-z_\varepsilon}{\varepsilon}$, and $\varphi = \mathcal{W}_{\varepsilon,l} - (g_\varepsilon - \text{Ave}(g_\varepsilon))$. g_ε satisfies the equation

$$-\left(\frac{\varepsilon}{l}\right)^2 g_\varepsilon'' + f'(H) g_\varepsilon + \frac{2}{l} f(H) = O(e^{-C/\varepsilon}).$$

Subtract this from (9.12) and use the fact $\text{Ave}(g_\varepsilon) = O(l)$ to deduce the equation for φ :

$$-\left(\frac{\varepsilon}{l}\right)^2 \varphi'' + f'(\mathcal{U}_{\varepsilon,l}) \varphi + l^2 (-D^2)^{-1} \mathcal{W}_{\varepsilon,l} + O(l) = \lambda_l. \quad (9.13)$$

After defining $A = (-D^2)^{-1}(g_\varepsilon - \text{Ave}(g_\varepsilon))$ and $B = (-D^2)^{-1}(\varphi - \text{Ave}(\varphi))$, we multiply the above equation by φ and integrate by parts.

$$\int_0^1 \left[\left(\frac{\varepsilon}{l}\right)^2 |\varphi'|^2 + f'(\mathcal{U}_{\varepsilon,l}) \varphi^2 + l^2 \varphi(A+B) + \varphi O(l) \right] dz = 0.$$

Note that $\int_0^1 \varphi B dz = \int_0^1 |B|^2 dz$ and $A = O(1/l)$. By Proposition 9.1 we find

$$c_7 l^2 \int_0^1 \varphi^2 dz \leq \int_0^1 \left[\left(\frac{\varepsilon}{l}\right)^2 |\varphi'|^2 + f'(\mathcal{U}_{\varepsilon,l}) \varphi^2 + l^2 \varphi B \right] dz \leq \int_0^1 |\varphi| dz O(l).$$

Hence

$$\|\varphi\|_2 = O(1/l). \quad (9.14)$$

Since $\|g_\varepsilon\|_2 = O(1/l)$, we conclude that $\|\mathcal{W}_{\varepsilon,l}\|_2 = O(1/l)$. Hence $A + B = O(1/l)$. This simplifies (9.13) to

$$-\left(\frac{\varepsilon}{l}\right)^2 \varphi'' + f'(\mathcal{U}_{\varepsilon,l})\varphi + O(l) = \lambda_l. \quad (9.15)$$

Multiply this by $h_\varepsilon = \frac{l}{\varepsilon} H'_\varepsilon\left(\frac{l(z-z_\varepsilon)}{\varepsilon}\right)$ (as in the proof of Proposition 9.1), and integrate by parts:

$$\int_0^1 [-\varphi h_\varepsilon'' + \varphi f'(\mathcal{U}_{\varepsilon,l})h_\varepsilon] dz = 2(\lambda_l - O(l)).$$

Thus by (9.14) and (9.6)

$$2(\lambda_l - O(l)) \leq \|\varphi\|_2 \|O(e^{-C/\varepsilon}) + (f'(\mathcal{U}_{\varepsilon,l}) - f'(H))h_\varepsilon\|_2 = O(1/l)O(l^3) = O(l^2),$$

which implies

$$\lambda_l = O(l). \quad (9.16)$$

The equation (9.15) becomes

$$-\left(\frac{\varepsilon}{l}\right)^2 \varphi'' + f'(\mathcal{U}_{\varepsilon,l})\varphi = O(l). \quad (9.17)$$

We decompose $\varphi = ch_\varepsilon + \varphi^\perp$, $\int_0^1 h_\varepsilon \varphi^\perp dz = 0$. By (9.14)

$$c = \frac{\int_0^1 \varphi h_\varepsilon dz}{\|h_\varepsilon\|_2^2} \leq \frac{\|\varphi\|_2}{\|h_\varepsilon\|_2} = O(1).$$

φ^\perp satisfies the equation

$$-\left(\frac{\varepsilon}{l}\right)^2 (\varphi^\perp)'' + f'(\mathcal{U}_{\varepsilon,l})\varphi^\perp + c(f'(\mathcal{U}_{\varepsilon,l}) - f'(H))h_\varepsilon = O(l).$$

However similar to the argument leading to (9.6)

$$\begin{aligned} |(f'(\mathcal{U}_{\varepsilon,l}) - f'(H))h_\varepsilon| &\leq C|\varepsilon\phi_0(z) + O(\varepsilon^2)||h_\varepsilon(z)| \\ &= C\varepsilon \left| \frac{\phi_0(\varepsilon t + z_\varepsilon) - \phi_0((1-m)/2)}{\varepsilon} + O(1) \right| \cdot |H'_\varepsilon(t)| \\ &= C\varepsilon |t| + O(1) \cdot |H'_\varepsilon(t)| \\ &= O(\varepsilon). \end{aligned}$$

So the equation for φ^\perp is

$$-\left(\frac{\varepsilon}{l}\right)^2 (\varphi^\perp)'' + f'(\mathcal{U}_{\varepsilon,l})\varphi^\perp = O(l).$$

Multiply this equation by φ^\perp , integrate by parts, and use (9.2) to find $\|\varphi^\perp\|_2 = O(l)$. \square

10 The convexity of E and E/l

As suggested in (8.1) we define

$$E(\epsilon, l) = \inf\{lJ_{\epsilon, l}(\mathcal{U}) : \mathcal{U} \in B_\delta\} = lJ_{\epsilon, l}(\mathcal{U}_{\epsilon, l}).$$

Through reversal we observe

$$E(\epsilon, l) = \inf\{lJ_{\epsilon, l}(\mathcal{U}) : \mathcal{U} \in B_\delta^R\} = lJ_{\epsilon, l}(\mathcal{U}_{\epsilon, l}^R),$$

where $\mathcal{U}_{\epsilon, l}^R$ is the reversal of $\mathcal{U}_{\epsilon, l}$.

Proposition 10.1 *In the range (8.3) both E and E/l are strictly convex with respect to l . More precisely*

$$\begin{aligned} \frac{\partial^2 E}{\partial l^2} &= \frac{(1-m^2)^2}{4}l + O(\epsilon^{2/3}), \\ \frac{\partial^2}{\partial l^2}\left(\frac{E}{l}\right) &= 2c_0\left(\frac{\epsilon}{l^3}\right) + \frac{(1-m^2)^2}{12} + O(\epsilon^{1/3}). \end{aligned}$$

Proof. Multiplying (8.6) by $\mathcal{U}_{\epsilon, l} - m$ and integrating by parts, we find the useful integral identity

$$\int_0^1 \left[\left(\frac{\epsilon}{l}\right)^2 |\mathcal{U}'_{\epsilon, l}|^2 + f(\mathcal{U}_{\epsilon, l})(\mathcal{U}_{\epsilon, l} - m) + l^2 |\mathcal{V}'_{\epsilon, l}|^2 \right] dz = 0, \quad (10.1)$$

where $\mathcal{V}_{\epsilon, l} = (-D^2)^{-1}(\mathcal{U}_{\epsilon, l} - m)$. This identity and Lemma 8.4 turn E to

$$\begin{aligned} E(\epsilon, l) &= l \int_0^1 \left[W(\mathcal{U}_{\epsilon, l}) - \frac{f(\mathcal{U}_{\epsilon, l})(\mathcal{U}_{\epsilon, l} - m)}{2} \right] dz \\ &= \epsilon \int_{-\infty}^{\infty} \left(W(H) - \frac{f(H)(H - m)}{2} \right) dt + \\ &\quad \epsilon \int_0^1 \left(f(\pm 1) - \frac{f'(\pm 1)(\pm 1 - m) + f(\pm 1)}{2} \right) \phi_0 dz + O(\epsilon^{5/3}) \\ &= c_0 \epsilon + \frac{(1-m^2)^2}{24} l^3 + O(\epsilon^{5/3}). \end{aligned} \quad (10.2)$$

The integral in the second line is c_0 because of (2.3) and (2.4). The computation of the integral in the third line uses the definition (8.10) of ϕ_0 and the expression (3.4) of \mathcal{V}_0 .

Differentiating E with respect to l yields

$$\frac{\partial E}{\partial l} = \int_0^1 \left[-\frac{\epsilon^2}{2l^2} |\mathcal{U}'_{\epsilon, l}|^2 + W(\mathcal{U}_{\epsilon, l}) + \frac{3l^2}{2} |\mathcal{V}'_{\epsilon, l}|^2 \right] dz. \quad (10.3)$$

We have used the fact that $\mathcal{U}_{\epsilon,l}$ is a critical point of $J_{\epsilon,l}$. By (10.1) and Lemma 8.4 this becomes

$$\begin{aligned}\frac{\partial E}{\partial l} &= \int_0^1 [W(\mathcal{U}_{\epsilon,l}) + \frac{f(\mathcal{U}_{\epsilon,l})(\mathcal{U}_{\epsilon,l} - m)}{2}] dz + \int_0^1 2l^2 |\mathcal{V}'_{\epsilon,l}|^2 dz \\ &= \frac{\epsilon}{l} \int_{-\infty}^{\infty} [W(H) + \frac{f(H)(H - m)}{2}] dt + \frac{\epsilon}{l} \int_0^1 \frac{f'(\pm 1)(\pm 1 - m)}{2} \phi_0 dz + \\ &\quad \int_0^1 2l^2 |\mathcal{V}'_{\epsilon,l}|^2 dz + O(\epsilon^{4/3})\end{aligned}$$

The first integral in the second line is 0 again by (2.3) and (2.4). The second integral is $-(1-m^2)^2/24$ multiplied by l^2/ϵ as in the estimate of E . To estimate the integral in the last line note (8.17), so using (3.4) we deduce

$$\int_0^1 2l^2 |\mathcal{V}'_{\epsilon,l}|^2 dz = \int_0^1 2l^2 |\mathcal{V}'_0|^2 dz + O(\epsilon^{4/3}) = \frac{(1-m^2)^2}{6} l^2 + O(\epsilon^{4/3}).$$

Altogether

$$\frac{\partial E}{\partial l} = \frac{(1-m^2)^2}{8} l^2 + O(\epsilon^{4/3}). \quad (10.4)$$

Differentiate (10.3) with respect to l . Denote the derivative of $\mathcal{U}_{\epsilon,l}$ with respect to l by $\mathcal{W}_{\epsilon,l}$ and the derivative of $\mathcal{V}_{\epsilon,l}$ with respect to l by $\mathcal{Z}_{\epsilon,l}$. Then

$$\frac{\partial^2 E}{\partial l^2} = \int_0^1 [\frac{\epsilon^2}{l^3} |\mathcal{U}'_{\epsilon,l}|^2 + 3l |\mathcal{V}'_{\epsilon,l}|^2] dz + \int_0^1 [-\frac{\epsilon^2}{l^2} \mathcal{U}'_{\epsilon,l} \mathcal{W}'_{\epsilon,l} + f(\mathcal{U}_{\epsilon,l}) \mathcal{W}_{\epsilon,l} + 3l^2 \mathcal{V}'_{\epsilon,l} \mathcal{Z}'_{\epsilon,l}] dz.$$

Call the first term on the right T_1 and the second term T_2 . The estimation of T_1 is similar to the earlier ones. Using (10.1) we find

$$\begin{aligned}T_1 &= \frac{1}{l} \int_0^1 [-f(\mathcal{U}_{\epsilon,l})(\mathcal{U}_{\epsilon,l} - m) + 2l^2 |\mathcal{V}'_{\epsilon,l}|^2] dz \\ &= \frac{\epsilon}{l^2} \int_{-\infty}^{\infty} -f(H)(H - m) dt + \frac{(1-m^2)^2}{12} l + \frac{(1-m^2)^2}{6} l + O(\epsilon) \\ &= \frac{\epsilon}{l^2} \int_{-\infty}^{\infty} -f(H)H dt + \frac{(1-m^2)^2}{4} l + O(\epsilon).\end{aligned}$$

To estimate T_2 , first use (8.6) and $\mathcal{Z}_{\epsilon,l} = (-D^2)^{-1} \mathcal{W}_{\epsilon,l}$ to simplify it to

$$T_2 = \int_0^1 [2f(\mathcal{U}_{\epsilon,l}) \mathcal{W}_{\epsilon,l} + 4l^2 \mathcal{V}_{\epsilon,l} \mathcal{W}_{\epsilon,l}] dz.$$

By Propositions 8.3 and 9.3

$$\begin{aligned}T_2 &= \int_0^1 [2f(H) + O(\epsilon^{2/3})] (\frac{z - z_\epsilon}{\epsilon} H' + \phi^\perp + O(\epsilon^{1/3})) dz \\ &= \frac{\epsilon}{l^2} \int_{-z_\epsilon/\epsilon}^{(1-z_\epsilon)/\epsilon} 2f(H(t)) H'(t) t dt + O(\epsilon^{2/3}) \\ &= \frac{\epsilon}{l^2} \int_{-\infty}^{\infty} -2W(H) dt + O(\epsilon^{2/3}).\end{aligned}$$

We have used the estimates

$$\begin{aligned}\int_0^1 \left| \frac{z - z_\epsilon}{\epsilon} H' \right| dz &= \frac{\epsilon}{l^2} \int_{-z_\epsilon/\epsilon}^{(1-z_\epsilon)/\epsilon} |H'(t)t| dt = O(\epsilon^{1/3}), \\ \int_0^1 |f(H)| dz &= \frac{\epsilon}{l} \int_{-z_\epsilon/\epsilon}^{(1-z_\epsilon)/\epsilon} |f(H(t))| dt = O(\epsilon^{2/3}) \\ \|2f(H) + O(\epsilon^{2/3})\|_2 &= O(\epsilon^{1/3}).\end{aligned}$$

Adding T_1 and T_2 , since $\int_{-\infty}^{\infty} (f(H)H + 2W(H)) dt = 0$ as in the estimation for $\partial E/\partial l$, we arrive at

$$\frac{\partial^2 E}{\partial l^2} = \frac{(1-m^2)^2}{4} l + O(\epsilon^{2/3}), \quad (10.5)$$

proving the estimate for $\partial^2 E/\partial l^2$. From (10.2), (10.4) and (10.5), we deduce

$$\frac{\partial^2}{\partial l^2} \left(\frac{E}{l} \right) = \frac{\frac{\partial^2 E}{\partial l^2} l^2 - 2 \frac{\partial E}{\partial l} l + 2E}{l^3} = 2c_0 \left(\frac{\epsilon}{l^3} \right) + \frac{(1-m^2)^2}{12} + O(\epsilon^{1/3}). \quad \square$$

We now prove the three theorems stated in Section 1. Recall that a global minimum of I_ϵ is denoted by u_ϵ , with N_ϵ α -points, denoted by $x_1, x_2, \dots, x_{N_\epsilon}$. Between them there are $N_\epsilon - 1$ zeros of the derivative of $v_\epsilon = (-D^2)^{-1}(u_\epsilon - m)$, denoted by $y_1, y_2, \dots, y_{N_\epsilon-1}$, satisfying $0 < x_1 < y_1 < x_2 < y_2 < \dots < x_{N_\epsilon-1} < y_{N_\epsilon-1} < x_{N_\epsilon} < 1$. We set $l_i = y_i - y_{i-1}$ for $i = 1, 2, \dots, N_\epsilon$ with $y_0 = 0$ and $y_{N_\epsilon} = 1$. There are two possibilities for u_ϵ on $(0, x_1)$: $u_\epsilon > \alpha$ or $u_\epsilon < \alpha$.

Proof of Theorem 1.1. Without the loss of generality we suppose that $u_\epsilon > \alpha$ on $(0, x_1)$. We construct a particular periodic solution u_ϵ^* with N_ϵ α -points (i.e. $N_\epsilon/2$ periods), and show that $u_\epsilon = u_\epsilon^*$.

Let $\mathcal{U}_{\epsilon, 1/N_\epsilon}$ be the unique minimum of $J_{\epsilon, l}$ in B_δ (Proposition 9.2), with $l = 1/N_\epsilon$, and $\mathcal{U}_{\epsilon, 1/N_\epsilon}^R$, its reversal, be the unique minimum of $J_{\epsilon, 1/N_\epsilon}$ in B_δ^R . Set $u_\epsilon^*(x) = \mathcal{U}_{\epsilon, 1/N_\epsilon}^R(N_\epsilon x)$ for $x \in (0, 1/N_\epsilon)$. Extend u_ϵ^* anti-periodically to $(0, 1)$, i.e. $u_\epsilon^*(x) = \mathcal{U}_{\epsilon, 1/N_\epsilon}(N_\epsilon x - 1)$ for $x \in (1/N_\epsilon, 2/N_\epsilon)$, $u_\epsilon^*(x) = \mathcal{U}_{\epsilon, 1/N_\epsilon}^R(N_\epsilon x - 2)$ for $x \in (2/N_\epsilon, 3/N_\epsilon)$... Clearly u_ϵ^* is periodic with $N_\epsilon/2$ periods.

For small ϵ by Lemma 4.6, Propositions 7.1 and 7.2, $u_\epsilon(l_i \cdot + y_{i-1}) \in B_\delta$ when i is even and $\in B_\delta^R$ when i is odd. Using the strict convexity of E in Proposition 10.1 and (8.1) we find

$$I_\epsilon(u_\epsilon^*) \geq I_\epsilon(u_\epsilon) \geq \sum_{i=1}^{N_\epsilon} l_i J_{\epsilon, l_i}(u_\epsilon(l_i \cdot + y_{i-1})) \geq \sum_{i=1}^{N_\epsilon} E(\epsilon, l_i) \geq N_\epsilon E(\epsilon, \frac{1}{N_\epsilon}) = I_\epsilon(u_\epsilon^*).$$

All the inequalities above must be equalities. Therefore $l_i = 1/N_\epsilon$ for all i , and $u_\epsilon(l_i \cdot + y_{i-1}) = \mathcal{U}_{\epsilon, 1/N_\epsilon}$ when i is even or $= \mathcal{U}_{\epsilon, 1/N_\epsilon}^R$ when i is odd by Proposition 9.2. Thus $u_\epsilon = u_\epsilon^*$.

If on $(0, x_1)$ $u_\epsilon < \alpha$, u_ϵ must be the reversal of u_ϵ^* . \square

Proof of Theorem 1.2. In the previous proof we have shown that if N_ϵ is known, there are exactly two global minima of I_ϵ , u_ϵ^* and its reversal, with N_ϵ α -points. Here we determine whether N_ϵ is unique.

By the strict convexity of E/l (Proposition 10.1), E/l attains its minimum at a unique l_* . But for $I_\epsilon(u_\epsilon) = N_\epsilon E(\epsilon, 1/N_\epsilon)$, its minimum with respect to N_ϵ is achieved at one or two integers.

If $1/l_*$ happens to be an integer, then there is only one $N_\epsilon = 1/l_*$. If $1/l_*$ is not an integer, there exist two consecutive integers, say N and $N + 1$, such that $N < 1/l_* < N + 1$. If $NE(\epsilon, 1/N) \neq (N + 1)E(\epsilon, 1/(N + 1))$, then again there is only one N_ϵ . It must be the one of N and $N + 1$ which offers the smaller of $NE(\epsilon, 1/N)$ and $(N + 1)E(\epsilon, 1/(N + 1))$. In these two case we have two global minima of I_ϵ .

In the less likely third case that $1/l_*$ is not an integer, and $NE(\epsilon, 1/N) = (N + 1)E(\epsilon, 1/(N + 1))$, we have two values, N and $N + 1$, for N_ϵ . Then there are four global minima of I_ϵ . \square

Proof of Theorem 1.3. Collecting the estimates (10.2) and (10.4), we have

$$\frac{\partial}{\partial l} \left(\frac{E}{l} \right) = \frac{\frac{\partial E}{\partial l} l - E}{l^2} = \frac{-c_0 \epsilon + \frac{(1-m^2)^2}{12} l^3 + O(\epsilon^{5/3})}{l^2}.$$

If E/l is minimized at $l = l_*$, the above estimate implies

$$-c_0 \epsilon + \frac{(1-m^2)^2}{12} l_*^3 + O(\epsilon^{5/3}) = 0,$$

which in turn yields

$$l_* = \left(\frac{12c_0}{(1-m^2)^2} \right)^{1/3} \epsilon^{1/3} + O(\epsilon). \quad (10.6)$$

N_ϵ , the number of α -points of u_ϵ , is either $1/l_*$ if it happens to be an integer, or one of the two consecutive integers, N and $N + 1$, such that $1/(N + 1) < l_* < 1/N$. In the first case the theorem is proved since $2l_* = 2/N_\epsilon$ is the period. In the second case since

$$\frac{1}{N} - \frac{1}{N + 1} < \frac{l_*^2}{1 - l_*} = O(\epsilon^{2/3}),$$

the period $2/N$ or $2/(N + 1)$ of u_ϵ is $2l_* + O(\epsilon^{2/3})$, proving the theorem by (10.6). \square

Acknowledgment

Some of the ideas in this paper emerged during our several stays in 1997 and 98 at Mathematisches Forschungsinstitut Oberwolfach, which we would like to thank for its hospitality. The first author also thanks the Department of Mathematics and the Institute of Mathematical Sciences at the Chinese University of Hong Kong where he spent several crucial weeks during the course of this work.

References

- [1] J. Carr, M.E. Gurtin and M. Slemrod, *Structured phase transitions on a finite interval*, Arch. Rational Mech. Anal. 86 (1984), no. 4, 317-351.
- [2] R. Choksi, *Scaling laws in microphase separation of diblock copolymers*, J. Nonlinear Sci. 11 (2001), 223-236.
- [3] P.C. Fife and D. Hilhorst, *The Nishiura-Ohnishi free boundary problem in the 1D case*, SIAM J. Math. Anal. 33 (2001), 589-606.
- [4] M. Henry, *Singular limit of a fourth order problem arising in the microphase separation of diblock copolymers*, Adv. Differential Equations 6 (2001), no. 9, 1049-1114.
- [5] C.B. Muratov, *Theory of domain patterns in systems with long-range interactions of Coulomb type*, preprint.
- [6] S. Müller, *Singular perturbations as a selection criterion for periodic minimizing sequences*, Calc. Var. Partial Differential Equations 1 (1993), no. 2, 169-204.
- [7] W.-M. Ni and I. Takagi, *On the shape of least energy solution to a semilinear Neumann problem*, Comm. Pure Appl. Math. 41 (1991), 819-851.
- [8] W.-M. Ni and I. Takagi, *Locating the peaks of least energy solutions to a semilinear Neumann problem*, Duke Math. J. 70 (1993), 247-281.
- [9] W.-M. Ni and J. Wei, *On the location and profile of spike-layer solutions to singularly perturbed semilinear Dirichlet problems*, Comm. Pure Appl. Math. 48 (1995), 723-761.
- [10] Y. Nishiura and I. Ohnishi *Some mathematical aspects of the microphase separation in diblock copolymers*, Phys. D, 84 (1995), 31-39.
- [11] I. Ohnishi, Y. Nishiura, M. Imai and Y. Matsushita, *Analytical solutions describing the phase separation driven by a free energy functional containing a long-range interaction term*, Chaos 9 (1999), no. 2, 329-341.
- [12] X. Ren and J. Wei, *On the multiplicity of solutions of two nonlocal variational problems*, SIAM J. Math. Anal. 31 (2000), 909-924.
- [13] X. Ren and J. Wei, *Concentrically layered energy equilibria of the di-block copolymer problem*, European J. Appl. Math. 31 (2002), 479-496.
- [14] X. Ren and J. Wei, *Triblock copolymer theory: Free energy, disordered phase and weak segregation*, Physica D, to appear.
- [15] X. Ren and J. Wei, *Triblock copolymer theory: Ordered ABC lamellar phase*, J. Nonlinear Sci., to appear.

- [16] X. Ren and J. Wei. *On the spectra of 3-D lamellar solutions of the diblock copolymer problem*. *SIAM J. Math. Anal.*, to appear.
- [17] X. Ren and J. Wei. *Wriggled lamellar solutions and their stability in the diblock copolymer problem*. preprint.
- [18] T. Ohta and K. Kawasaki, *Equilibrium morphology of block copolymer melts*, *Macromolecules*, 19, 2621-2632 (1986), 2621-2632.
- [19] L. Truskinovsky and G. Zanzotto, *Finite scale microstructures and metastability in one dimensional elasticity*, *Meccanica*, 30 (1995) 577-589.
- [20] L. Truskinovsky and G. Zanzotto, *Ericksen's bar revisited: Energy wiggles*, *J. Mech. Phys. Solids*, 44, 8 (1996) 1371-1408.
- [21] J. Wei, *Conditions for two-peaked solutions of singularly perturbed elliptic equations*, *Manuscripta Math.* 96 (1998), 113-131.