

# STURMIAN EXPANSIONS AND ENTROPY

E. ARTHUR ROBINSON, JR.

## 1. INTRODUCTION

Sturmian sequences were introduced by Morse and Hedlund [10] as the sequences that code the orbits of the geodesic flow on a flat 2-torus. In this paper, we restrict our attention to (1-sided) aperiodic Sturmian sequences, which may be defined to be those sequences  $\mathbf{d} = .d_1d_2d_3 \cdots \in \{0, 1\}^{\mathbb{N}}$  that have exactly  $n + 1$  distinct *factors* (subsequences  $\mathbf{u} = d_jd_{j+1} \cdots d_{j+n-1}$ ) of length  $n$ . This property is often expressed by saying that a Sturmian sequence  $\mathbf{d}$  has *complexity function*  $c_{\mathbf{d}}(n) = n + 1$ . If  $c_{\mathbf{e}}(n)$  is the complexity function of a sequence  $\mathbf{e} \in \{0, 1\}^{\mathbb{N}}$ , it is known (see [5], Chapter 6) that  $c_{\mathbf{e}}(k) = k$  for some  $k$  if and only if  $\mathbf{e}$  is eventually periodic. Thus Sturmian sequences are the least complex among aperiodic sequences.

A sequence  $\mathbf{d} = .d_1d_2d_3 \cdots \in \{0, 1\}^{\mathbb{N}}$  is said to be *balanced* if for any  $i, j, \ell \geq 1$

$$\left| \sum_{k=i}^{i+\ell-1} d_k - \sum_{k=j}^{j+\ell-1} d_k \right| \leq 1.$$

It can be shown that a sequence is balanced if and only if it is Sturmian, and from this, one can prove (see [5], Chapter 6) that the limit

$$(1) \quad \alpha = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^{\infty} d_k$$

exists, and  $\alpha$  is irrational. The number  $\alpha$  is called the *slope* of  $\mathbf{d}$ .

Morse and Hedlund [10] showed that if  $\mathbf{d} = .d_1d_2d_3 \cdots$  is a Sturmian sequence with slope  $\alpha \in (0, 1) \setminus \mathbb{Q}$ , then there is a unique  $x \in [0, 1)$ , called the *intercept*, so that  $\mathbf{d}$  either has the form

$$(2) \quad d_n = \lfloor \alpha(n+1) + x \rfloor - \lfloor \alpha n + x \rfloor,$$

for all  $n \in \mathbb{N}$ , or

$$(3) \quad d_n = \lceil \alpha(n+1) + x \rceil - \lceil \alpha n + x \rceil.$$

Note that (2) and (3) are the same unless  $n\alpha + x = 0 \pmod{1}$  for some  $n > 1$ , in which case they disagree in exactly one or two adjacent digits.

Given a Sturmian sequence  $\mathbf{d}$ , one can easily determine its slope  $\alpha$  using (1). The goal of this paper is to exhibit a similarly simple formula for the intercept  $x$ . In particular, the intercept  $x$  can be obtained using a well know generalization of continued fraction and radix expansions, called an *f*-expansions. Another way to say this is that a Sturmian sequence  $\mathbf{d} = .d_1d_2d_2 \cdots \in \{0, 1\}^{\mathbb{N}}$  can be regarded

*Date:* October 15, 2010.

*2010 Mathematics Subject Classification.* 11K16, 28D20, 37E05, 37B10.

*Key words and phrases.* numeration, entropy, interval maps, symbolic dynamics.

as type “binary expansion” of its intercept  $x$ . We refer to this as the *Sturmian  $\alpha$  expansion* of  $x$ .

After a brief discussion of  $f$ -expansions in general, we discuss general some properties of Sturmian  $\alpha$ -expansions. In particular, Sturmian  $\alpha$ -expansions differs significantly from nearly all other the familiar numeration systems, including continued fraction expansions and  $\beta$ -expansions — two examples we use to draw this contrast. We conclude by mentioning a few other examples that have properties similar to Sturmian  $\alpha$ -expansions.

## 2. $f$ -EXPANSIONS

Let  $f : \mathbb{R} \rightarrow [0, 1]$  be a continuous monotonic function with  $\overline{f(\mathbb{R})} = [0, 1]$ . An  $f$ -*expansion* is an expression of the form

$$(4) \quad x = f(d_1 + f(d_2 + f(d_3 + \dots))),$$

where the *digits*  $d_k$  are integers. We call  $\mathbf{d} = .d_1d_2d_3\dots$  the *digit sequence* of the expansion (4). In particular, the expression (4) means that  $x_n \rightarrow x$ , where

$$(5) \quad x_n = f(d_1 + f(d_2 + \dots + f(d_n))).$$

This idea goes back to Kakeya [7], who observed in 1926 that examples of  $f$ -expansions include<sup>1</sup> both regular continued fractions and base  $\beta$  radix expansions  $\beta > 1$ . In particular, regular continued fractions

$$x = f(d_1 + f(d_2 + f(d_3 + \dots))) = \frac{1}{d_1 + \frac{1}{d_2 + \frac{1}{d_3 + \dots}}},$$

correspond to the case  $f(x) = 1/x$ , whereas base- $\beta$  radix expansions

$$x = f(d_1 + f(d_2 + f(d_3 + \dots))) = \frac{d_1 + \frac{d_2 + \frac{d_3 + \dots}{\beta}}{\beta}}{\beta} = \sum_{k=1}^{\infty} \frac{d_k}{\beta^k}$$

correspond to  $f(x) = x/\beta$ .

Although more than one digit sequence in (4) may yield the same number  $x \in [0, 1)$  (just as  $0.099\dots = 0.100\dots$  in base 10), there is a standard algorithm that takes  $x$  and produces a digit particular digit sequence  $\mathbf{d} = .d_1d_2d_3\dots$  that we call the *proper  $f$ -expansion of  $x$* . As Rényi [14] observed in 1951, this algorithm is perhaps best described in terms of a dynamical system. Using the given function  $f$ , we define an almost everywhere map  $T : [0, 1) \rightarrow [0, 1)$ , called the  *$f$ -transformation*, by

$$(6) \quad Tx = f^{-1}(x) \bmod 1.$$

We also define a *labeled interval partition* a.e. on  $[0, 1)$ , by which simply mean an integer valued function on  $[0, 1)$  defined  $\xi(x) = \lfloor f^{-1}(x) \rfloor$ . The “interval partition” here is the partition of  $[0, 1)$  into the nonempty intervals  $\Delta(d) = [a, b) = \{x : \xi(x) = d\}$  where the function  $\xi$  is constant. These sets  $\Delta(d)$  are called *fundamental*

<sup>1</sup>The same observation was made independently by Bissinger [2] for  $f$  increasing and Everett [4] for  $f$  decreasing.

intervals. Assuming  $x$  is such that  $T^{n-1}x$  exists for all  $n \in \mathbb{N}$ , the proper digit sequence  $\mathbf{d} = .d_1d_2d_3\dots$  is defined by

$$(7) \quad d_n = \xi(T^{n-1}x), \quad n \in \mathbb{N}.$$

If the proper digit sequence  $\mathbf{d} = .d_1d_2d_3\dots$  is used in (4) we call it the *proper  $f$ -expansion* of  $x$ . Note that  $d_n \in \mathcal{D}$  where  $\mathcal{D} := \xi([0, 1))$  is called the *digit set*.

For continued fractions, the  $f$ -transformation is the *Gauss map*  $Tx = 1/x \bmod 1$ , and  $\mathcal{D} = \mathbb{N}$ . (In cases like this, where some of the digits  $d_n$  are multi-digit numbers when written in base 10, it will sometimes be convenient to write the digit sequence as  $\mathbf{d} = [d_1, d_2, d_3, \dots]$  rather than  $\mathbf{d} = .d_1d_2d_3\dots$ .) For base- $\beta$  radix expansions, the  $f$ -transformation  $Tx = \beta x \bmod 1$  is called the  $\beta$ -*transformation*, and  $\mathcal{D} = \{0, 1, \dots, [\beta] - 1\}$ . The case  $\beta \in \mathbb{N}$  gives the usual radix expansions (e.g., base 2, base 10).

We say  $f$ -expansions have *unique proper digits* if the proper digit sequence map  $x \mapsto \mathbf{d}$  is injective, and we say  $f$ -expansions are *valid* if for each  $x$ , such that  $T^n x$  exists for all  $n \geq 0$ , the proper  $f$ -expansion converges to  $x$ . A typical approach to this problem is the following (see [7] and also [17]).

**Theorem 1** (Kakeya's theorem). *Assume  $f$  is strictly monotone on an interval  $(a, b) \subseteq \mathbb{R}$  with*

$$(8) \quad \begin{aligned} -\infty \leq a < a + 1 < b \leq +\infty, \\ \text{and} \\ \overline{f((a, b))} = [0, 1]. \end{aligned}$$

*If the  $f$ -transformation  $T$  satisfies*

$$(9) \quad |T'(x)| > 1 \quad \text{a.e.},$$

*then  $f$  has unique proper digits and  $f$ -expansions are valid.*

Note that (9) is equivalent to

$$(10) \quad |f'(x)| < 1 \quad \text{a.e. on } (a, b).$$

Similar results due to Bissinger [2] and Everett [4] require that  $f$  satisfies a Lipschitz condition with constant  $K < 1$  instead of (10).

### 3. STURMIAN $\alpha$ -EXPANSIONS

Let us fix an irrational number  $\alpha \in [0, 1) \setminus \mathbb{Q}$  and consider as the *irrational rotation transformation*  $Tx = x + \alpha \bmod 1$ . This can be interpreted as the  $f$ -transformation (6) for the function

$$(11) \quad f(x) = \begin{cases} 0 & \text{if } x < \alpha \\ x - \alpha & \text{if } \alpha \leq x \leq \alpha + 1 \\ 1 & \text{if } x > \alpha + 1. \end{cases}$$

The corresponding labeled partition is given by

$$\xi(x) = \begin{cases} 0 & \text{if } 0 \leq x < 1 - \alpha \\ 1 & \text{if } 1 - \alpha < x \leq 1, \end{cases}$$

with digit set  $\mathcal{D} = \{0, 1\}$ .

It is easy to see that for the function  $f$ , defined by (11), the proper digit sequence  $\mathbf{d} = .d_1d_2d_3\dots$  for  $x \in [0, 1)$  is equal to the Sturmian sequence  $\mathbf{d}$  given by (2) with slope  $\alpha$  and intercept  $x$ . These are our *Sturmian  $\alpha$ -expansions*. The irrational rotation transformation  $T$  clearly fails to satisfy Kakeya's hypotheses (9). However, we can still obtain the following result<sup>2</sup>.

**Theorem 2.** *Sturmian  $\alpha$ -expansions have unique proper digit sequences, and are valid.*

*Proof.* Given  $x \in [0, 1)$ , let  $x \mapsto \mathbf{d} = .d_1d_2d_3\dots$  be the proper digit sequence map, and let  $\mathbf{d}_n(x) = d_1d_2\dots d_n$ . Let  $\xi^{(n)}$  be the partition of  $[0, 1)$  into those subintervals  $[a_k^n, b_k^n)$ ,  $k = 1, 2, \dots$ , of  $[0, 1)$ , on which  $\mathbf{d}_n(x)$  is constant. We claim that for each  $n = 1, 2, \dots$ , there are  $|\xi^{(n)}| = n + 1$  such intervals, and if they are arranged so that  $b_{k+1}^n = a_k^n$ , then the *cut points*  $a_1^n, a_2^n, \dots, a_{n+1}^n$  are the first  $n + 1$  points orbit  $O_{T^{-1}}^+(0) = \{T^{-n+1}0 : n \in \mathbb{N}\}$  of 0 under the irrational rotation transformation  $T^{-1}x = x + (1 - \alpha) \bmod 1$ .

The claim is true for  $n = 1$ , so assume it holds for  $n$ . It is well known that  $O_{T^{-1}}^+(0)$  is dense in  $[0, 1)$  (since  $T^{-1}$  is an irrational rotation, see [19]), so  $T^{-(n+1)}0$  is in the interior of  $[a_\ell^n, b_\ell^n)$  for some  $\ell$ . We then have

$$[a_k^{n+1}, b_k^{n+1}) = \begin{cases} [a_k^n, b_k^n) & \text{if } 1 \leq k < \ell, \\ [a_k^n, T^{-(n+1)}0) & \text{if } k = \ell, \\ [T^{-(n+1)}0, b_k^n) & \text{if } k = \ell + 1, \\ [a_{k-1}^n, b_{k-1}^n) & \text{if } \ell + 1 < k \leq n + 1, \end{cases}$$

so the claim holds for  $n + 1$ .

By the claim, the cutpoints of  $\xi^{(n)}$  satisfy  $\{a_1^n, a_2^n, \dots, a_{n+1}^n\} = \{T^{-k}0 : k = 0, \dots, n\}$  for all  $n$ . Let  $|\xi^{(n)}| = \max\{b - a : \Delta = [a, b) \in \xi^{(n)}\}$ . Since  $O_{T^{-1}}^+(x)$  is dense, it follows that  $|\xi^{(n)}| \rightarrow 0$ . This shows Sturmian base- $\alpha$  expansions have unique proper digit sequences.

For  $x \in [0, 1)$  and  $n \in \mathbb{N}$  we have

$$(12) \quad x \in [a^n(x), b^n(x)) := [a_k^n, b_k^n) \in \xi^{(n)}$$

for some unique  $k = k(n)$ . We claim that

$$a^n(x) = f(d_1 + f(d_2 + \dots + f(d_n)))$$

and

$$b^n(x) = f(d_1 + f(d_2 + \dots + f(d_n + 1)))$$

Indeed,  $[f(d_1), f(d_1 + 1)) = [f(0), f(1)) = [0, 1 - \alpha) = [a^1(x), b^1(x))$  if  $d_1 = 0$  and  $[f(d_1), f(d_1 + 1)) = [f(1), f(2)) = [1 - \alpha, 1) = [a^1(x), b^1(x))$  if  $d_1 = 1$ , so the claim holds for  $n = 1$ . We proceed by induction. Note that

$$[a^n(x), b^n(x)) = [a^1(x), b^1(x)) \cap T^{-1}[a^{n-1}(x'), b^{n-1}(x'))$$

where  $x' = Tx$ . By induction,

$$a^{n-1}(x') = f(d_2 + f(d_3 + \dots + f(d_n)))$$

and

$$b^{n-1}(x') = f(d_2 + f(d_3 + \dots + f(d_n + 1))).$$

<sup>2</sup>This fact was noted in passing by Parry in [12].

Thus

$$\begin{aligned} a^n(x) &= T^{-1}(a^{n-1}(x')) \cap [a^1(x), b^1(x)) \\ &= f(d_1 + a^{n-1}(x')) \\ &= f(d_1 + f(d_2 + \cdots + f(d_n))), \end{aligned}$$

and

$$\begin{aligned} b^n(x) &= T^{-1}(b^{n-1}(x')) \cap [a^1(x), b^1(x)) \\ &= f(d_1 + b^{n-1}(x')) \\ &= f(d_1 + f(d_2 + \cdots + f(d_n + 1))). \end{aligned}$$

Finally, since  $\|\xi^{(n)}\| \rightarrow 0$ , it follows that  $x_n = a^n(x) \rightarrow x$ , so Sturmian base- $\alpha$  are valid.  $\square$

As an example, let the base  $\alpha = \sqrt{2} - 1$ . Then for  $x = 1/2$

$$\mathbf{d} = .010100101010010100101001010010100101001010010101000\dots,$$

admittedly, not a very intuitive expansion for  $1/2$ . The first 30 partial convergents (partial  $f$ -expansions) are shown in Table 1. Note that (in general) the convergents

$n$	$x_n$	$\sim x_n$	$n$	$x_n$	$\sim x_n$	$n$	$x_n$	$\sim x_n$
1	0	.00000	11	$16 - 11\sqrt{2}$	.44365	21	$16 - 11\sqrt{2}$	.44365
2	$3 - 2\sqrt{2}$	.17157	12	$16 - 11\sqrt{2}$	.44365	22	$16 - 11\sqrt{2}$	.44365
3	$3 - 2\sqrt{2}$	.17157	13	$16 - 11\sqrt{2}$	.44365	23	$33 - 23\sqrt{2}$	.47309
4	$6 - 4\sqrt{2}$	.34315	14	$16 - 11\sqrt{2}$	.44365	24	$33 - 23\sqrt{2}$	.47309
5	$6 - 4\sqrt{2}$	.34315	15	$16 - 11\sqrt{2}$	.44365	25	$33 - 23\sqrt{2}$	.47309
6	$6 - 4\sqrt{2}$	.34315	16	$16 - 11\sqrt{2}$	.44365	26	$33 - 23\sqrt{2}$	.47309
7	$6 - 4\sqrt{2}$	.34315	17	$16 - 11\sqrt{2}$	.44365	27	$33 - 23\sqrt{2}$	.47309
8	$6 - 4\sqrt{2}$	.34315	18	$16 - 11\sqrt{2}$	.44365	28	$33 - 23\sqrt{2}$	.47309
9	$6 - 4\sqrt{2}$	.34315	19	$16 - 11\sqrt{2}$	.44365	29	$33 - 23\sqrt{2}$	.47309
10	$6 - 4\sqrt{2}$	.34315	20	$16 - 11\sqrt{2}$	.44365	30	$33 - 23\sqrt{2}$	.47309

TABLE 1. First 20 convergents of  $x = 1/2 = .5$  in the Sturmian expansion base  $\alpha = \sqrt{2} - 1$ .

$x_n$  lie in the set  $\mathbb{Z} + \alpha\mathbb{Z}$  ( $\mathbb{Z}[\sqrt{2}]$  in the example). Table 1 suggests that convergence is very slow. This is reinforced by Figure 1, which shows a plot of the first 1000 convergents. Note that  $x_{1000} = 1105 - 781\sqrt{2} \sim .49921$ , still correct to only three decimal places. Figure 1 and Table 1 both suggest that there are long intervals of  $n$  where the convergents  $x_n$  remain constant.

As a second example, again for  $\alpha = \sqrt{2} - 1$ , let  $x = 16 - 11\sqrt{2} \sim .44365$ . Then we get

$$\mathbf{d} = .010100101010010100101001010010100101001010010101001010010100101\dots$$

A calculation shows that  $x_1 = 0$ ,  $x_2 = x_3 = 3 - 2\sqrt{2}$ ,  $x_4 = \cdots = x_9 = 6 - 4\sqrt{2}$  and  $x_{10} = x_{11} = x_{12} = \cdots = 16 - 11\sqrt{2}$ . So the convergents reach  $x$  after a finite number of steps (and never change) even though the representation  $\mathbf{d}$  is infinite. These facts are explained by the following proposition.




FIGURE 1. A listplot of the first 1000 convergents of  $x = 1/2 = .5$  in the Sturmian base  $\alpha = \sqrt{2} - 1$ .

**Proposition 3.** Fix  $\alpha \in [0, 1) \setminus \mathbb{Q}$ , and for  $x \in [0, 1)$ , define

$$\underline{x}_n = \min((\{0\} \cup \{T^{-k}0 : k = 1, \dots, n\}) \cap [0, x])$$

and

$$\bar{x}_n = \min((\{T^{-k}0 : k = 1, \dots, n\} \cup \{1\}) \cap (x, 1]),$$

so that  $x \in [\underline{x}_n, \bar{x}_n)$  for all  $n \in \mathbb{N}$ . Then there exist strictly increasing sequences  $\underline{n}_k, \bar{n}_k \in \mathbb{N}$  so that

$$\underline{x}_n = T^{-\underline{n}_k}0 \text{ for } n \in [\underline{n}_k, \underline{n}_{k+1})$$

and

$$\bar{x}_n = T^{-\bar{n}_k}0 \text{ for } n \in [\bar{n}_k, \bar{n}_{k+1}).$$

Moreover, if  $a^n(x)$  and  $b^n(x)$  are defined as in (12), then  $[a^n(x), b^n(x)] = [\underline{x}_n, \bar{x}_n)$ .

Note that  $x_n = \underline{x}_n$ . We are now in a position to give a qualitative description of the long intervals in  $\mathbb{N}$  on which the convergents are constant. By Proposition 3, these are the intervals  $[\underline{n}_k, \underline{n}_{k+1})$ . At step  $\underline{n}_k$  there are  $\underline{n}_k + 1$  intervals in  $\xi^{(\underline{n}_k)}$ , one of which,  $[a_{\underline{x}^k}^{\underline{n}_k}, b_{\underline{x}^k}^{\underline{n}_k})$  contains  $x$ . Thus  $O_{T^{-1}}^+(x)$  will visit every other interval in  $\xi^{(\underline{n}_k)}$  at least once before its first return to  $[a_{\underline{x}^k}^{\underline{n}_k}, b_{\underline{x}^k}^{\underline{n}_k})$ . Thus,  $\underline{n}_{k+1}$  will be a least twice  $\underline{n}_k$ . We will discuss the relation between Sturmian  $\alpha$ -expansions and Ostrowski numeration (see [5], Chapter 5) in a later paper [15].

Now let us consider Sturmian  $\alpha$ -expansions for digit sequences  $\mathbf{c} \in \mathcal{D}^{\mathbb{N}}$  that are not necessarily proper. In particular, given any  $\mathbf{c} = .c_1c_2c_3 \dots \in \{0, 1\}^{\mathbb{N}}$  let

$$\varepsilon(\mathbf{c}) = f(c_1 + f(c_2 + f(c_3 + \dots))).$$

Let  $\prec$  denote lexicographic order on  $\{0, 1\}^{\mathbb{N}}$ . That is,  $\mathbf{c} \prec \mathbf{e}$ ,  $\mathbf{e} = .e_1e_2e_3 \dots$ , if and only if for some  $n \geq 1$ ,  $c_1 \dots c_n = e_1 \dots e_n$ ,  $c_{n+1} = 0$  and  $e_{n+1} = 1$ .

**Lemma 4.** If  $f$  is given by (11), then for any  $\mathbf{c} = .c_1c_2c_3 \dots \in \{0, 1\}^{\mathbb{N}}$ , the  $f$ -expansion  $\varepsilon(\mathbf{c}) = f(c_1 + f(c_2 + f(c_3 + \dots)))$  converges. Moreover, if  $\mathbf{c} \prec \mathbf{e}$  then  $\varepsilon(\mathbf{c}) \leq \varepsilon(\mathbf{e})$ .

*Proof.* Since  $f$  is nondecreasing,  $x_{n+1} \geq x_n$ , and moreover,  $x_n \leq 1$  since  $f(x) \leq 1$ . Thus  $\varepsilon(\mathbf{c})$  converges.

Suppose  $c_1 = 0$  and  $e_1 = 1$  so that  $\mathbf{c} \prec \mathbf{e}$ . Since  $f(c_2 + f(c_3 + \dots)) \leq 1$  and  $f(e_2 + f(e_3 + \dots)) \geq 0$ , it follows that  $c_1 + f(c_2 + \dots) \leq e_1 + f(e_2 + \dots)$ , and so  $f(c_1 + f(c_2 + \dots)) \leq f(e_1 + f(e_2 + \dots))$ .

Now suppose  $\mathbf{c} \prec \mathbf{e}$ . Let  $n >$  be such that  $c_1 c_2 \dots c_{n-1} = e_1 e_2 \dots e_{n-1}$ ,  $c_n = 0$  and  $e_n = 1$ . By the previous paragraph  $f(c_n + f(c_{n+1} + \dots)) \leq f(e_n + f(e_{n+1} + \dots))$ . Since  $f$  is increasing it follows that  $f(c_1 + \dots + f(c_n + f(c_{n+1} + \dots))) \leq f(e_1 + \dots + f(e_n + f(e_{n+1} + \dots)))$ .  $\square$

#### 4. ERGODIC PROPERTIES OF STURMIAN $\alpha$ -EXPANSIONS

As Rényi observed in his landmark paper [14], many properties of  $f$ -expansions reflect the “ergodic” properties of the corresponding  $f$ -transformation  $T$ . In this section, we compare the irrational rotation map  $Tx = x + \alpha \pmod{1}$  to the Gauss map  $Tx = 1/x \pmod{1}$  and the  $\beta$ -transformations  $Tx = \beta x \pmod{1}$ . A  $T$ -invariant probability measure  $\mu$  on  $[0, 1)$  is a Borel measure so that  $\mu([0, 1)) = 1$  and  $\mu(T^{-1}E) = \mu(E)$  for every Borel set  $E$ . A measure  $\mu$  is an *absolutely continuous* if there is a density  $\rho(x) \geq 0$  on  $[0, 1)$  with  $\mu(E) = \int_E \rho(x) dx$ , and “Lebesgue-equivalent” if  $\rho(x) > 0$  a.e. (i.e.,  $\rho(x) = d\mu/dx$  is the Radon-Nikodym derivative of  $\mu$ ). A measure is *ergodic* if  $\mu(T^k E) = 0$  implies  $\mu(A) = 0$  or 1.

All of the transformations we are discussing have an ergodic Lebesgue-equivalent invariant probability measure (“ELEM” for short). For the Gauss map, the density for this measure, called the *Gauss measure*, is given by  $\rho(x) = \frac{1}{\log 2} \frac{1}{1+x}$ . For  $\beta$ -transformations  $T$  with  $\beta \in \mathbb{N}$ , Lebesgue measure itself is invariant (i.e.,  $\rho(x) = 1$ ). When  $\beta \notin \mathbb{N}$ ,  $\rho(x)$  is a step function, and  $\mu$  is called the *Parry measure* (see [14] and [11]). Finally, if  $T$  is an irrational rotation transformation the ELEM is, again, Lebesgue measure.

For all three of these transformations, the existence of a ELEM implies that for the corresponding proper  $f$ -representation  $\mathbf{d}$ , almost every  $x \in [0, 1)$  is a *normal number*. In particular, for  $\mathbf{u} \in \mathcal{D}^n$  let

$$L_n(\mathbf{u}, \mathbf{d}) = |\{j \in [1, \dots, n] : \mathbf{d}|_{[j, \dots, j+|\mathbf{u}|-1]} = \mathbf{u}\}|$$

denote the number of *occurrences* of  $\mathbf{u}$  in the first  $n$  places in  $\mathbf{d}$ . A standard argument using Birkhoff ergodic theorem (see e.g. [19]) shows that for almost every  $x \in [0, 1)$ , for any  $\mathbf{u} \in \mathcal{D}^n$ ,

$$(13) \quad \lim_{n \rightarrow \infty} \frac{1}{n} L_n(\mathbf{u}, \mathbf{d}) = \int_{\Delta(d_1) \cap T^{-1}\Delta(d_2) \cap \dots \cap T^{-n+1}\Delta(d_n)} \rho(x) dx.$$

Thus, in a typical proper  $f$ -expansion, every finite sequence of digits occurs with a well defined frequency (which may sometimes be zero).

Beyond these simple facts, however, irrational rotation transformations  $T$  is very different from either the Gauss map or any  $\beta$ -transformation, and this leads to some unusual properties for Sturmian  $\alpha$ -expansions. To begin with, both the Gauss map and the  $\beta$ -transformations have other ergodic invariant measures besides their ELEMs. On the other hand, Lebesgue measure is the unique invariant measure for an irrational rotation  $T$ . This property is known as *unique ergodicity*. One corollary of unique ergodicity is that the ergodic theorem (13) converges for all  $x$  (see [19]) rather than just almost everywhere. Thus *every*  $x \in [0, 1)$  is a normal number for Sturmian  $\alpha$ -expansions.

Another consequence of unique ergodicity is  $O_T^+(x)$  is dense for every  $x$ . A minimal map  $T$  has no periodic or eventually periodic points. This means that there

are no periodic or eventually periodic proper Sturmian  $\alpha$ -expansions. In contrast, periodic points are dense for both the Gauss map and  $\beta$ -transformations. For both continued fractions and  $\beta$ -expansions, periodic expansions have important number theoretic consequences. However, even though there are no periodic expansions for Sturmian  $\alpha$ -expansions, minimality implies that all proper digit sequences for have the following “almost periodicity” property. Suppose a finite sequence  $\mathbf{u} \in \{0, 1\}^n$  occurs in the proper Sturmian  $\alpha$ -expansion  $\mathbf{c}$  of some  $y$ . Then there is a constant  $K > 0$  ( $K = K(\mathbf{u}, \alpha)$ ) so that  $\mathbf{u}$  occurs within  $K$  of an arbitrary location in the proper Sturmian  $\alpha$ -expansion  $\mathbf{d}$  of any  $x$ . Qualitatively, all proper Sturmian  $\alpha$ -expansions look pretty much alike.

A special case of eventual periodicity for  $\beta$ -expansions occurs when the proper expansion of  $x$  ends in zeros. In base  $\beta = 2$ , for example, such numbers are the dyadic rationals. We say the  $\beta$ -expansion of  $x$  is *finite*. Note that the  $\beta$ -expansion of  $x$  is a finite sum in this case. A similar situation can be imposed on continued fractions by defining  $T0 = \xi(0) = f(0) = 0$  and allowing  $0 \in \mathcal{D}$ . In this case, a number  $x \in [0, 1)$  has a finite continued fraction expansion if and only if it is rational.

At first it appears that there is no analogous situation for Sturmian expansions, since no proper Sturmian expansions end in infinitely many zeros. On the other hand, let  $x \in O_+(0)$  and let  $\mathbf{d} = .d_1d_2d_3\dots$  be the proper Sturmian expansion of  $x$ . Then there exists  $n_0 \in \mathbb{N}$  so that  $x = f(d_1 + f(d_2 + \dots + f(d_n)))$  for  $n \geq n_0$  (namely,  $T^{n_0}0 = x$ ). Moreover, it is easy to see that the digit sequence  $\mathbf{d}' = .d_1d_2\dots d_{n_0}0000\dots$  gives an improper Sturmian  $\alpha$ -expansion of  $x$ . It follows that there are uncountably many  $\mathbf{d}'' \in \{0, 1\}^{\mathbb{N}}$  satisfying  $\mathbf{d}' \prec \mathbf{d}'' \prec \mathbf{d}$  that are all Sturmian  $\alpha$ -expansions of  $x$ . So in this sense Sturmian  $\alpha$ -expansions can be highly non-unique.

## 5. ENTROPY AND GENERATORS

Let  $T$  be a *measure preserving transformation* of  $[0, 1)$ , with  $\mu$  the invariant Borel probability measure. Let  $\xi$  be a finite or countable partition of  $[0, 1)$  into positive measure Borel sets  $C$ . In general, we do not assume  $\mu$  is an ELEM or that  $\xi$  is a labeled interval partition. We say a Borel set  $A$  satisfies  $A \leq \xi$  if  $A$  is a union of elements  $C \in \xi$ . Let  $\xi \vee \xi' := \{C \cap C' : C \in \xi, C' \in \xi', \mu(C \cap C') > 0\}$ . Define  $\xi^{(n)} = \xi \vee T\xi \vee \dots \vee T^{-n+1}\xi$ , and if  $T$  is invertible, also define  $\xi^{(-n,n)} := T^{-n}\xi \vee \dots \vee \xi \vee \dots \vee T^n\xi$ . A partition  $\xi$  is called a *1-sided generator*  $T$  if for any Borel set  $A$ , and  $n \in \mathbb{N}$ , there exists  $A_n \in \xi^{(n)}$  so that  $\mu(A_n \Delta A) \rightarrow 0$ . If  $T$  is invertible,  $\xi$  is a *2-sided generator* if there is an  $A_n \leq \xi^{(-n,n)}$  such that  $\mu(A_n \Delta A) \rightarrow 0$ .

Let  $T$  be an  $f$ -transformation with an ELEM  $\mu$  and let  $\xi$  be the corresponding partition into fundamental intervals. It follows from the Lebesgue Density Theorem that  $\xi$  is a 1-sided generator if and only if  $\|\xi^{(n)}\| \rightarrow 0$ . This is equivalent to unique proper digits, and holds for all three transformations under consideration.

The entropy of a finite partition  $\xi$  is given by  $H(\xi) = -\sum_{C \in \xi} \mu(C) \log(C)$ . Note that  $H(\xi) \leq \log(|\xi|)$ . The entropy of  $T$  with respect to  $\xi$  is defined by  $h_\mu(T, \xi) = \lim_{n \rightarrow \infty} \frac{1}{n} H(\xi^{(n)})$ , and *entropy* of  $T$  is defined by  $h_\mu(T) = \sup_{H(\xi) < \infty} h_\mu(T, \xi)$ . In practice, the supremum in the definition of entropy often makes it difficult to apply directly, but the Kolmogorov-Sinai theorem, says the supremum is achieved,  $h_\mu(T) = h_\mu(T, \xi)$ , provided  $\xi$  is a (1- or 2-sided) generator.



In the case of an irrational rotation transformation  $T$ , we have  $|\xi^{(n)}| = c_{\mathbf{d}} = n+1$  (for any  $x$ ), so  $H(\xi^{(n)}) \leq \log(n+1)$ . Since  $\xi$  is (1-sided) a generator for  $T$  (by Theorem 2), the Kolmogorov-Sinai theorem implies  $h_{\mu}(T) = 0$ . The Kolmogorov-Sinai theorem also shows (using Kakeya's theorem) that for the  $\beta$ -transformation  $T$ , when  $\beta \in \mathbb{N}$ ,  $h_{\mu}(T) = \log \beta$ . This theorem does not apply directly when  $\beta \notin \mathbb{N}$ , or to the Gauss map. On the other hand, for a lot of  $f$ -transformations  $T$ , entropy is given by *Rohlin's entropy formula*:

$$(14) \quad h_{\mu}(T) = \int_0^1 \log |T'(x)| d\mu.$$

In particular (14) gives the well known result  $h_{\mu}(T) = \pi^2/(6 \log 2)$  for the the Gauss map (with Gauss measure) and gives  $h_{\mu}(T) = \log \beta$  for all  $\beta$ -transformations  $T$ . Note that the entropy is positive in both of these cases.

The validity of Rohlin's formula (14) can be deduced under various hypotheses (see e.g., [16],[13]), which always seem, at least implicitly, to include Kakeya's hypothesis (9). This suggests that (14) is valid only in the case  $h_{\mu}(T) > 0$ . We note, however, that for Rohlin's entropy formula gives the correct answer  $h_{\mu}(T) = 0$  for irrational rotation transformations  $T$ , if only by coincidence, since they satisfy  $T'(x) \equiv 1$ .

The fact that irrational rotation transformations  $T$  have zero entropy contributes to the strangeness of Sturmian  $\alpha$ -expansions. As our calculation above, using the Kolmogorov-Sinai theorem shows, zero entropy is due to the low complexity  $c_{\mathbf{d}}(n) = n+1$  of Sturmian sequences. Thus entropy zero is closely related to the slow convergence Sturmian  $\alpha$ -expansions. Heuristically, each additional digit in a Sturmian  $\alpha$ -expansion contributes very little new information about the number  $x$ . However, even more significant is fact that irrational rotation transformations  $T$  are invertible, whereas  $\beta$ -transformations and the Gauss map are not.

It follows from the invertibility of an irrational rotation transformation  $T$  that the Sturmian  $\alpha$ -expansion of any  $x$  extends to a two-sided sequence

$$\mathbf{d} = \dots d_{-2}d_{-1}d_0.d_1d_2\dots,$$

where  $d_n = \xi(T^{n-1}x)$ . Since the digits to the right of the "radix point" completely determine  $x$ , it follows that the digits to the left contribute no new information. Equivalently, a typical one-sided Sturmian sequence has a unique two-sided extension. The only exceptions to this (which are countable in number) occur when  $x = n\alpha \bmod 1$  for  $n > 1$ . In such a case there are exactly two left-extensions, which differ on exactly two adjacent digits.

For  $\beta$ -expansions, allowing finitely many (nonzero) digits to the left of the radix point gives an expansion of any  $x \in \mathbb{R}$ . In particular, the  $\beta$ -expansion of the digit sequence  $\mathbf{d} = d_{-N}d_{-N+1}\dots d_0.d_1d_2\dots$  is  $x = \sum_{k=-N}^{\infty} d_k\beta^{-k}$ . For continued fractions, expansions of all  $x \in \mathbb{R}$  are obtained with a single non-zero digit to the left of the radix-point. The digits for the expansion

$$x = d_0 + \frac{1}{d_1 + \frac{1}{d_2 + \frac{1}{d_3 + \dots}}}.$$

are usually written  $\mathbf{d} = [d_0; d_1, d_2, d_3, \dots]$ . In both cases, this works because the corresponding  $f$ -transformation  $T$  is not invertible.

To interpret continued fraction and  $\beta$ -expansions with more non-zero digits to the left of the radix point, however, one needs to consider the natural extension of  $T$ . This is the smallest invertible measure preserving transformation  $\tilde{T}$  having  $T$  as a factor. For example, if  $Tx = 2x \bmod 1$  on  $[0, 1)$  (the  $f$ -transformation for ordinary base 2 expansions), the natural extension is the Lebesgue measure preserving map  $\tilde{T} : [0, 1)^2 \rightarrow [0, 1)^2$ , defined  $\tilde{T}(x, y) = (2x \bmod 1, (\lfloor 2x \rfloor + y)/2)$ . This Lebesgue measure preserving mapping, called the *baker's transformation*, is isomorphic to the 2-sided Bernoulli shift with entropy  $\log 2$ . It is known that the natural extensions for any  $\beta$ -transformations is isomorphic to a Bernoulli shift, as is the natural extension of the Gauss map (see [3]).

Since an irrational rotation transformation  $T$  is already invertible, it is its own natural extension. There is no new information to be obtained by an extension to a bijection. Entropy theory provides another way to understand this phenomenon. A well-known theorem says that any invertible map  $T$  with a 1-sided generator (like the irrational rotation transformation) must have entropy zero (see [19]). Thus, no finite partition can be a 1-sided generator for any invertible transformation  $\tilde{T}$  with positive entropy. It is easy to see that the partition  $\tilde{\xi} = \{[0, 1/2) \times [0, 1), [1/2, 1) \times [0, 1)\}$  is a 2-sided generator for the baker's transformation  $\tilde{T}$ , since  $\tilde{\xi}^{(-n, n)}$  is the partition of  $[0, 1)^2$  into  $2^{-n} \times 2^{-n}$  squares. But  $\tilde{\xi}^{(n)}$  is the partition of  $[0, 1)^2$  into  $2^{-n} \times 1$  squares, and the factor corresponding to this partition is just  $Tx = 2x \bmod 1$ . Thus  $\tilde{\xi}$  is not a 1-sided generator for  $\tilde{T}$ .

## 6. GENERALIZATIONS

Let  $\xi$  and  $\xi'$  be partitions of  $[0, 1)$  into finitely or countably many intervals of the form  $\Delta = [a, b)$ . Assume, moreover, that there is a nondecreasing function  $\xi : [0, 1) \rightarrow \mathbb{Z}$  that is constant on each  $\Delta \in \xi$ , and is unequal on different  $\Delta, \Delta' \in \xi$ . The existence of such a function is automatic if  $|\xi| = d < \infty$ , in which case we usually take  $\mathcal{D} := \xi([0, 1)) = \{0, 1, \dots, d-1\}$ . It is a more substantial restriction if  $|\xi| = \infty$ . In particular, the only limit points of the set of endpoints of  $\Delta \in \xi$  can be 0 and 1 (and at least one must be a limit point).

Let  $\mu$  denote Lebesgue measure, and suppose  $\tau : \xi \rightarrow \xi'$  is such that  $\mu(\tau(\Delta)) = \mu(\Delta)$  for every  $\Delta \in \xi$ . Let  $T : [0, 1) \rightarrow [0, 1)$  be the mapping so that  $T$  maps each  $\Delta \in \xi$  by translation to  $\tau(\Delta)$ . We call  $T$  an *interval exchange transformation* (IET) if  $|\xi| < \infty$ , or an *infinite interval exchange transformation* (IIET) if  $|\xi| = \infty$ . In either case,  $T$  preserves Lebesgue measure.

Let  $F(x) = T(x) + \xi(x)$  and note that  $F : [0, 1) \rightarrow \mathbb{R}$  is increasing, continuous on each  $\Delta \in \xi$  and continuous from the right on  $[0, 1)$ . We define  $f(x) = F^{-1}(x)$ , extended to continuous non-decreasing  $f : \mathbb{R} \rightarrow [0, 1)$  with  $f(\mathbb{R}) = [0, 1)$ .

Let  $T$  be an IET or IIET. We call  $a \in [0, 1)$  a *cut-point* of  $\xi$  if it is the left endpoint of some  $\Delta \in \xi$ . Given a cutpoint  $a$ , we define  $\xi_a$  to be all the intervals in  $\Delta \in \xi$  so that  $x < a$  for  $x \in \Delta$ . In particular,  $\xi_a$  is a partition of  $[0, a)$  into intervals. We say  $T$  is *reducible* if there is an  $a \in [0, 1)$  that is a cut-point for both  $\xi$  and  $\xi'$ , and such that  $\tau(\xi_a) = \xi'_a$ . If there are no such  $a$ , we say  $T$  is *irreducible*. If  $T$  is reducible,  $T([0, a)) = [0, a)$ , and it cannot be minimal or ergodic.

If  $T$  is an irreducible IET, Keane [8] showed that  $T$  is minimal if and only if the left endpoints of all the intervals  $\Delta \in \xi$  have *infinite* and *distinct orbits* (this is abbreviated IDOC). He proved that if the lengths  $\ell_0, \ell_1, \dots, \ell_{d-1}$  of the intervals in  $\xi$

are rationally independent then IDOC follows. The case  $|\xi| = 2$  is just an irrational rotation transformation  $T$ . If  $T$  is an IET, we call the  $f$ -expansions IET-expansions.

**Proposition 5.** *If  $T$  is an irreducible IET with rationally independent interval lengths (or that satisfies IDOC) then the corresponding IET-expansions are valid.*

The proof of Proposition 5 is almost exactly the same as the proof of Theorem 2. It depends on the fact that IDOC implies  $\|\xi^{(n)}\| \rightarrow 0$ .

Unique ergodicity for an irreducible IET  $T$  is a bit stronger (and more difficult to prove) than minimality, but it holds for almost every choice of lengths  $\ell_0, \ell_1, \dots, \ell_{d-1}$  of intervals in  $\xi$  (see [18], [9]), as does weak (but never strong) mixing, (see [1]). The entropy of an IET  $T$  is always zero. In summary, IET-expansions have many of the same properties as Sturmian  $\alpha$ -expansions, with at least one notable difference. An interval exchange transformation  $T$  can be minimal but not uniquely ergodic. In such a case there will be non-normal numbers  $x$  for the expansions, as well as up to  $d$  different kinds of normal numbers (corresponding to, possibly,  $d$  different ergodic invariant measures).

We conclude by considering expansions based on the well-known *von Neumann adding machine* (or *odometer*) transformation  $T$ . Let  $a_n = 1 - 1/2^n$ ,  $b_n = 1/2^n$ ,  $\xi = \{[a_n, a_{n+1}) : n = 0, 1, 2, \dots\}$ ,  $\xi' = \{[b_{n+1}, b_n) : n = 0, 1, 2, \dots\}$ , and  $\tau([a_n, a_{n+1})) = [b_{n+1}, b_n)$ . Let  $T$  be the corresponding IIET, and define the labels  $\xi([a_n, a_{n+1})) = n$ , noting that  $|\xi| = \infty$  and  $\mathcal{D} = \xi([0, 1)) = \mathbb{N} \cup \{0\}$ . Define  $f : \mathbb{R} \rightarrow [0, 1)$  as the extension of  $F^{-1}$ , where  $F(x) = T(x) + \xi(x)$ , so that  $\overline{f(\mathbb{R})} = [0, 1]$ . We call the corresponding  $f$ -expansions of  $x \in [0, 1)$  *von Neumann expansions*. The fact that von Neumann expansions are valid follows from the unique ergodicity of  $T$ , which is well known. In particular, the endpoints of the  $\Delta \in \xi$  have dense orbits, and this can be used to show that  $\|\xi^{(n)}\| \rightarrow 0$ . The entropy of  $T$  is zero.

To find the von Neumann expansion of  $x \in [0, 1)$ , we first identify  $x$  with its ordinary binary expansion, i.e.,  $x = .x_1x_2x_3\dots$  means  $x = \sum_{k=1}^{\infty} x_k\beta^{-k}$ . It is easy to see that

$$T(.x_1x_2x_3\dots) = \begin{cases} .1x_2x_3x_4\dots & \text{if } x_1 = 0, \\ .00\dots 01x_{n+1}x_{n+2} & \text{if } x_1x_2\dots x_{n-1} = 11\dots 1 \text{ and } x_n = 0. \end{cases}$$

So  $T$  adds .1 to  $.x_1x_2x_3\dots$  with right carry, which is why  $T$  is called an “adding machine”. Moreover,  $\xi(.1^n0x_{n+2}x_{n+3}\dots) = n$ , where  $n \geq 0$ .

As an example, if  $x = 1/3 = .01010101\dots$ , then

$x = .010101010101010101\dots$	0
$Tx = .110101010101010101\dots$	2
$T^2x = .001101010101010101\dots$	0
$T^3x = .101101010101010101\dots$	1
$T^4x = .011101010101010101\dots$	0
$T^5x = .111101010101010101\dots$	4
$T^6x = .000011010101010101\dots$	0
$T^7x = .100011010101010101\dots$	1
$\dots$	

where the numbers in the right column are  $d_n$  for  $n = 1, 2, 3, \dots$ . Thus we have the digit sequence  $\mathbf{d} = [0, 2, 0, 1, 0, 4, 0, 1, 0, 2, 0, 1, 0, 3, 0, 1, 0, 2, 0, 1, 0, 6, \dots]$ .

Notice the list  $x, Tx, T^2x\dots$ , the first column alternates 0 and 1, the second 00 and 11, third 0000 and 1111 (the first 0000 being truncated to 00), etc. Moreover,

0s in earlier columns mask 1s in later columns. This implies that every 2nd digit of any von Neumann expansion  $\mathbf{d} = [d_1, d_2, d_3, \dots]$  is a 0, every 4th digit a 1, every 8th digit a 2,  $\dots$ , every  $2^{n+1}$ st digit an  $n$ . About  $2^{n+1}$  digits of  $\mathbf{d}$  determines one binary digit of  $x$ . So like Sturmian  $\alpha$ -expansions, von Neumann expansions converge slowly.

As a final remark, we note that if we define  $\mathbf{e} = .e_1e_2e_3\dots$ , by  $e_n = d_n \bmod 2$ , then the resulting sequence is a *Toeplitz sequence* (see [6]). For example, the Toeplitz sequence corresponding to  $1/3$  is  $\mathbf{c} = .00010001000101010001000\dots$ . Since it is possible to recover the von Neumann sequence from the Toeplitz sequence, the map  $x \mapsto \mathbf{c}$  is injective. However, we don't know if it is possible to recover  $x$  from  $\mathbf{c}$  by a simple formula like an  $f$ -expansion.

#### REFERENCES

- [1] Artur Avila and Giovanni Forni. Weak mixing for interval exchange transformations and translation flows. *Ann. of Math. (2)*, 165(2):637–664, 2007.
- [2] B. H. Bissinger. A generalization of continued fractions. *Bull. Amer. Math. Soc.*, 50:868–876, 1944.
- [3] Karma Dajani and Cor Kraaikamp. *Ergodic theory of numbers*, volume 29 of *Carus Mathematical Monographs*. Mathematical Association of America, Washington, DC, 2002.
- [4] C. J. Everett. Representations for real numbers. *Bull. Amer. Math. Soc.*, 52:861–869, 1946.
- [5] N. Pytheas Fogg. *Substitutions in dynamics, arithmetics and combinatorics*, volume 1794 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2002. Edited by V. Berthé, S. Ferenczi, C. Mauduit and A. Siegel.
- [6] Konrad Jacobs and Michael Keane. 0 – 1-sequences of Toeplitz type. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 13:123–131, 1969.
- [7] S. Kakeya. On the generalized scale of notation. *Japan J. Math.*, 1:95–108, 1924.
- [8] Michael Keane. Interval exchange transformations. *Math. Z.*, 141:25–31, 1975.
- [9] Howard Masur. Interval exchange transformations and measured foliations. *Ann. of Math. (2)*, 115(1):169–200, 1982.
- [10] Marston Morse and Gustav A. Hedlund. Symbolic dynamics II. Sturmian trajectories. *Amer. J. Math.*, 62:1–42, 1940.
- [11] W. Parry. On the  $\beta$ -expansions of real numbers. *Acta Math. Acad. Sci. Hungar.*, 11:401–416, 1960.
- [12] W. Parry. Representations for real numbers. *Acta Math. Acad. Sci. Hungar.*, 15:95–105, 1964.
- [13] Mark Pollicott and Michiko Yuri. *Dynamical systems and ergodic theory*, volume 40 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1998.
- [14] A. Rényi. Representations for real numbers and their ergodic properties. *Acta Math. Acad. Sci. Hungar.*, 8:477–493, 1957.
- [15] E. A. Robinson. Sturmian and ostrowski numeration. preprint, 2010.
- [16] V. A. Rohlin. Exact endomorphisms of a Lebesgue space. *Izv. Akad. Nauk SSSR Ser. Mat.*, 25:499–530, 1961.
- [17] Fritz Schweiger. *Ergodic theory of fibred systems and metric number theory*. Oxford Science Publications. The Clarendon Press Oxford University Press, New York, 1995.
- [18] William A. Veech. Gauss measures for transformations on the space of interval exchange maps. *Ann. of Math. (2)*, 115(1):201–242, 1982.
- [19] Peter Walters. *An introduction to ergodic theory*, volume 79 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.

DEPARTMENT OF MATHEMATICS, GEORGE WASHINGTON UNIVERSITY, 2115 G ST. NW, WASHINGTON, DC 20052

*E-mail address:* robinson@gwu.edu