

THE WIGNER-FOKKER-PLANCK EQUATION: STATIONARY STATES AND LARGE TIME BEHAVIOR

ANTON ARNOLD, IRENE M. GAMBA, MARIA PIA GUALDANI, STÉPHANE MISCHLER,
CLÉMENT MOUHOT, AND CHRISTOF SPARBER

ABSTRACT. We consider the linear Wigner-Fokker-Planck equation subject to confining potentials which are smooth perturbations of the harmonic oscillator potential. For a certain class of perturbations we prove that the equation admits a unique stationary solution in a weighted Sobolev space. A key ingredient of the proof is a new result on the existence of spectral gaps for Fokker-Planck type operators in certain weighted L^2 -spaces. In addition we show that the steady state corresponds to a positive density matrix operator with unit trace and that the solutions of the time-dependent problem converge towards the steady state with an exponential rate.

1. INTRODUCTION

This work is devoted to the study of the *Wigner-Fokker-Planck equation* (WFP), considered in the following dimensionless form (where all physical constants are normalized to one for simplicity):

$$(1.1) \quad \begin{cases} \partial_t w + \xi \cdot \nabla_x w + \Theta[V]w = \Delta_\xi w + 2 \operatorname{div}_\xi (\xi w) + \Delta_x w, \\ w|_{t=0} = w_0(x, \xi), \end{cases}$$

where $x, \xi \in \mathbb{R}^d$, for $d \geq 1$, and $t \in \mathbb{R}_+$. Here, $w(t, x, \xi)$ is the (real valued) *Wigner transform* [38] of a quantum mechanical *density matrix* $\rho(t, x, y)$, as defined by

$$(1.2) \quad w(t, x, \xi) := \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \rho \left(t, x + \frac{\eta}{2}, x - \frac{\eta}{2} \right) e^{-i\xi \cdot \eta} d\eta.$$

Recall that, for any time $t \in \mathbb{R}_+$, a quantum mechanical (mixed) state is given by a positive, self-adjoint *trace class operator* $\rho(t) \in \mathcal{T}_1^+$. Here we denote by $\mathcal{B}(L^2(\mathbb{R}^d))$ the set of bounded operators on $L^2(\mathbb{R}^d)$ and by

$$\mathcal{T}_1 := \{ \rho \in \mathcal{B}(L^2(\mathbb{R}^d)) : \operatorname{tr} |\rho| < \infty \},$$

the corresponding set of trace-class operators. We consequently write $\rho \in \mathcal{T}_1^+ \subset \mathcal{T}_1$, if in addition $\rho \geq 0$ (in the sense of non-negative operators). Since $\mathcal{T}_1 \subset \mathcal{T}_2$, the space of *Hilbert-Schmidt operators*, i.e.

$$\mathcal{T}_2 := \{ \rho \in \mathcal{B}(L^2(\mathbb{R}^d)) : \operatorname{tr}(\rho^* \rho) < \infty \},$$

2000 *Mathematics Subject Classification.* 82C10, 35S10, 74H40, 81Q15.

Key words and phrases. Wigner transform, Fokker Planck operator, spectral gap, stationary solution, large time behavior.

A. Arnold acknowledges partial support from the FWF (project “Quantum Transport Equations: Kinetic, Relativistic, and Diffusive Phenomena” and Wissenschaftskolleg “Differentialgleichungen”), the ÖAD (Amadeus project), and the Newton Institute of Cambridge University. I. M. Gamba is supported by NSF-DMS0507038. M. P. Gualdani is supported by NSF-DMS 0807636. C. Mouhot would like to thank Cambridge University who provided repeated hospitality in 2009 thanks to the Award No. KUK-II-007-43, funded by the King Abdullah University of Science and Technology (KAUST). C. Sparber has been supported by the Royal Society through his Royal Society University Research Fellowship.

we can identify the operator $\rho(t)$ with its corresponding integral kernel $\rho(t, \cdot, \cdot) \in L^2(\mathbb{R}^{2d})$, the so-called *density matrix*. Consequently, $\rho(t)$ acts on any given function $\varphi \in L^2(\mathbb{R}^d)$ via

$$(\rho(t)\varphi)(x) = \int_{\mathbb{R}^d} \rho(t, x, y) \varphi(y) dy.$$

Using the Wigner transformation (1.2), which by definition yields a *real-valued* function $w(t, \cdot, \cdot) \in L^2(\mathbb{R}^{2d})$, one obtains a *phase-space description* of quantum mechanics, reminiscent of classical statistical mechanics, with $x \in \mathbb{R}^d$ being the position and $\xi \in \mathbb{R}^d$ the momentum. However, in contrast to classical phase space distributions, $w(t, x, \xi)$ in general also takes *negative values*.

Equation (1.1) governs the time evolution of $w(t, x, \xi)$ in the framework of so-called *open quantum systems*, which model both the Hamiltonian evolution of a quantum system and its interaction with an environment (see [14], e.g.). Here, we specifically describe these interactions by the Fokker-Planck (FP) type diffusion operator on the r.h.s. of (1.1). For notational simplicity we use here only normalized constants in the quantum FP operator. However, all of the subsequent analysis also applies to the general WFP model presented in [35] (cf. Remark 2.4 below). Potential forces acting on $w(t, \cdot, \cdot)$ are taken into account by the pseudo-differential operator

$$(1.3) \quad (\Theta[V]f)(x, \xi) := -\frac{i}{(2\pi)^d} \iint_{\mathbb{R}^{2d}} \delta V(x, \eta) f(x, \xi') e^{i\eta \cdot (\xi - \xi')} d\xi' d\eta,$$

where the symbol δV is given by

$$(1.4) \quad \delta V(x, \eta) = V\left(x + \frac{\eta}{2}\right) - V\left(x - \frac{\eta}{2}\right),$$

and V is a given real valued function. The WFP equation is a kinetic model for quantum mechanical charge-transport, including diffusive effects, as needed, e.g., in the description of quantum Brownian motion [16], quantum optics [19], and semiconductor device simulations [17]. It can be considered as a quantum mechanical generalization of the usual kinetic Fokker-Planck equation (or Kramer's equation), to which it is known to converge in the classical limit $\hbar \rightarrow 0$, after an appropriate rescaling of the appearing physical parameters [10]. The WFP equation has been partly derived in [11] as a rigorous scaling limit for a system of particles interacting with a heat bath of phonons. Additional "derivations" (based on formal arguments from physics) can also be found in [15, 16, 36, 37].

In recent years, mathematical studies of WFP type equations mainly focused on the *Cauchy problem* (with or without self-consistent Poisson-coupling), see [2, 3, 4, 7, 9, 12]. In these works, the task of establishing a rigorous definition of the particle density $n(t, x)$ has led to various functional analytical settings. To this end, it is important to note that the dynamics induced by (1.1) maps $\mathcal{S}_1^+(L^2(\mathbb{R}^d))$ into itself, since the so-called *Lindblad condition* is fulfilled (see again Remark 2.4 below). For more details on this we refer to [7, 9] and the references given therein. In the present work we shall be mainly interested in the asymptotic behavior as $t \rightarrow +\infty$ of solutions to (1.1). To this end, we first need to study the stationary problem corresponding to (1.1). Let us remark, that stationary equations for open quantum systems, based on the Wigner formalism, seem to be rather difficult to treat as only very few results exist (in spite of significant efforts, cf. [6] where the stationary, inflow-problem for the linear Wigner equation in $d = 1$ was analyzed). In fact the only result for the WFP equation is given in [35], where the existence of a unique steady state for a quadratic potential $V(x) \propto |x|^2$ has been proved. However, the cited work is based on several explicit calculations, which can *not* be applied in the case of a more general potential $V(x)$.

The goal of the present paper is *twofold*: First, we aim to establish the existence of a normalized steady state $w_\infty(x, \xi)$ for (1.1) in the case of confining potentials $V(x)$, which are given by a suitable class of perturbations of quadratic potentials (thus, $V(x)$ can be considered as a perturbed harmonic oscillator potential). The second goal is to study the long-time behavior of (1.1). We shall prove exponential convergence of the time-dependent solution $w(t, x, \xi)$ towards w_∞ as $t \rightarrow +\infty$. In a subsequent step, we shall also prove that the stationary Wigner function w_∞ corresponds to a density matrix operator $\rho_\infty \in \mathcal{T}_1^+$. Remarkably, this proof exploits the positivity preservation of the *time-dependent* problem (using results from [9]), via a stability property of the steady states.

To establish the existence of a (unique) steady state w_∞ , the basic idea is to prove the existence of a spectral gap for the unperturbed Wigner-Fokker-Planck operator with quadratic potential. This implies invertibility of the (unperturbed) WFP-operator on the orthogonal of its kernel. Assuming that the perturbation potential is sufficiently small with respect to this spectral gap, we can set up a fixed point iteration to obtain the existence of w_∞ . The key difficulty in doing so, is the choice of a suitable functional setting: On the one hand a Gaussian weighted L^2 -space seems to be a natural candidate, since it ensures dissipativity of the unperturbed WFP-operator (see Section 3). Indeed, this space is classical in the study of the long-time behavior of the classical (kinetic) Fokker-Planck equation, see [27]. However, it *does not* allow for feasible perturbations through $\Theta[V_0]$. In fact, even for smooth and compactly supported perturbation potentials V_0 , the operator $\Theta[V_0]$ would be *unbounded* in such an L^2 -space (due to the non-locality of $\Theta[V_0]$, see Remark 5.2). We therefore have to enlarge the functional space and to show that the unperturbed WFP-operator then still has a (now smaller) spectral gap. This is a key step in our approach. It is a result from spectral and semigroup theory (cf. Proposition 4.8) which is related to a more general mathematical theory of spectral gap estimates for kinetic equations, developed in parallel in [25] (see also [30]). We also remark that for $V(x) = |x|^2$ the WFP equation corresponds to a differential operator with quadratic symbol [35] and thus our approach is closely related to recent results for hypo-elliptic and sub-elliptic operators given in [18, 27, 32].

Comparing our methods to closely related results in the quantum mechanical literature, we first cite [21], where several criteria for the existence of stationary density matrices for quantum dynamical semigroups (in Lindblad form) were obtained by means of compactness methods. In [5] the applicability of this general approach to the WFP equation was established. In [23, 22] sufficient conditions (based on commutator relations for the Lindblad operators) for the large-time convergence of open quantum systems were derived. However, these techniques do not provide a rate of convergence towards the steady states. In comparison to that, the novelty of the present work consists in establishing steady states in a kinetic framework and in proving exponential convergence rates. However, the optimality of such rates for the WFP equation remains an open problem. In this context one should also mention the recent work [26], in which explicit estimates on the norm a semigroup in terms of bounds on the resolvent of its generator are obtained, very much along the same lines as in present paper and in [25].

The paper is organized as follows: In Section 2 we present the basic mathematical setting (in particular the class of potentials covered in our approach) and state our two main theorems. In Section 3 we collect some known results for the case of a purely quadratic potential and we introduce the Gaussian weighted L^2 -space for this unperturbed WFP operator. This basic setting is then generalized in Section 4, which contains the core of our (enlarged) functional framework: We shall prove

new spectral gap estimates for the WFP operator with a harmonic potential in L^2 -spaces with only *polynomial weights*. In Section 5 we prove the boundedness of the operator $\Theta[V_0]$ in these spaces. Finally, Section 6 concludes the proof of our main result by combining the previously established elements. Appendix A includes the rather technical proof of a preliminary step which guarantees the applicability of the spectral method developed in [25].

2. SETTING OF THE PROBLEM AND MAIN RESULTS

2.1. Basic definitions. In this work we shall use the following convention for the Fourier transform of a function $\varphi(x)$:

$$\widehat{\varphi}(k) := \int_{\mathbb{R}^d} \varphi(x) e^{-ik \cdot x} dx.$$

From now on we shall assume that the (real valued, time-independent) potential V , appearing in (1.1), is of the form

$$(2.1) \quad V(x) = \frac{1}{2} |x|^2 + \lambda V_0(x),$$

with $\lambda \in \mathbb{R}$ some given *perturbation parameter*. In other words we consider a perturbation V_0 of the harmonic oscillator potential. An easy calculation shows that for such a V the stationary equation, corresponding to (1.1), can be written as

$$(2.2) \quad Lw = \lambda \Theta[V_0]w,$$

where L is the linear operator

$$(2.3) \quad Lw := -\xi \cdot \nabla_x w + x \cdot \nabla_\xi w + \Delta_\xi w + 2 \operatorname{div}_\xi(\xi w) + \Delta_x w.$$

Remark 2.1. When considering the slightly more general class of potentials

$$V(x) = \frac{1}{2} |x|^2 + \alpha \cdot x + \lambda V_0(x), \quad \lambda_0 \in \mathbb{R},$$

we would find, instead of (2.3), the following operator: $L_\alpha w := Lw + \alpha \cdot \nabla_\xi w$. Thus, by the change of variables $x \mapsto x + \alpha$ we are back to (2.3).

The basic idea for establishing the existence of (stationary) solutions to (2.2) is the use of a fixed point iteration. However, L has a non-trivial kernel. Indeed it has been proved in [35] that, in the case $\lambda = 0$, there exists a unique stationary solution $\mu \in \mathcal{S}(\mathbb{R}^{2d})$, satisfying

$$(2.4) \quad L\mu = 0$$

and the normalization condition

$$(2.5) \quad \iint_{\mathbb{R}^{2d}} \mu(x, \xi) dx d\xi = 1.$$

Explicitly, μ can be written as

$$(2.6) \quad \mu = c e^{-A(x, \xi)},$$

where the function A is given by

$$(2.7) \quad A(x, \xi) := \frac{1}{4} (|x|^2 + 2x \cdot \xi + 3|\xi|^2),$$

and the constant $c > 0$ is chosen such that (2.5) holds. Note that for any $\rho \in \mathcal{T}_1$ such that $w \in L^1(\mathbb{R}^{2d})$ the following formal identity

$$\operatorname{tr} \rho = \int_{\mathbb{R}^d} \rho(x, x) dx = \iint_{\mathbb{R}^{2d}} w(x, \xi) dx d\xi,$$

can be rigorously justified by a limiting procedure in \mathcal{T}_1 , see [1]. Since $\text{tr} \rho$ is proportional to the total mass of the quantum system, we can interpret condition (2.5) as a mass normalization.

In the following, we shall denote by $\sigma > 0$ the biggest constant such that

$$(2.8) \quad \text{Hess}A - \sigma \mathbf{I} \geq 0, \quad \text{for all } (x, \xi) \in \mathbb{R}^{2d},$$

in the sense of positive definite matrices, where \mathbf{I} denotes the identity matrix on \mathbb{R}^{2d} . In the analysis of the classical FP equation, condition (2.8) is referred to as the *Bakry-Emery criterion* [8]. In our case one easily computes

$$\sigma = 1 - 1/\sqrt{2}.$$

In a Gaussian weighted L^2 -space, σ will be the spectral gap of the unperturbed WFP-operator and hence the decay rate towards the corresponding steady μ (cf. (3.9), (3.10) below).

The functional setting of our problem will be based on the following weighted Hilbert spaces. While the stationary and transient Wigner function are real valued, we need to consider function spaces over \mathbb{C} , for the upcoming spectral analysis.

Definition 2.2. For any $m \in \mathbb{N}$, we define $\mathcal{H}_m := L^2(\mathbb{R}^{2d}, \nu_m^{-1} dx d\xi)$, where the weight is

$$\nu_m^{-1} := 1 + A^m(x, \xi).$$

We equip \mathcal{H}_m with the inner product

$$\langle f, g \rangle_{\mathcal{H}_m} = \iint_{\mathbb{R}^{2d}} \frac{f \bar{g}}{\nu_m} dx d\xi.$$

Clearly, we have that $\mathcal{H}_{m+1} \subset \mathcal{H}_m$, for all $m \in \mathbb{N}$.

2.2. Main results. With these definitions at hand, we can now state the main theorems of our work. Note that for the sake of transparency we did not try to optimize the appearing constants.

Theorem 1. *Let $m \geq Kd$ be some fixed integer, where $K = K(A) \in (1, 144]$ is a constant depending only on $A(x, \xi)$ (defined in Lemma 4.2). Assume that the perturbation potential V_0 satisfies*

$$(2.9) \quad \Gamma_m := C_m \max_{|j| \leq m} \|\partial_x^j V_0\|_{L^\infty(\mathbb{R}^d)} < +\infty,$$

where $C_m > 0$ depends only on m and d , as seen in the proof of Proposition 5.1. Next we fix some $\tilde{\gamma}_m \in (0, \gamma_m)$, where $\gamma_m > 0$ is given in (4.7). Furthermore, let the perturbation parameter λ satisfy

$$(2.10) \quad |\lambda| < \frac{\tilde{\gamma}_m}{\Gamma_m \delta_m},$$

where $\delta_m = \delta_m(\tilde{\gamma}_m) > 1$ is defined in (4.15). Then it holds:

- (i) *The stationary Wigner-Fokker-Planck equation (2.2) admits a unique weak solution $w_\infty \in \mathcal{H}_m \cap H^1(\mathbb{R}^{2d})$, satisfying $\iint_{\mathbb{R}^{2d}} w_\infty dx d\xi = 1$. Moreover, w_∞ is real valued and satisfies $w_\infty \in H_{\text{loc}}^2(\mathbb{R}^{2d})$.*
- (ii) *Denoting by $w \in C([0, \infty), \mathcal{H}_m)$ the unique mild solution to equation (1.1) with some initial data $w_0 \in \mathcal{H}_m$, such that $\iint_{\mathbb{R}^{2d}} w_0 dx d\xi = 1$, we have*

$$\|w(t) - w_\infty\|_{\mathcal{H}_m} \leq \delta_m e^{-\kappa_m t} \|w_0 - w_\infty\|_{\mathcal{H}_m}, \quad \forall t \geq 0,$$

with an exponential decay rate

$$\kappa_m := \tilde{\gamma}_m - |\lambda| \delta_m \Gamma_m > 0.$$

(iii) Concerning the continuity of $w_\infty = w_\infty(\lambda)$ w.r.t. λ , we have

$$\|w_\infty - \mu\|_{\mathcal{H}_m} \leq \frac{|\lambda|\delta_m\Gamma_m}{\tilde{\gamma}_m - |\lambda|\delta_m\Gamma_m} \|\mu\|_{\mathcal{H}_m}.$$

Remark 2.3. In this result, the constant $\sigma_m := \tilde{\gamma}_m/\delta_m > 0$, roughly speaking, plays the same role for L on \mathcal{H}_m as $\sigma > 0$ does in the case of \mathcal{H} , where it is nothing but the size of the spectral gap, see Proposition 3.5. For L on \mathcal{H} , σ also gives the exponential decay rate in the unperturbed case $\lambda = 0$. For L on \mathcal{H}_m the situation is more complicated. Here, assertion (ii) yields an exponential decay of the unperturbed semi-group with rate $\kappa_m = \tilde{\gamma}_m \in (0, \gamma_m)$ and $\gamma_m \neq \sigma_m$ (but possibly equal to σ , as can be seen from (4.15)). In addition, one should note that $\delta_m > 1$ may blow-up as $\tilde{\gamma}_m \nearrow \gamma_m$, cf. estimate (4.11).

Theorem 1 is formulated in the Wigner picture of quantum mechanics. We shall now turn our attention to the corresponding density matrix operators $\rho(t)$. This is important since it is *a priori* not clear that w_∞ is physically meaningful – in the sense of being the Wigner transform of a positive trace class operator. To this end we denote by ρ_∞ the Hilbert-Schmidt operator corresponding to the kernel $\rho_\infty(x, y)$, which is obtained from $w_\infty(x, \xi)$ by the *inverse Wigner transform*, i.e.

$$\rho_\infty(x, y) = \int_{\mathbb{R}^d} w_\infty\left(\frac{x+y}{2}, \xi\right) e^{-i\xi \cdot (x-y)} d\xi.$$

Analogously we denote by ρ_0 the Hilbert-Schmidt operator corresponding to the initial Wigner function $w_0 \in \mathcal{H}_m$.

We remark that the existence of a unique mild solution of equation (1.1) on \mathcal{H}_m will be a byproduct of our analysis.

Theorem 2. *Let $m \geq Kd$ be some fixed integer. Let V_0 , λ , and w_0 satisfy the same assumptions as in Theorem 1. Then we have:*

- (i) *The steady state ρ_∞ is a positive trace-class operator on $L^2(\mathbb{R}^d)$, satisfying $\text{tr } \rho_\infty = 1$.*
- (ii) *Let $\rho \in C([0, \infty), \mathcal{T}_2)$ be the unique density matrix trajectory corresponding to the mild solution of (1.1). Then, the steady state ρ_∞ is exponentially stable, in the sense that*

$$\|\rho(t) - \rho_\infty\|_{\mathcal{T}_2} \leq (2\pi)^{\frac{d}{2}} \delta_m e^{-\kappa_m t} \|w_0 - w_\infty\|_{\mathcal{H}_m}, \quad \forall t \geq 0.$$

- (iii) *If the initial state $w_0 \in \mathcal{H}_m$ corresponds to a density matrix $\rho_0 \in \mathcal{T}_1^+$ (and hence w_0 is real valued, $\text{tr } \rho_0 \equiv \iint w_0 dx d\xi = 1$), then we also have*

$$\lim_{t \rightarrow \infty} \|\rho(t) - \rho_\infty\|_{\mathcal{T}_1} = 0.$$

Note that, in the presented framework, we do not obtain exponential convergence towards the steady state in the \mathcal{T}_1 -norm but only in the sense of Hilbert-Schmidt operators. This is due to the weak compactness methods involved in the proof of Gr\"umm's theorem (cf. the proof of Th. 2 in §6).

Remark 2.4. Consider now the following, more general quantum Fokker-Planck type operator replacing the r.h.s. of (1.1):

$$Qw := D_{\text{pp}} \Delta_\xi w + 2D_{\text{pq}} \text{div}_x (\nabla_\xi w) + 2D_{\text{f}} \text{div}_\xi (\xi w) + D_{\text{qq}} \Delta_x w.$$

It is straightforward to extend our results to this case as long as the *Lindblad condition* holds, i.e.

$$(2.11) \quad D_{\text{pp}} \geq 0, \quad D_{\text{pp}} D_{\text{qq}} - \left(D_{\text{pq}}^2 + \frac{D_{\text{f}}^2}{4} \right) \geq 0.$$

The modified quadratic function $A(x, \xi)$ is given in [35]. The Lindblad condition (2.11) implies that discarding in (1.1) the diffusion in x , and hence reducing the r.h.s. to the classical Fokker-Planck operator $Q_{cl}w := \Delta_\xi w + 2 \operatorname{div}_\xi(\xi w)$, would *not* describe a “correct” open quantum system. Nevertheless, this is a frequently used model in applications [39], yielding reasonable results in numerical simulations.

3. BASIC PROPERTIES OF THE UNPERTURBED OPERATOR L

3.1. Functional framework. It has been shown in [35] that the operator L , defined in (2.3), can be rewritten in the following form

$$(3.1) \quad Lw = \operatorname{div}(\nabla w + w(\nabla A + F)),$$

with

$$(3.2) \quad \operatorname{div}(Fe^{-A}) = \frac{1}{c} \operatorname{div}(F\mu) = 0.$$

Here and in the sequel, all differential operators act with respect to both x and ξ (if not indicated otherwise). In (3.1), the function A is defined by (2.7) and

$$(3.3) \quad F := \begin{pmatrix} -\xi \\ x + 2\xi \end{pmatrix} - \nabla A = \frac{1}{2} \begin{pmatrix} -x - 3\xi \\ x + \xi \end{pmatrix}.$$

The reason to do so is that (3.1) belongs to a class of non-symmetric Fokker-Planck operators considered in [8]. From this point of view, a natural functional space to study the unperturbed operator L is given by the following definition.

Definition 3.1. Let $\mathcal{H} := L^2(\mathbb{R}^{2d}, \mu^{-1} dx d\xi)$, equipped with the inner product

$$\langle f, g \rangle_{\mathcal{H}} = \iint_{\mathbb{R}^{2d}} \frac{f\bar{g}}{\mu} dx d\xi.$$

We can now decompose L into its symmetric and anti-symmetric part in \mathcal{H} , i.e.

$$(3.4) \quad L = L^s + L^{\text{as}},$$

where

$$(3.5) \quad L^s w := \operatorname{div}(\nabla w + w\nabla A), \quad L^{\text{as}} w := \operatorname{div}(Fw).$$

It has been shown in [35], that the following property holds:

$$(3.6) \quad L^s \mu = 0, \quad L^{\text{as}} \mu = 0.$$

where μ is the stationary state defined in (2.6). Next we shall properly define the operator L . To this end we first consider $L|_{C_0^\infty}$, which is closable (w.r.t. the \mathcal{H} -norm) since it is dissipative:

Lemma 3.2. $L|_{C_0^\infty}$ is dissipative, i.e. it satisfies $\operatorname{Re} \langle Lw, w \rangle_{\mathcal{H}} \leq 0$, for all $w \in C_0^\infty(\mathbb{R}^{2d})$.

Proof. Using $\nabla A = -\mu^{-1} \nabla \mu$ we have, on the one hand

$$\begin{aligned} \langle L^s w, w \rangle_{\mathcal{H}} &= \iint_{\mathbb{R}^{2d}} \frac{\bar{w}}{\mu} \operatorname{div}(\nabla w + w\nabla A) dx d\xi = \iint_{\mathbb{R}^{2d}} \frac{\bar{w}}{\mu} \operatorname{div} \left(\mu \nabla \left(\frac{w}{\mu} \right) \right) dx d\xi \\ &= - \iint_{\mathbb{R}^{2d}} \mu \left| \nabla \left(\frac{w}{\mu} \right) \right|^2 dx d\xi \leq 0. \end{aligned}$$

On the other hand, it follows from (3.2) that

$$w \operatorname{div} F = -\frac{w}{\mu} F \cdot \nabla \mu,$$

and thus

$$\operatorname{div}(Fw) = -\mu F \cdot \left(\frac{w}{\mu^2} \nabla \mu - \frac{\nabla w}{\mu} \right) = \mu F \cdot \nabla \left(\frac{w}{\mu} \right).$$

An easy calculation then shows

$$\begin{aligned} \operatorname{Re} \langle L^{\text{as}} w, w \rangle_{\mathcal{H}} &= \operatorname{Re} \iint_{\mathbb{R}^{2d}} \frac{\bar{w}}{\mu} \operatorname{div}(Fw) \, dx \, d\xi = \operatorname{Re} \iint_{\mathbb{R}^{2d}} \frac{\bar{w}}{\mu} F \cdot \nabla \left(\frac{w}{\mu} \right) \mu \, dx \, d\xi \\ &= -\frac{1}{2} \iint_{\mathbb{R}^{2d}} \left| \frac{w}{\mu} \right|^2 \operatorname{div}(F\mu) \, dx \, d\xi = 0, \end{aligned}$$

by (3.2). To sum up we have shown that $\operatorname{Re} \langle Lw, w \rangle_{\mathcal{H}} \leq 0$ holds. \square

The operator $L \equiv \overline{L|_{C_0^\infty}}$ is now closed, densely defined on \mathcal{H} and dissipative. Moreover one easily sees that $L^* = L^s - L^{\text{as}}$. The main goal of this section is to prove that L admits a spectral gap and is invertible on the orthogonal complement of its kernel. For the first property, we start showing that L is the generator of a C_0 -semigroup of contractions on \mathcal{H} . For this we recall the following result from [9].

Lemma 3.3. *Let the operator $P = p_2(x, \xi, \nabla_x, \nabla_\xi)$, where p_2 is a second order polynomial, be defined on the domain $\mathcal{D}(P) = C_0^\infty(\mathbb{R}^{2d})$. Then P is closable and $\overline{P|_{C_0^\infty}}$ is the maximum extension of P in $L^2(\mathbb{R}^{2d})$.*

Several variants of such a result (on different functional spaces) can be found in [2, 3]. We can use this result now in order to prove that L is the generator of a C_0 -semigroup.

Lemma 3.4. *L generates a C_0 -semigroup of contractions on \mathcal{H} .*

Proof. Defining $v := w/\sqrt{\mu}$ transforms the evolution problem

$$\partial_t w = Lw, \quad w|_{t=0} = w_0 \in \mathcal{H}$$

into its analog on $L^2(\mathbb{R}^{2d})$. The new unknown $v(t, x, \xi)$ then satisfies the following equation

$$\partial_t v = Hv, \quad v|_{t=0} = w_0/\sqrt{\mu},$$

where H is (formally) is given by

$$Hv = \Delta v + F \cdot \nabla v + Uv,$$

and the new “potential” $U = U(x, \xi)$ reads

$$U = \frac{1}{2} \Delta A - \frac{1}{4} |\nabla A|^2.$$

Defining H on $\mathcal{D}(H) = C_0^\infty(\mathbb{R}^{2d})$, we have

$$Lw = \sqrt{\mu} H \left(\frac{w}{\sqrt{\mu}} \right),$$

and thus the dissipativity of L on \mathcal{H} directly carries over to $H \equiv \overline{H|_{C_0^\infty}}$ on $L^2(\mathbb{R}^{2d})$.

Next, we consider $H^*|_{C_0^\infty}$, defined via $\langle Hf, g \rangle_{L^2} = \langle f, H^*g \rangle_{L^2}$, for $f, g \in C_0^\infty(\mathbb{R}^{2d})$.

Due to the definitions (2.7) and (3.3), the operator $H^*|_{C_0^\infty}$ is exactly of the form needed in order to apply Lemma 3.3. Thus, $H^* \equiv \overline{H^*|_{C_0^\infty}}$ is also dissipative (on all of its domain). Hence, the Lumer-Phillips Theorem (see [31], Section 1.4) implies that H is the generator of a C_0 -semigroup on $L^2(\mathbb{R}^{2d})$, denoted by e^{Ht} .

Reversing the transformation $w \rightarrow v$ then implies that L is the generator of the C_0 -semigroup U_t on \mathcal{H} , given by

$$U_t w_0 = \sqrt{\mu} e^{Ht} \left(\frac{w_0}{\sqrt{\mu}} \right).$$

This finishes the proof. \square

3.2. Semigroup properties on \mathcal{H} . The above lemma shows that the unperturbed WFP equation

$$\partial_t w = Lw, \quad w|_{t=0} = w_0 \in \mathcal{H},$$

has, for all $w_0 \in \mathcal{H}$, a unique mild solution $w \in C([0, \infty), \mathcal{H})$, where $w(t, x, \xi) = U_t w_0(x, \xi)$, with U_t defined above. Obviously we also have $U_t \mu = \mu$, by (2.4). Moreover, in [35] the Green's function of U_t was computed explicitly. It shows that U_t conserves mass, i.e.

$$\iint_{\mathbb{R}^{2d}} w(t, x, \xi) dx d\xi = \iint_{\mathbb{R}^{2d}} w_0(x, \xi) dx d\xi, \quad \forall t \geq 0.$$

Next, we define

$$(3.7) \quad \mathcal{H}^\perp := \{w \in \mathcal{H} : w \perp \mu\} \subset \mathcal{H},$$

which is a closed subset of \mathcal{H} . Note that $w \perp \mu$ simply means that

$$\langle w, \mu \rangle_{\mathcal{H}} \equiv \iint_{\mathbb{R}^{2d}} w(x, \xi) dx d\xi = 0.$$

Hence, we have for $w \in C_0^\infty(\mathbb{R}^{2d})$, using (3.5):

$$\langle L^{\text{as}} w, \mu \rangle_{\mathcal{H}} \equiv \iint_{\mathbb{R}^{2d}} L^{\text{as}} w(x, \xi) dx d\xi = 0.$$

Thus, $L^{\text{as}} : \mathcal{H}^\perp \cap \mathcal{D}(L^{\text{as}}) \rightarrow \mathcal{H}^\perp$. Moreover, $L^s : \mathcal{H}^\perp \cap \mathcal{D}(L^s) \rightarrow \mathcal{H}^\perp$, since \mathcal{H}^\perp is spanned by the eigenfunctions of L^s (except of μ). To sum up, the operators L^s and L^{as} are simultaneously reducible on the two subspaces $\mathcal{H} = \text{span}[\mu] \oplus \mathcal{H}^\perp$.

We also have that U_t maps \mathcal{H}^\perp into itself, since for $w_0 \in \mathcal{H}^\perp$ the conservation of mass implies

$$(3.8) \quad \langle U_t w_0, \mu \rangle_{\mathcal{H}} \equiv \iint_{\mathbb{R}^{2d}} w(t, x, \xi) dx d\xi = \iint_{\mathbb{R}^{2d}} w_0(x, \xi) dx d\xi = 0, \quad \forall t \geq 0.$$

Lemma 3.4 allows us to prove that L has a spectral gap in \mathcal{H} , in the sense that

$$(3.9) \quad \sigma(L) \setminus \{0\} \subset \{z \in \mathbb{C} : \text{Re } z \leq -\sigma\}.$$

Proposition 3.5. *It holds*

$$\|L^{-1}\|_{\mathcal{B}(\mathcal{H}^\perp)} \leq \frac{1}{\sigma},$$

where $\sigma > 0$ is defined in (2.8).

Proof. Condition (2.8) implies that L^s has a spectral gap of size $\sigma > 0$ (cf. §3.2 in [8], e.g.). Moreover, [8, Theorem 2.19] also yields exponential decay (with the same rate) for the non-symmetric WFP equation:

$$(3.10) \quad \|U_t(w_0 - \mu)\|_{\mathcal{H}} \leq e^{-\sigma t} \|w_0 - \mu\|_{\mathcal{H}}.$$

Here, $w_0 \in \mathcal{H}$ has to satisfy $\iint_{\mathbb{R}^{2d}} w_0 dx d\xi = \iint_{\mathbb{R}^{2d}} \mu dx d\xi = 1$. By the discussion above, we know that $L|_{\mathcal{H}^\perp}$ is the generator of $U_t|_{\mathcal{H}^\perp}$. Hence, (3.10) implies

$$(3.11) \quad \|(L - z)^{-1}\|_{\mathcal{B}(\mathcal{H}^\perp)} \leq \frac{1}{\text{Re } z + \sigma}, \quad \forall z \in \mathbb{C}, \text{Re } z > -\sigma,$$

which proves the assertion for $z = 0$. \square

As a final preparatory step in this section, we shall prove more detailed coercivity properties of L within \mathcal{H}^\perp . We shall denote $\mathcal{H}^1 := \{w \in \mathcal{H} : \nabla w \in \mathcal{H}\}$, and \mathcal{H}^{-1} will denote its dual.

Lemma 3.6. *In \mathcal{H}^\perp the operator L satisfies*

$$(3.12) \quad -\operatorname{Re} \langle Lw, w \rangle_{\mathcal{H}} \geq \sigma \|w\|_{\mathcal{H}}^2.$$

Similarly, there exists a constant $0 < \alpha < \sigma$, such that

$$(3.13) \quad -\operatorname{Re} \langle Lw, w \rangle_{\mathcal{H}} \geq \alpha \|w\|_{\mathcal{H}^1}^2, \quad \forall w \in \mathcal{H}^\perp \cap \mathcal{H}^1.$$

Proof. We shall use the weighted Poincaré inequality (see [8]): For any function $f \in L^2(\mathbb{R}^{2d}, \mu dx d\xi)$, such that $\iint_{\mathbb{R}^{2d}} f \mu dx d\xi = 0$, it holds:

$$(3.14) \quad \iint_{\mathbb{R}^{2d}} |f|^2 \mu dx d\xi \leq \frac{1}{\sigma} \iint_{\mathbb{R}^{2d}} \mu |\nabla f|^2 dx d\xi.$$

Estimate (3.12) then readily follows by setting $f = w/\mu$:

$$\operatorname{Re} \langle Lw, w \rangle_{\mathcal{H}} = - \iint_{\mathbb{R}^{2d}} \mu \left| \nabla \left(\frac{w}{\mu} \right) \right|^2 dx d\xi \leq -\sigma \iint_{\mathbb{R}^{2d}} \frac{|w|^2}{\mu} dx d\xi.$$

In order to prove assertion (3.13), we note that

$$\mu \left| \nabla \left(\frac{w}{\mu} \right) \right|^2 = \frac{|\nabla w|^2}{\mu} - 2d \frac{|w|^2}{\mu} - \operatorname{div} \left(|w|^2 \frac{\nabla \mu}{\mu^2} \right),$$

taking into account that $\Delta(\log \mu) = -2d$. Next, let $0 < \alpha < 1$ (to be chosen later), and write

$$\begin{aligned} \operatorname{Re} \langle Lw, w \rangle_{\mathcal{H}} &= -\alpha \iint_{\mathbb{R}^{2d}} \mu \left| \nabla \left(\frac{w}{\mu} \right) \right|^2 dx d\xi - (1-\alpha) \iint_{\mathbb{R}^{2d}} \mu \left| \nabla \left(\frac{w}{\mu} \right) \right|^2 dx d\xi \\ &= -\alpha \iint_{\mathbb{R}^{2d}} \frac{|\nabla w|^2}{\mu} dx d\xi + 2d\alpha \iint_{\mathbb{R}^{2d}} \frac{|w|^2}{\mu} dx d\xi \\ &\quad - (1-\alpha) \iint_{\mathbb{R}^{2d}} \mu \left| \nabla \left(\frac{w}{\mu} \right) \right|^2 dx d\xi. \end{aligned}$$

Inequality (3.14) for $f = w/\mu$ then implies:

$$-(1-\alpha) \iint_{\mathbb{R}^{2d}} \mu \left| \nabla \left(\frac{w}{\mu} \right) \right|^2 dx d\xi \leq -\sigma(1-\alpha) \iint_{\mathbb{R}^{2d}} \frac{|w|^2}{\mu} dx d\xi.$$

Therefore

$$\operatorname{Re} \langle Lw, w \rangle_{\mathcal{H}} \leq -\alpha \iint_{\mathbb{R}^{2d}} \frac{|\nabla w|^2}{\mu} dx d\xi + (2d\alpha - \sigma(1-\alpha)) \iint_{\mathbb{R}^{2d}} \frac{|w|^2}{\mu} dx d\xi.$$

The choice

$$\alpha = \frac{\sigma}{\sigma + 2d + 1}$$

yields assertion (3.13). \square

In the next section we shall study the operator L in the larger functional spaces \mathcal{H}_m (see Definition 2.2). This is necessary since the perturbation operator $\Theta[V_0]$ is unbounded in \mathcal{H} , even for $V_0 \in C_0^\infty(\mathbb{R}^d)$, cf. Remark 5.2.

4. STUDY OF THE UNPERTURBED PROBLEM IN \mathcal{H}_m

In this section, we adapt the general procedure outlined in [25, 30] to the specific model at hand. One of the main differences to the models studied in [25] is the fact that the WFP operator includes a diffusion in x . Nevertheless, we shall follow the main ideas of [25]. In a first step, this requires us to gain sufficient control on the action of U_t on \mathcal{H}_m . After that, we establish a new decomposition of L (not to be confused with the decomposition $L = L^s + L^{\text{as}}$ used above) in order to lift resolvent estimates onto the enlarged space $\mathcal{H}_m \supset \mathcal{H}$. Together with the Gearhart-Prüss Theorem (cf. Theorem V.1.11 in [20]), these estimates will finally allow us to infer exponential decay of U_t on \mathcal{H}_m .

4.1. Mathematical preliminaries. In (3.5) we decomposed the unperturbed evolution operator as $L = L^s + L^{\text{as}}$. As a first, from basic property of the spaces \mathcal{H}_m (see Definition 2.2), we note that L^{as} is still anti-symmetric in \mathcal{H}_m , $m \in \mathbb{N}$.

Lemma 4.1. *It holds*

$$(4.1) \quad \operatorname{Re} \langle L^{\text{as}} w, w \rangle_{\mathcal{H}_m} = 0, \quad \forall m \in \mathbb{N}.$$

Proof. A straightforward calculation yields $\operatorname{div}(F) = 0$. Hence

$$\begin{aligned} \operatorname{Re} \langle L^{\text{as}} w, w \rangle_{\mathcal{H}_m} &= \operatorname{Re} \iint_{\mathbb{R}^{2d}} \frac{\bar{w}}{\nu_m} \operatorname{div}(F w) \, dx \, d\xi = \frac{1}{2} \iint_{\mathbb{R}^{2d}} F \cdot \nabla (|w|^2) \nu_m^{-1} \, dx \, d\xi \\ &= -\frac{1}{2} \iint_{\mathbb{R}^{2d}} |w|^2 m A^{m-1} F \cdot \nabla A \, dx \, d\xi, \end{aligned}$$

after integrating by parts and using $\nu_m^{-1} = 1 + A^m(x, \xi)$. Now, it is easily seen from (3.3) that $F \cdot \nabla A = 0$, which implies (4.1). \square

The proof shows that, in the definition of \mathcal{H}_m , it is important to choose the weight ν_m as a (smooth) function of $A = -\log \frac{\mu}{c}$. Otherwise the fundamental property (4.1) would no longer be true. Also note that in contrast to L^{as} , the operator L^s is *not* symmetric in \mathcal{H}_m .

Before studying further properties of L in \mathcal{H}_m we state the following technical lemma. In order to keep the presentation simple, we shall not attempt to give the optimal constants.

Lemma 4.2. *Let $A = -\log \frac{\mu}{c}$, as given in (2.7). Then the following properties hold:*

(a) *There exists a constant $a_1 > 0$, such that for all $m \in \mathbb{N}$ it holds:*

$$a_1 (1 + A^m) \leq A^{m-1} |\nabla A|^2, \quad \forall |x|^2 + |\xi|^2 \geq 12.$$

(b) *Choosing $K := \frac{4}{a_1}$ it holds for all integer $m \geq Kd$:*

$$4d (1 + A^m) \leq m A^{m-1} |\nabla A|^2, \quad \forall |x|^2 + |\xi|^2 \geq 12.$$

(c) *There exists a constant $a_2 > 1$ such that*

$$|\nabla A|^2 \leq a_2 A, \quad \forall x, \xi \in \mathbb{R}^d.$$

(d) *For any $|x|, |\xi| \geq \frac{1}{\varepsilon}$ and $m \geq 1$, it holds*

$$\Delta (1 + A^m) \leq m A^{m-1} |\nabla A|^2 \varepsilon^2 6(m-1+3d).$$

Proof. Using Young's inequality we easily obtain

$$(4.2) \quad \frac{1}{12} (|x|^2 + |\xi|^2) \leq A(x, \xi) \leq |x|^2 + |\xi|^2,$$

$$(4.3) \quad \frac{1}{18} (|x|^2 + |\xi|^2) \leq |\nabla A(x, \xi)|^2 \leq 3(|x|^2 + |\xi|^2).$$

This yields assertion (c). To show (a), we note from (4.2) that

$$1 \leq A \leq A^m, \quad \forall |x|^2 + |\xi|^2 \geq 12.$$

Hence, we obtain with (4.2), (4.3):

$$1 + A^m \leq 2 A^m \leq 36 A^{m-1} |\nabla A|^2,$$

which is assertion (a). Finally, to prove assertion (d), we compute

$$\begin{aligned} \Delta (1 + A^m) &= m A^{m-1} |\nabla A|^2 \left(\frac{m-1}{A} + \frac{2d}{|\nabla A|^2} \right) \\ &\leq m A^{m-1} |\nabla A|^2 \varepsilon^2 6(m-1+3d), \quad \text{for } |x|, |\xi| \geq \frac{1}{\varepsilon}. \end{aligned}$$

□

Remark 4.3. Note that the constants $a_1 \geq \frac{1}{36}$, $a_2 \leq 36$, and $K = \frac{4}{a_1} \leq 144$ can be chosen *independent* of $m \in \mathbb{N}$ and of the spatial dimension $d \in \mathbb{N}$. Moreover, $K = 1$ is not possible for $d = 1$.

4.2. Semigroup properties on \mathcal{H}_m . Analogously to (3.7), we now define the following closed subset of \mathcal{H}_m :

$$\mathcal{H}_m^\perp := \{w \in \mathcal{H}_m : w \perp \nu_m\}, \quad m \in \mathbb{N},$$

which is again characterized by the zero-mass condition

$$(4.4) \quad \langle w, \nu_m \rangle_{\mathcal{H}_m} \equiv \iint_{\mathbb{R}^{2d}} w(x, \xi) \, dx \, d\xi = 0.$$

Thus we have $\mathcal{H}^\perp \subset \mathcal{H}_m^\perp \forall m \in \mathbb{N}$. As before, we define L on \mathcal{H}_m via $L \equiv \overline{L|_{C_0^\infty}}$ which yields a closed, densely defined operator on \mathcal{H}_m for each $m \in \mathbb{N}$. In addition, we also have the following result.

Lemma 4.4. *For each $m \in \mathbb{N}$, the operator L generates a C_0 -semigroup of bounded operators on \mathcal{H}_m , satisfying*

$$(4.5) \quad \|U_t\|_{\mathcal{B}(\mathcal{H}_m)} \leq e^{\beta_m t}, \quad \beta_m \in \mathbb{R}.$$

Proof. We compute

$$\begin{aligned} \operatorname{Re} \langle Lw, w \rangle_{\mathcal{H}_m} &= - \iint_{\mathbb{R}^{2d}} \frac{|\nabla w|^2}{\nu_m} \, dx \, d\xi \\ &\quad + \frac{1}{2} \iint_{\mathbb{R}^{2d}} |w|^2 \left(\Delta A^m - m A^{m-1} |\nabla A|^2 + \frac{2d}{\nu_m} \right) \, dx \, d\xi, \end{aligned}$$

by taking into account (4.1) and the fact that $\nu_m^{-1} = 1 + A^m(x, \xi)$. Using assertion (c) of Lemma 4.2, we can estimate

$$\Delta(A^m) = m A^{m-2} ((m-1) |\nabla A|^2 + 2d A) \leq m ((m-1) a_2 + 2d) A^{m-1}.$$

Moreover, since, for all $x, \xi \in \mathbb{R}^d$: $A^{m-1}(x, \xi) \leq 1 + A^m(x, \xi)$, we consequently obtain

$$\Delta(A^m) - m A^{m-1} |\nabla A|^2 + \frac{2d}{\nu_m} \leq \Delta(A^m) + \frac{2d}{\nu_m} \leq \beta_m (1 + A^m),$$

where

$$\beta_m := 2d + m((m-1)a_2 + 2d).$$

In summary, this yields

$$\operatorname{Re} \langle Lw, w \rangle_{\mathcal{H}_m} \leq \beta_m \|w\|_{\mathcal{H}_m}^2.$$

Thus, for the unperturbed evolution equation $\partial_t w = Lw$ we infer

$$\frac{d}{dt} \|w\|_{\mathcal{H}_m}^2 = 2 \operatorname{Re} \langle Lw, w \rangle_{\mathcal{H}_m} \leq 2 \beta_m \|w\|_{\mathcal{H}_m}^2,$$

and the assertion follows. □

Remark 4.5. Note that β_m *cannot be negative* in Lemma 4.4 since $U_t(\mu) = \mu$. However, using some refined estimates below, we shall find (see Proposition 4.8) that the restricted semigroup $U_t|_{\mathcal{H}_m^\perp}$ is exponentially decaying, provided $m \in \mathbb{N}$ is sufficiently large. To this end, we note that the two (non-orthogonal) subspaces $\mathcal{H}_m = \operatorname{span}[\mu] \oplus \mathcal{H}_m^\perp$ are invariant under L and under U_t ($\forall m \in \mathbb{N}$) due to mass conservation (3.8) and (4.4).

As a final preparatory step, we shall need the following decomposition result for L , where we denote

$$\mathcal{H}_m^1 := \{w \in \mathcal{H}_m : \nabla w \in \mathcal{H}_m\}.$$

Proposition 4.6. *Let $m \geq Kd$ be some fixed integer, and K was defined in Lemma 4.2. Then there exists an $0 < \varepsilon < 1$ such that the operator L can be split into $L = L_1^\varepsilon + L_2^\varepsilon$, with $L_1^\varepsilon, L_2^\varepsilon$ satisfying:*

- (1) $L_1^\varepsilon : \mathcal{H}_m \rightarrow \mathcal{H}_m$ is a closed and unbounded operator, while $L_1^\varepsilon : \mathcal{H}_m^1 \rightarrow \mathcal{H}$ and $L_1^\varepsilon : \mathcal{H}_m \rightarrow \mathcal{H}^{-1}$ are bounded operators.
- (2) $(L_2^\varepsilon - z) : \mathcal{H} \rightarrow \mathcal{H}$ and $(L_2^\varepsilon - z) : \mathcal{H}_m \rightarrow \mathcal{H}_m$ is a closed, unbounded and invertible operator for every $z \in \Omega := \{z \in \mathbb{C} : \operatorname{Re} z > -\Lambda_m\}$, where $\Lambda_m > 0$ is a positive constant defined in (A.3).
- (3) The operator

$$L_1^\varepsilon (L_2^\varepsilon - z)^{-1} : \mathcal{H}_m \rightarrow \mathcal{H} \subset \mathcal{H}_m$$

is bounded for any $z \in \Omega$.

The proof is lengthy and rather technical and therefore deferred to Appendix A.

Remark 4.7. Note that in Proposition 4.6, ε has to be chosen positive, in order to ensure assertion (1). In fact, while L_2^ε continues to be coercive also for $\varepsilon = 0$, the operators $L_1^\varepsilon : \mathcal{H}_m^1 \rightarrow \mathcal{H}$ and $L_1^\varepsilon : \mathcal{H}_m \rightarrow \mathcal{H}^{-1}$ become unbounded as $\varepsilon \rightarrow 0$. The fact that L_1^ε is bounded for $\varepsilon > 0$ is essential, in order to obtain the decay estimate (4.6), cf. the proof of Proposition 4.8.

Indeed, introducing the decomposition $L = L_1^\varepsilon + L_2^\varepsilon$ is one of the key ideas in [25, 30] in order to lift estimates for the resolvent $R(z) = (L - z)^{-1}$ onto the larger space \mathcal{H}_m . The general decomposition procedure introduced in [25] applies to the WFP equation and provides the following exponential decay of U_t on \mathcal{H}_m .

Proposition 4.8. *Let $\sigma > 0$ be the spectral gap of L^s in \mathcal{H} , and let $w_0 \in \mathcal{H}_m$ with $\iint_{\mathbb{R}^{2d}} w_0 \, dx d\xi = 1$. Then, for every integer $m \geq Kd$, it holds*

$$(4.6) \quad \|U_t(w_0 - \mu)\|_{\mathcal{H}_m} \leq \delta_m e^{-\tilde{\gamma}_m t} \|w_0 - \mu\|_{\mathcal{H}_m},$$

for any $\tilde{\gamma}_m \in (0, \gamma_m)$, where

$$(4.7) \quad \gamma_m := \min\{\Lambda_m; \sigma\} > 0,$$

and $\delta_m = \delta_m(\tilde{\gamma}_m) > 1$ is given in (4.15). Furthermore, we have for the resolvent set

$$(4.8) \quad \varrho(L|_{\mathcal{H}_m}) \supseteq \Omega_1 := \{z \in \mathbb{C} : \operatorname{Re} z > -\gamma_m, z \neq 0\}.$$

Proof. For the sake of completeness we briefly present the proof which follows the ones of Theorem 2.1, Theorem 3.1, and Theorem 4.1 in [25]. The spirit of the proof is the following: By using the operator factorization from Proposition 4.6, we shall infer an estimate for the resolvent on \mathcal{H}_m . Restricting the resolvent to \mathcal{H}_m^\perp removes its singularity at $z = 0$ and consequently yields a uniform estimate on the complex half plane $\{z \in \mathbb{C} : \operatorname{Re} z \geq -\tilde{\gamma}_m\}$. The Gearhart-Prüss Theorem [24, 33] then yields the exponential decay of U_t on \mathcal{H}_m^\perp . The proof now follows in several steps:

Step 1: Following [29] we define on \mathcal{H}_m the operator

$$(4.9) \quad R(z) := (L_2^\varepsilon - z)^{-1} - ((L - z)|_{\mathcal{H}})^{-1} L_1^\varepsilon (L_2^\varepsilon - z)^{-1}, \quad z \in \Omega_1.$$

Theorem 2.1 and Remark 2.2 in [25] implies that $R(z)$ is the inverse operator of $(L - z)$ in \mathcal{H}_m for any $z \in \Omega_1$ and therefore statement (4.8) holds.

Step 2: The next step is devoted to obtaining uniform estimates for $R(z) = ((L - z)|_{\mathcal{H}_m})^{-1}$, for $z \in \mathbb{C}$ on some appropriately defined half planes, cf. [25,

Theorem 3.1 (4)]. To this end, we shall first prove the following bound for the resolvent on \mathcal{H} :

$$(4.10) \quad \sup_s \|(L - (a + is))^{-1}\|_{\mathcal{B}(\mathcal{H})} = K_0 < \infty, \quad \forall a \in (-\sigma, 0).$$

Indeed, the constant $K_0 = K_0(\sigma, a)$ can be explicitly obtained by considering the resolvent equation for $\operatorname{Re} z > -\sigma$ and $z \neq 0$:

$$(L - z)f = g \quad \text{on } \mathcal{H},$$

Using the orthogonal decomposition $f = f^\perp + c_1\mu$, $g = g^\perp + c_2\mu$, having in mind that L maps \mathcal{H}^\perp into \mathcal{H}^\perp , we infer

$$f^\perp = (L - z)|_{\mathcal{H}^\perp}^{-1} g^\perp, \quad c_1 = -\frac{c_2}{z}.$$

In view of (3.11) this yields

$$\|f\|_{\mathcal{H}}^2 \leq \frac{1}{(\operatorname{Re} z + \sigma)^2} \|g^\perp\|_{\mathcal{H}}^2 + \frac{|c_2|^2}{|z|^2} \leq \max \left\{ \frac{1}{(\operatorname{Re} z + \sigma)^2}; \frac{1}{|z|^2} \right\} \|g\|_{\mathcal{H}}^2.$$

Hence,

$$\|(L - z)^{-1}\|_{\mathcal{B}(\mathcal{H})} \leq \max \left\{ \frac{1}{\operatorname{Re} z + \sigma}; \frac{1}{|z|} \right\} \quad \text{for } \operatorname{Re} z > -\sigma, z \neq 0;$$

and $K_0 \leq \max \left\{ \frac{1}{a+\sigma}; \frac{1}{|a|} \right\}$.

From this bound on $(L - z)|_{\mathcal{H}}^{-1}$ we can deduce a bound on $(L - z)|_{\mathcal{H}_m}^{-1}$ for $z \in \Omega_1 \equiv \{z \in \mathbb{C} : \operatorname{Re} z > -\gamma_m, z \neq 0\}$. Using (4.9), (A.2), and $L_1^\varepsilon \in \mathcal{B}(\mathcal{H}_m^1 \rightarrow \mathcal{H})$, we infer

$$(4.11) \quad \begin{aligned} & \|(L - z)^{-1}\|_{\mathcal{B}(\mathcal{H}_m)} \\ & \leq \|(L_2^\varepsilon - z)^{-1}\|_{\mathcal{B}(\mathcal{H}_m \rightarrow \mathcal{H}_m^1)} \left(1 + \|(L - z)^{-1}\|_{\mathcal{B}(\mathcal{H})} \|L_1^\varepsilon\|_{\mathcal{B}(\mathcal{H}_m^1 \rightarrow \mathcal{H})}\right) \\ & \leq \max \left\{ \frac{1}{\operatorname{Re} z + \Lambda_m}; \frac{1}{\Lambda_m} \right\} \left(1 + \max \left\{ \frac{1}{\operatorname{Re} z + \sigma}; \frac{1}{|z|} \right\} \|L_1^\varepsilon\|_{\mathcal{B}(\mathcal{H}_m^1 \rightarrow \mathcal{H})}\right) \\ & =: \vartheta_m(z). \end{aligned}$$

Next we consider the resolvent of $L|_{\mathcal{H}_m^\perp}$. First we note that both subspaces of $\mathcal{H}_m = \operatorname{span}[\mu] \oplus \mathcal{H}_m^\perp$ are invariant for $(L - z)^{-1}$ (cf. Remark 4.5). Hence $(L - z)|_{\mathcal{H}_m}^{-1}$ and $(L - z)|_{\mathcal{H}_m^\perp}^{-1}$ coincide on \mathcal{H}_m^\perp . Since $z = 0$ is an isolated and non-degenerate eigenvalue of $L|_{\mathcal{H}_m}$, we conclude $\sigma(L|_{\mathcal{H}_m^\perp}) = \sigma(L|_{\mathcal{H}_m}) \setminus \{0\}$. Since L generates a C_0 -semigroup on \mathcal{H}_m^\perp , it is closed and its resolvent is analytic on

$$\varrho(L|_{\mathcal{H}_m^\perp}) \supseteq \{z \in \mathbb{C} : \operatorname{Re} z > -\gamma_m\}.$$

For any fixed $\tilde{\gamma}_m \in (0, \gamma_m)$ we henceforth conclude from (4.11) that the resolvent of $L|_{\mathcal{H}_m^\perp}$ is uniformly bounded (on a whole right half space)

$$(4.12) \quad \|(L - z)^{-1}\|_{\mathcal{B}(\mathcal{H}_m^\perp)} \leq M(\tilde{\gamma}_m) < \infty, \quad \operatorname{Re} z \geq -\tilde{\gamma}_m.$$

Note, however, that the constant $M(\tilde{\gamma}_m)$ is not known explicitly.

Step 3: Next we shall show that this resolvent estimate yields an exponential decay estimate for the semigroup U_t on \mathcal{H}_m^\perp . In order to do so, we will apply the Gearhart-Prüss-Theorem to the rescaled semigroup $e^{\tilde{\gamma}_m t} U_t$, cf. Theorem V.1.11 in

[20] (see also [25, Theorem 3.1] and [26]). This is possible in view of the uniform bound (4.12) and yields the following estimate for U_t :

$$(4.13) \quad \|U_t\|_{\mathcal{B}(\mathcal{H}_m^\perp)} \leq \frac{(1 + M\omega)^2 C_L^2}{2\pi t} e^{-\tilde{\gamma}_m t}, \quad t > 0,$$

where $\omega > \beta_m + \tilde{\gamma}_m + 1$, $C_L \leq \pi(\omega - \beta_m - \tilde{\gamma}_m)^{-1}$, and

$$(4.14) \quad M := \sup_{s \in \mathbb{R}} \|(L - (-\tilde{\gamma}_m + is))^{-1}\|_{\mathcal{B}(\mathcal{H}_m^\perp)} \leq \vartheta_m(-\tilde{\gamma}_m),$$

where the second inequality follows directly from (4.11) and the definition of M . Note that (4.14) asserts a bound on the resolvent along the (fixed) line

$$\{z \in \mathbb{C} : z = -\tilde{\gamma}_m + is\},$$

in contrast to (4.12). Keeping this in mind, we conclude that the estimates in the proof of Theorem V.1.11 in [20] in fact only depend on the resolvent evaluated at $z = -\tilde{\gamma}_m + is$. Interpolating (4.13) with (4.5), we consequently conclude

$$\|U_t\|_{\mathcal{B}(\mathcal{H}_m^\perp)} \leq \delta_m e^{-\tilde{\gamma}_m t},$$

where (after optimizing in $\omega > \beta_m + \tilde{\gamma}_m + 1$)

$$(4.15) \quad \delta_m := \max \left\{ \frac{\pi}{2} \vartheta_m(-\tilde{\gamma}_m)^2; e^{\beta_m + \tilde{\gamma}_m} \right\} > 1.$$

This finishes the proof. \square

Proposition 4.8 implies that the operator L is invertible in the space \mathcal{H}_m^\perp , for $m \geq Kd$. More precisely, invoking classical arguments (cf. Theorem 1.5.3 in [31]), we infer

$$(4.16) \quad \forall m \geq Kd : \quad \|L^{-1}\|_{\mathcal{B}(\mathcal{H}_m^\perp)} \leq \frac{\delta_m}{\tilde{\gamma}_m} =: \frac{1}{\sigma_m}.$$

Note that the constant $\sigma_m > 0$ depends on m and thus on d and A . The reason why we obtain exponential decay of U_t on \mathcal{H}_m with a rate $\tilde{\gamma}_m < \gamma_m$ can be understood from the fact that in order to apply the Gearhart-Prüss Theorem one needs to guarantee a uniform resolvent estimates on some complex half plane, which is not sharp, in contrast to e.g. the Hille-Yoshida theorem (used on \mathcal{H}).

5. BOUNDEDNESS OF THE PERTURBATION

In this section we shall prove the boundedness of the operator $\Theta[V_0]$ in \mathcal{H}_m which is the key technical result for our perturbation analysis. We recall that our potential V from (2.1) consists of the harmonic potential plus the perturbation λV_0 . Hence, we shall now consider $-\lambda\Theta[V_0]$ as a perturbation of L .

Proposition 5.1. *Let $m \in \mathbb{N}$ and the potential $V_0 \in C_b^m(\mathbb{R}^d)$. Then the operator $\Theta[V_0]$ maps \mathcal{H}_m into \mathcal{H}_m^\perp and*

$$\|\Theta[V_0]\|_{\mathcal{B}(\mathcal{H}_m)} \leq \Gamma_m := C_m \max_{|j| \leq m} \|\partial_x^j V_0\|_{L^\infty(\mathbb{R}^d)},$$

where $C_m > 0$ denotes some positive constant, depending only on m and d .

Proof. We first note that, in view of (4.2), the norm $\|f\|_{\mathcal{H}_m}^2$ and the norm

$$\|f\|_m^2 := \iint_{\mathbb{R}^{2d}} |f|^2(x, \xi) \left(1 + (|x|^2 + |\xi|^2)^m\right) dx d\xi$$

are equivalent. It is therefore enough to prove that

$$(5.1) \quad \iint_{\mathbb{R}^{2d}} |\Theta[V_0]w|^2 \left(1 + (|x|^2 + |\xi|^2)^m\right) dx d\xi \leq C_0 \|w\|_m^2 :$$

for some $C_0 \geq 0$. In the following we denote by

$$(5.2) \quad (\mathcal{F}_{\xi \rightarrow \eta} w)(x, \eta) \equiv \widehat{w}(x, \eta) := \int_{\mathbb{R}^d} w(x, \xi) e^{-i\xi \cdot \eta} d\xi,$$

the partial Fourier transform with respect to the variable $\xi \in \mathbb{R}^d$ only. Recall from (1.3) that the operator $\Theta[V_0]$ acts via

$$(5.3) \quad \Theta[V_0]w = -i\mathcal{F}_{\eta \rightarrow \xi}^{-1}(\delta V_0(x, \eta) \cdot \mathcal{F}_{\xi \rightarrow \eta} w(x, \eta)).$$

Using Plancherel's formula and Hölder's inequality, this implies

$$\|\Theta[V_0]w\|_{L^2} \leq 2\|V_0\|_{L^\infty} \|w\|_{L^2}.$$

Thus, in order to prove (5.1), we only need to estimate

$$\iint_{\mathbb{R}^{2d}} |\Theta[V_0]w(x, \xi)|^2 (|x|^2 + |\xi|^2)^m dx d\xi.$$

We rewrite this term, using

$$(|x|^2 + |\xi|^2)^m = \sum_{j=0}^m \binom{m}{j} |x|^{2(m-j)} |\xi|^{2j},$$

in the following form:

$$\begin{aligned} & \iint_{\mathbb{R}^{2d}} |\Theta[V_0]w(x, \xi)|^2 (|x|^2 + |\xi|^2)^m dx d\xi \\ &= \sum_{j=0}^m \binom{m}{j} \iint_{\mathbb{R}^{2d}} |x|^{2(m-j)} |\xi|^{2j} |\Theta[V_0]w(x, \xi)|^2 dx d\xi. \end{aligned}$$

It holds

$$|\xi|^{2j} = \sum_{|n|=j} c_{n,j} \xi^{2n_1} \dots \xi^{2n_d},$$

where $c_{n,j}$ are some coefficients depending only on $n \in \mathbb{N}^d$. Therefore

$$\begin{aligned} & \iint_{\mathbb{R}^{2d}} |\Theta[V_0]w(x, \xi)|^2 (|x|^2 + |\xi|^2)^m dx d\xi \\ &= \sum_{j=0}^m \binom{m}{j} \left(\sum_{|n|=j} c_{n,j} \int_{\mathbb{R}^{2d}} |x|^{2(m-j)} \xi^{2n} |\Theta[V_0]w(x, \xi)|^2 dx d\xi \right), \end{aligned}$$

where we denote $\xi^{2n} := \xi^{2n_1} \dots \xi^{2n_d}$. From (5.2) we see that

$$(5.4) \quad \|\xi^n (\Theta[V_0]w)\|_{L^2(\mathbb{R}^{2d})}^2 = (2\pi)^{-d} \|\partial_\eta^n (\delta V_0 \widehat{w})\|_{L^2(\mathbb{R}^{2d})}^2,$$

with $\widehat{w}(x, \eta)$ defined by (5.2). We expand the right hand side of this identity by using the Leibniz formula (see also [28]), and we apply

$$\sup_{x, \eta \in \mathbb{R}^d} |\partial_{\eta_k}^j (\delta V_0)(x, \eta)| \leq 2^{(1-j)} \sup_{y \in \mathbb{R}^d} |\partial_{y_k}^j V_0(y)|,$$

(cf. Definition (1.4)). Then we can estimate (5.4) as follows:

$$\iint_{\mathbb{R}^{2d}} \xi^{2n} |\Theta[V_0]w(x, \xi)|^2 dx d\xi \leq C(n) \max_{|k| \leq |n|} \|\partial_x^k V_0\|_{L^\infty(\mathbb{R}^d)}^2 \|\xi^{n-k} w\|_{L^2(\mathbb{R}^{2d})}^2,$$

where $C(n) > 0$ depends only on binomial coefficients. In summary, we obtain

$$\begin{aligned} & \iint_{\mathbb{R}^{2d}} |\Theta[V_0]w(x, \xi)|^2 (|x|^2 + |\xi|^2)^m dx d\xi \\ & \leq \tilde{C}_m \sum_{j=0}^m \binom{m}{j} \left(\sum_{|n|=j} c_{n,j} \max_{|k| \leq |n|} \|\partial_x^k V_0\|_{L^\infty(\mathbb{R}^d)}^2 \| |x|^{m-j} \xi^{n-k} w \|_{L^2(\mathbb{R}^{2d})}^2 \right) \\ & \leq C_m^2 \max_{|j| \leq m} \|\partial_x^j V_0\|_{L^\infty(\mathbb{R}^d)}^2 \iint_{\mathbb{R}^{2d}} |w|^2(x, \xi) (|x|^2 + |\xi|^2)^m dx d\xi, \end{aligned}$$

with $\tilde{C}_m, C_m > 0$ depending only on binomial coefficients. Thus, the assertion is proved. \square

Remark 5.2. The unboundedness of $\Theta[V_0]$ in \mathcal{H} (the exponentially weighted Hilbert space) is due to its non-locality, which can be seen from the following reformulation of (1.3) (cf. §3 of [7]):

$$(\Theta[V]f)(x, \xi) = -2\pi^{-d} W(x, \xi) *_{\xi} f(x, \xi),$$

with

$$W(x, k) := \text{Im} [e^{2ix \cdot k} \hat{V}(2k)].$$

To illustrate the situation let us take $V_0(x) = \sin(x \cdot k_0)$ for some $k_0 \in \mathbb{R}^d \setminus \{0\}$. This implies

$$(\Theta[V]f)(x, \xi) = 2^d \cos(2x \cdot k_0) [f(x, \xi - k_0) - f(x, \xi + k_0)].$$

The problem is that such a shift operator cannot be bounded in an L^2 -space with an inverse Gaussian weight, as the following computation shows:

$$\int_{\mathbb{R}^d} |f(\xi - k_0)|^2 e^{|\xi|^2} d\xi = e^{|k_0|^2} \int_{\mathbb{R}^d} |f(\xi)|^2 e^{|\xi|^2} e^{2k_0 \cdot \xi} d\xi, \quad f \in L^2(\mathbb{R}^d, e^{|\xi|^2} d\xi).$$

6. PROOF OF THE MAIN THEOREMS

The results of the preceding sections allow us to give the proofs of Theorem 1 and Theorem 2.

Proof of Theorem 1. We start with Assertion (i): Let m be the integer fixed in the assertion. Any solution $w_\infty \in \mathcal{H}_m$ of (2.2) that is subject to the normalization $\iint w_\infty dx d\xi = 1$, satisfies the unique decomposition $w_\infty = \mu + w_*$ with $w_* \in \mathcal{H}_m^\perp$, i.e. $\iint w_* dx d\xi = 0$. Therefore, we consider the following fixed point iteration for w_* :

$$T : \mathcal{H}_m^\perp \rightarrow \mathcal{H}_m^\perp, \quad w_{n-1} \mapsto T(w_{n-1}) \equiv w_n,$$

where $w_n \in \mathcal{H}_m^\perp$ solves

$$Lw_n = \lambda \Theta[V_0](w_{n-1} + \mu).$$

To be able to apply Banach's fixed point theorem, we have to prove that the mapping T is a contraction on \mathcal{H}_m^\perp . To this end we write, for any $w_{n-1}, \tilde{w}_{n-1} \in \mathcal{H}_m^\perp$,

$$\|w_n - \tilde{w}_n\|_{\mathcal{H}_m^\perp} = \|\lambda L^{-1} \Theta[V_0](w_{n-1} - \tilde{w}_{n-1})\|_{\mathcal{H}_m^\perp}$$

and estimate

$$\|w_n - \tilde{w}_n\|_{\mathcal{H}_m^\perp} \leq |\lambda| \|L^{-1}\|_{\mathcal{B}(\mathcal{H}_m^\perp)} \|\Theta[V_0](w_{n-1} - \tilde{w}_{n-1})\|_{\mathcal{H}_m^\perp}.$$

From (4.16) and Proposition 5.1 we obtain

$$\|w_n - \tilde{w}_n\|_{\mathcal{H}_m^\perp} \leq \frac{\Gamma_m |\lambda|}{\sigma_m} \|w_{n-1} - \tilde{w}_{n-1}\|_{\mathcal{H}_m^\perp},$$

since the potential V_0 satisfies (2.9). Since $|\lambda| < \sigma_m/\Gamma_m$, there exists a unique fixed point $w_* = T(w_*) \in \mathcal{H}_m^\perp$. Thus, the unique (stationary) solution of (2.2) is obtained as $w_\infty = \mu + w_* \in \mathcal{H}_m$. Note, however, that $\mu \not\perp w_*$ in the sense of \mathcal{H}_m .

The obtained solution w_∞ is real valued, since T maps real valued functions to real valued functions. Moreover, $w_\infty \in \mathcal{H}_m$ satisfies (2.2), at least in the distributional sense. Moreover, $\Theta[V_0]w_\infty \in \mathcal{H}_m$ and $Lw_\infty \in H^{-2}(\mathbb{R}^{2d})$ and thus (2.2) also holds in $H^{-2}(\mathbb{R}^{2d})$. To explore *a posteriori* the regularity of w_∞ , we rewrite (2.2) in the following weak form

$$\iint_{\mathbb{R}^{2d}} (\nabla_x w_\infty \cdot \nabla_x \varphi + \nabla_\xi w_\infty \cdot \nabla_\xi \varphi + w_\infty \varphi) dx d\xi = {}_{H^{-1}} \langle F(w_\infty), \varphi \rangle_{H^1},$$

for any $\varphi \in H^1(\mathbb{R}^{2d})$, where

$$F(w_\infty) := w_\infty - \operatorname{div}_x(\xi w_\infty) + \operatorname{div}_\xi(x w_\infty + 2\xi w_\infty) - \lambda \Theta[V_0]w_\infty.$$

Clearly $F(w_\infty) \in H^{-1}(\mathbb{R}^{2d})$ and thus $w_\infty \in H^1(\mathbb{R}^{2d})$ follows. Moreover, since $F(w_\infty) \in L^2_{loc}(\mathbb{R}^{2d})$, we also have $w_\infty \in H^2_{loc}(\mathbb{R}^{2d})$.

For the proof of Assertion (ii), we first note that $\Theta[V_0]$ is a bounded perturbation of L on \mathcal{H}_m and thus (1.1) admits a unique mild solution $w \in C([0, \infty), \mathcal{H}_m)$. Since $\Theta[V_0]$ maps \mathcal{H}_m into \mathcal{H}_m^\perp , we also know that along this solution the mass is conserved, i.e. $\iint w(t) dx d\xi = 1$, for all $t \geq 0$.

Next, consider the new unknown $g(t) := w(t) - w_\infty$ with $g_0 = w_0 - w_\infty$. Due to mass conservation $g(t) \in \mathcal{H}_m^\perp$ for all $t \geq 0$, and we also have

$$\partial_t g = Lg - \lambda \Theta[V_0]g,$$

since w_∞ is a stationary solution of (1.1). Taking into account that the semigroup U_t associated with L in the space \mathcal{H}_m satisfies (4.6), it holds:

$$\begin{aligned} \|g(t)\|_{\mathcal{H}_m} &\leq \delta_m e^{-\tilde{\gamma}_m t} \|g_0\|_{\mathcal{H}_m} + \delta_m |\lambda| \int_0^t e^{-\tilde{\gamma}_m(t-s)} \|\Theta[V_0]g(s)\|_{\mathcal{H}_m} ds \\ &\leq \delta_m e^{-\tilde{\gamma}_m t} \|g_0\|_{\mathcal{H}_m} + \delta_m |\lambda| \Gamma_m \int_0^t e^{-\tilde{\gamma}_m(t-s)} \|g(s)\|_{\mathcal{H}_m} ds, \end{aligned}$$

for any $\tilde{\gamma}_m \in (\delta_m |\lambda| \Gamma_m, \gamma_m)$ and with Γ_m defined in Proposition 5.1. Gronwall's lemma then implies

$$\|g(t)\|_{\mathcal{H}_m} \leq \delta_m e^{-t(\tilde{\gamma}_m - \delta_m |\lambda| \Gamma_m)} \|g_0\|_{\mathcal{H}_m}.$$

It remains to prove Assertion (iii): As before we write $w_\infty = \mu + w_*$, where $w_* \in \mathcal{H}_m^\perp$ solves

$$(L - \lambda \Theta[V_0]) w_* = \lambda \Theta[V_0] \mu.$$

Since $\|\Theta[V_0]\|_{\mathcal{B}(\mathcal{H}_m)} \leq \Gamma_m$, we obtain $\|\lambda \Theta[V_0] \mu\|_{\mathcal{H}_m} \leq |\lambda| \Gamma_m \|\mu\|_{\mathcal{H}_m}$. Next, we consider L on \mathcal{H}_m^\perp . From the proof of Proposition 4.8 we conclude for its resolvent set:

$$\varrho \left(L \Big|_{\mathcal{H}_m^\perp} \right) \supseteq \{z \in \mathbb{C} : \operatorname{Re} z > -\gamma_m\},$$

and thus

$$\varrho \left((L - \lambda \Theta[V_0]) \Big|_{\mathcal{H}_m^\perp} \right) \supseteq \{z \in \mathbb{C} : \operatorname{Re} z > |\lambda| \Gamma_m - \gamma_m\}.$$

Since $|\lambda| \Gamma_m - \gamma_m < 0$ (see (2.10)) we have

$$(L - \lambda \Theta[V_0])^{-1} = L^{-1} (\operatorname{Id} - \lambda \Theta[V_0] L^{-1})^{-1} \quad (\text{on } \mathcal{H}_m^\perp).$$

Using (4.16) and $|\lambda|\Gamma_m < \sigma_m$ (see (2.10)) we conclude

$$\|(L - \lambda\Theta[V_0])\Big|_{\mathcal{H}_m^\perp}^{-1}\|_{\mathcal{B}(\mathcal{H}_m^\perp)} \leq \frac{1}{\sigma_m} \frac{1}{1 - |\lambda|\Gamma_m \frac{1}{\sigma_m}} = \frac{1}{\sigma_m - |\lambda|\Gamma_m}.$$

Thus, by writing

$$w_* = \left((L - \lambda\Theta[V_0])\Big|_{\mathcal{H}_m^\perp} \right)^{-1} (\lambda\Theta[V_0]\mu),$$

we infer

$$\|w_*\|_{\mathcal{H}_m} \equiv \|w_\infty - \mu\|_{\mathcal{H}_m} \leq \frac{|\lambda|\Gamma_m \|\mu\|_{\mathcal{H}_m}}{\sigma_m - |\lambda|\Gamma_m}$$

and the assertion is proved. \square

Remark 6.1. Due to the mass normalization $\iint w_\infty dx d\xi = \iint \mu dx d\xi = 1$, the fixed point w_* must take both positive and negative values. Thus, $w_\infty = \mu + w_*$ may, in general, also take negative values.

Proof of Theorem 2. We start with assertion (ii), which follows from the fact that

$$\|\rho\|_{\mathcal{T}_2} = \|\rho(\cdot, \cdot)\|_{L^2} = (2\pi)^{d/2} \|w(\cdot, \cdot)\|_{L^2} \leq (2\pi)^{d/2} \|w(\cdot, \cdot)\|_{\mathcal{H}_m}, \quad \forall \rho \in \mathcal{T}_2.$$

Thus, we infer

$$\rho(t) \xrightarrow{t \rightarrow \infty} \rho_\infty \quad \text{in } \mathcal{T}_2,$$

with the exponential rate obtained from Theorem 1 (ii).

To prove assertion (i) we consider the transient equation (1.1) as an auxiliary problem: Choose any $\rho_0 \in \mathcal{T}_1^+$ such that $\text{tr } \rho_0 = 1$ and the corresponding $w_0 \in \mathcal{H}_m$. Due to the results on the linear Cauchy problem given in [9] we know that (1.1) gives rise to a unique mild solution $\rho \in C([0, \infty); \mathcal{T}_1^+)$, satisfying $\text{tr } \rho(t) = 1$, for all $t \geq 0$. Hence, the trajectory $\{\rho(t), t \geq 0\}$ is bounded in \mathcal{T}_1 . Since \mathcal{T}_1 has a predual, i.e. the compact operators on $L^2(\mathbb{R}^d)$, the Banach-Alaoglu Theorem then asserts the existence of a sequence $\{t_n\}_{n \in \mathbb{N}} \subset \mathbb{R}_+$ with $t_n \rightarrow \infty$, such that

$$\rho(t_n) \xrightarrow{n \rightarrow \infty} \tilde{\rho} \quad \text{in } \mathcal{T}_1 \text{ weak-}\star$$

for some limiting $\tilde{\rho} \in \mathcal{T}_1$. The already obtained \mathcal{T}_2 -convergence of $\rho(t)$ towards $\rho_\infty \in \mathcal{T}_2$ implies $\rho_\infty = \tilde{\rho} \in \mathcal{T}_1$. And the *uniqueness* of the steady state yields the convergences of the whole t -dependent function $\rho(t) \rightarrow \tilde{\rho}$ in \mathcal{T}_1 weak- \star . Finally, we also conclude positivity of the operator ρ_∞ by the \mathcal{T}_2 -convergence and the fact that we already know from [9]: $\rho(t) \geq 0$, for all $t \geq 0$.

It remains to prove $\text{tr } \rho_\infty = 1$. To this end, we recall that for any $\rho \in \mathcal{T}_1^+$ the corresponding kernel

$$\vartheta(x, \eta) := \rho\left(x + \frac{\eta}{2}, x - \frac{\eta}{2}\right)$$

satisfies $\vartheta \in C(\mathbb{R}_\eta^d, L_+^1(\mathbb{R}_x^d))$, see [1], and it also holds

$$(6.1) \quad \text{tr } \rho = \int_{\mathbb{R}^d} \vartheta(x, 0) dx.$$

Further, note that $\vartheta(x, \eta) = (\mathcal{F}_{\xi \rightarrow \eta} w)(x, \eta) \equiv \hat{w}(x, \eta)$, by (1.2). On the other hand, for any $w \in \mathcal{H}_m$ we know that $\hat{w} \in C(\mathbb{R}_\eta^d, L^1(\mathbb{R}_x^d))$, due to the polynomial L^2 -weight ν_m^{-1} in $x \in \mathbb{R}^d$ and a simple Sobolev imbedding w.r.t. the variable $\eta \in \mathbb{R}^d$ (for both embeddings we used $m > \frac{d}{2}$). Hence the normalization condition $\iint w_\infty dx d\xi = 1$ implies $\text{tr } \rho_\infty = 1$, via (6.1), and assertion (ii) is proved.

Finally, we prove claim (iii) by first noting that the \mathcal{T}_2 -convergence of $\rho(t)$ implies convergence in the strong operator topology. Thus, having in mind that $\|\rho(t)\|_{\mathcal{T}_1} = \|\rho_0\|_{\mathcal{T}_1} = 1$, we infer from Gr\"umm's theorem (Th. 2.19 in [34]) that $\rho(t)$ also converges in the \mathcal{T}_1 -norm towards ρ_∞ . This concludes the proof of Theorem 2. \square

APPENDIX A. PROOF OF PROPOSITION 4.6

The proof will be divided into several steps:

Step 1: Let $\chi \in C_0^\infty(\mathbb{R}^{2d})$ be such that $\chi = 1$ on $B_1(0)$, with $\text{supp}(\chi) \subseteq B_2(0)$, $\|\nabla\chi\|_{L^\infty} \leq \sqrt{2}$, and let $\chi_\varepsilon(y) := \chi(\varepsilon y)$, for any $y = (x, \xi) \in \mathbb{R}^{2d}$, $0 < \varepsilon < 1$. We define

$$L_1^\varepsilon w := (d w - \nu_m \nabla \nu_m^{-1} \cdot \nabla w) \chi_\varepsilon,$$

as well as

$$\begin{aligned} L_2^\varepsilon w &:= \nu_m \operatorname{div}(\nu_m^{-1} \nabla w) + \nabla w \cdot \nabla A + d w + \operatorname{div}(w F) \\ &\quad + (d w - \nu_m \nabla \nu_m^{-1} \cdot \nabla w) (1 - \chi_\varepsilon). \end{aligned}$$

It is easily seen that L_1^ε indeed satisfies property (1).

Step 2: In order to prove properties (2) and (3), we need to show that

$$(L_2^\varepsilon - z)^{-1} : \mathcal{H}_m \rightarrow \mathcal{H}_m^1,$$

is bounded for $z \in \Omega \subset \mathbb{C}$. To this end, it suffices to show that $(L_2^\varepsilon - z)$ satisfies $\forall z \in \Omega$:

$$(A.1) \quad -\operatorname{Re} \langle (L_2^\varepsilon - z)w, w \rangle_{\mathcal{H}_m} \geq c \|w\|_{\mathcal{H}_m^1}^2,$$

with some $c = c(\operatorname{Re} z) > 0$. Indeed, suppose (A.1) holds for all (complex valued) $w \in \mathcal{H}_m^2 := \{w \in \mathcal{H}_m : \nabla w, (\nabla A + F) \cdot \nabla w, \Delta w \in \mathcal{H}_m\}$. Then, $L_2^\varepsilon - z$ is densely defined on $\mathcal{D}(L_2^\varepsilon) := \mathcal{H}_m^2 \subset \mathcal{H}_m$ and dissipative. Hence, it has a maximal dissipative (and thus surjective on \mathcal{H}_m) extension, which we shall consider in the sequel. From (A.1) we conclude

$$c \|w\|_{\mathcal{H}_m^1}^2 \leq -\operatorname{Re} \langle (L_2^\varepsilon - z)w, w \rangle_{\mathcal{H}_m} \leq \|w\|_{\mathcal{H}_m^1} \|(L_2^\varepsilon - z)w\|_{\mathcal{H}_m}, \quad c > 0,$$

which implies

$$(A.2) \quad \|(L_2^\varepsilon - z)^{-1} \tilde{w}\|_{\mathcal{H}_m^1} \leq \frac{1}{c} \|\tilde{w}\|_{\mathcal{H}_m}, \quad \text{for all } \tilde{w} \in \mathcal{H}_m.$$

For future reference we note that this bound will turn out to be uniform on the lines $z = a + is$, $s \in \mathbb{R}$ (with fixed $a > -\Lambda_m$), since $c = c(\operatorname{Re} z)$.

Step 3: Now, we have to prove (A.1). To this end, we decompose

$$\begin{aligned} \operatorname{Re} \langle L_2^\varepsilon w, w \rangle_{\mathcal{H}_m} &= - \iint_{\mathbb{R}^{2d}} (1 + A^m) |\nabla w|^2 \, dx \, d\xi \\ &\quad - \frac{m}{2} \iint_{\mathbb{R}^{2d}} |w|^2 A^{m-1} |\nabla A|^2 \, dx \, d\xi \\ &\quad + d \iint_{\mathbb{R}^{2d}} |w|^2 (1 + A^m) (1 - \chi_\varepsilon) \, dx \, d\xi \\ &\quad + \frac{1}{2} \iint_{\mathbb{R}^{2d}} |w|^2 \nabla(1 - \chi_\varepsilon) \cdot \nabla(1 + A^m) \, dx \, d\xi \\ &\quad + \frac{1}{2} \iint_{\mathbb{R}^{2d}} |w|^2 (1 - \chi_\varepsilon) \Delta(1 + A^m) \, dx \, d\xi \\ &=: I_1 + I_2 + I_3 + I_4 + I_5. \end{aligned}$$

Using Lemma 4.2 (b), we readily obtain for $\varepsilon \leq \frac{1}{\sqrt{3}}$:

$$\begin{aligned} |I_3| &\leq \frac{m}{4} \iint_{\mathbb{R}^{2d}} |w|^2 A^{m-1} |\nabla A|^2 (1 - \chi_\varepsilon) \, dx \, d\xi \\ &\leq \frac{m}{4} \iint_{\mathbb{R}^{2d}} |w|^2 A^{m-1} |\nabla A|^2 \, dx \, d\xi. \end{aligned}$$

In order to treat the term I_4 , we note that

$$|\nabla \chi_\varepsilon| = \varepsilon |\nabla \chi|, \quad |\nabla \chi| \leq \sqrt{2}, \quad \text{supp } \{\nabla \chi_\varepsilon\} \subset \left\{ y \in \mathbb{R}^{2d} \mid \frac{1}{\varepsilon} \leq |y| \leq \frac{2}{\varepsilon} \right\}.$$

Therefore, by (4.3),

$$\frac{1}{|\nabla A|} \leq \sqrt{18} \varepsilon \quad \text{for } y = (x, \xi) \in \text{supp } \{\nabla \chi_\varepsilon\}.$$

With $\varepsilon < 1$ this allows us to estimate

$$\begin{aligned} |\nabla(1 - \chi_\varepsilon) \cdot \nabla(1 + A^m)| &\leq \varepsilon m A^{m-1} |\nabla \chi| |\nabla A| \\ &\leq 6\varepsilon^2 m A^{m-1} |\nabla A|^2 \leq 6\varepsilon m A^{m-1} |\nabla A|^2, \end{aligned}$$

which implies

$$|I_4| \leq 3m\varepsilon \iint_{\mathbb{R}^{2d}} |w|^2 A^{m-1} |\nabla A|^2 \, dx \, d\xi.$$

The term I_5 can be estimated using Lemma 4.2 (d):

$$\begin{aligned} I_5 &= \frac{1}{2} \iint_{\mathbb{R}^{2d}} |w|^2 (1 - \chi_\varepsilon) \Delta(1 + A^m) \, dx \, d\xi \\ &\leq 3m\varepsilon^2 (m - 1 + 3d) \iint_{\mathbb{R}^{2d}} |w|^2 (1 - \chi_\varepsilon) A^{m-1} |\nabla A|^2 \, dx \, d\xi \\ &\leq 3m\varepsilon^2 (m - 1 + 3d) \iint_{\mathbb{R}^{2d}} |w|^2 A^{m-1} |\nabla A|^2 \, dx \, d\xi. \end{aligned}$$

In summary, we obtain

$$\frac{1}{4} I_2 + |I_4| \leq -\left(\frac{m}{8} - 3m\varepsilon\right) \iint_{\mathbb{R}^{2d}} |w|^2 A^{m-1} |\nabla A|^2 \, dx \, d\xi,$$

and

$$\frac{3}{4} I_2 + I_3 + I_5 \leq -\left(\frac{m}{8} - 3m\varepsilon^2 (m - 1 + 3d)\right) \iint_{\mathbb{R}^{2d}} |w|^2 A^{m-1} |\nabla A|^2 \, dx \, d\xi.$$

Now choosing $\varepsilon \leq \min\left\{\frac{1}{24}; \frac{1}{12\sqrt{m}}\right\}$, we can estimate (using $m \geq d$)

$$m \left(\frac{1}{8} - 3\varepsilon\right) \geq 0, \quad m \left(\frac{1}{8} - 3\varepsilon^2 (m - 1 + 3d)\right) \geq \frac{m}{24}.$$

Therefore

$$\begin{aligned} -\text{Re} \langle L_2^\varepsilon w, w \rangle_{\mathcal{H}_m} &\geq \iint_{\mathbb{R}^{2d}} (1 + A^m) |\nabla w|^2 \, dx \, d\xi \\ &\quad + \left(\frac{m}{8} - 2\right) \iint_{\mathbb{R}^{2d}} |w|^2 A^{m-1} |\nabla A|^2 \, dx \, d\xi, \end{aligned}$$

which we estimate further using Lemma 4.2 (b):

$$\begin{aligned} -\text{Re} \langle L_2^\varepsilon w, w \rangle_{\mathcal{H}_m} &\geq \iint_{\mathbb{R}^{2d}} (1 + A^m) |\nabla w|^2 \, dx \, d\xi \\ &\quad + \frac{d}{6} \iint_{\mathbb{R}^{2d}} |w|^2 (1 + A^m) (1 - \chi_{1/\sqrt{12}}) \, dx \, d\xi. \end{aligned}$$

This establishes the desired estimate outside of $(x, \xi) \in B_{1/\sqrt{12}}(0)$. In order to take into account the contribution near $|y| = 0$, we consider

$$\iint_{\mathbb{R}^{2d}} (1 + A^m) |\nabla w|^2 \, dx \, d\xi.$$

Applying Sobolev's inequality we obtain for any $d > 1$:

$$\iint_{\mathbb{R}^{2d}} (1 + A^m) |\nabla w|^2 \, dx \, d\xi \geq \iint_{\mathbb{R}^{2d}} |\nabla w|^2 \, dx \, d\xi \geq C_d^2 \|w\|_{L^q(\mathbb{R}^{2d})}^2,$$

with $q = \frac{4d}{2d-2}$. This can be estimated further via

$$C_d^2 \|w\|_{L^q(\mathbb{R}^{2d})}^2 \geq C_d^2 \|w\|_{L^q(B_{1/\sqrt{12}}(0))}^2 \geq \frac{C_1^2}{\|1 + A^m\|_{L^\infty(B_{1/\sqrt{12}}(0))}} \|w\|_{\mathcal{H}_m(B_{1/\sqrt{12}}(0))}^2,$$

where C_1 depends only on the Sobolev constant C_d and on the measure of the ball $B_{1/\sqrt{12}}(0)$. Finally, in order to deal with $d = 1$ we apply Cauchy-Schwarz to obtain

$$\begin{aligned} \iint_{\mathbb{R}^2} |\nabla w| \, dx \, d\xi &= \iint_{\mathbb{R}^2} |\nabla w| (1 + A^m)^{1/2} (1 + A^m)^{-1/2} \, dx \, d\xi \\ &\leq \left(\iint_{\mathbb{R}^2} |\nabla w|^2 (1 + A^m) \, dx \, d\xi \right)^{1/2} \left(\iint_{\mathbb{R}^2} (1 + A^m)^{-1} \, dx \, d\xi \right)^{1/2}. \end{aligned}$$

By assumption we have $m > 1$ (cf. Remark 4.3). Hence, the second factor on the r.h.s. is a finite constant, denoted by $C_{A,m}$. Applying again Sobolev's inequality (for $d = 1$), we obtain

$$\|\nabla w\|_{L^1(\mathbb{R}^2)} \geq \tilde{C}_1 \|w\|_{L^2(\mathbb{R}^2)} \geq \frac{\tilde{C}_1}{\|1 + A^m\|_{L^\infty(B_{1/\sqrt{12}}(0))}^{1/2}} \|w\|_{\mathcal{H}_m(B_{1/\sqrt{12}}(0))}.$$

By combining all the above estimates we infer for all (complex valued) $w \in \mathcal{H}_m^2$:

$$-\operatorname{Re} \langle L_2^\varepsilon w, w \rangle_{\mathcal{H}_m} \geq \Lambda_m \langle w, w \rangle_{\mathcal{H}_m^1},$$

where $\Lambda_m > 0$ is given by

$$(A.3) \quad \Lambda_m := \min \left\{ \frac{C_1^2}{2\|1 + A^m\|_{L^\infty(B_{1/\sqrt{12}}(0))}}; \frac{\tilde{C}_1^2}{2C_{A,m}^2\|1 + A^m\|_{L^\infty(B_{1/\sqrt{12}}(0))}}; \frac{1}{2}; \frac{d}{6} \right\}.$$

In summary, inequality (A.1) holds on \mathcal{H}_m^2 for any $\operatorname{Re} z > -\Lambda_m$. And we easily see that $c(z) = \min\{\Lambda_m; \Lambda_m + \operatorname{Re} z\}$. Hence, the operator $(L_2^\varepsilon - z)$ is invertible in \mathcal{H}_m^1 . And we have to choose $\varepsilon = \varepsilon(m) > 0$ such that

$$(A.4) \quad \varepsilon \leq \min \left\{ \frac{1}{12\sqrt{m}}; \frac{1}{24} \right\}.$$

□

REFERENCES

1. A. Arnold, *Self-Consistent Relaxation-Time Models in Quantum Mechanics*, Comm. Part. Diff. Equ. **21** (1996), no. 3/4, 473–506.
2. A. Arnold, J. Carrillo, and E. Dhamo, *On the periodic Wigner-Poisson-Fokker-Planck system*, J. Math. Anal. Appl. **275** (2002), 263–276.
3. A. Arnold, E. Dhamo, and C. Manzini, *The Wigner-Poisson-Fokker-Planck system: global-in-time solutions and dispersive effects*, Annales de l'IHP (C) - Analyse non lineaire **24** (2007), no. 4, 645–676.
4. A. Arnold, E. Dhamo, and C. Manzini, *Dispersive effects in quantum kinetic equations*, Indiana Univ. Math. J. **56** (2007), no. 3, 1299–1332.

5. A. Arnold, F. Fagnola, and L. Neumann, *Quantum Fokker-Planck models: the Lindblad and Wigner approaches*, in “Quantum Probability and related Topics - Proceedings of the 28th Conference” (Series: QP-PQ: Quantum Probability and White Noise Analysis - Vol. 23), J.C. Garcia, R. Quezada, S.B. Sontz (Eds.), World Scientific 2008.
6. A. Arnold, H. Lange, and P.F. Zweifel, *A discrete-velocity, stationary Wigner equation*, J. Math. Phys. **41** (2000), no. 11, 7167–7180.
7. A. Arnold, J.L. López, P. Markowich, and J. Soler, *An Analysis of Quantum Fokker-Planck Models: A Wigner Function Approach*, Rev. Mat. Iberoam. **20** (2004), no. 3, 771–814.
8. A. Arnold, P. Markowich, G. Toscani, and A. Unterreiter, *On convex Sobolev inequalities and the rate of convergence to equilibrium for Fokker-Planck type equations*, Comm. Partial Diff. Equ. **26** (2001), 43–100.
9. A. Arnold and C. Sparber, *Quantum dynamical semigroups for diffusion models with Hartree interaction*, Comm. Math. Phys. **251** (2004), no.1, 179–207.
10. R. Bosi, *Classical limit for linear and nonlinear quantum Fokker-Planck systems*, Commun. Pure Appl. Anal. **8** (2009), no. 3, 845–870.
11. F. Castella, L. Erdős, F. Frommlet, and P. Markowich, *Fokker-Planck equations as Scaling Limit of Reversible Quantum Systems*, J. Stat. Physics **100** (2000), no. 3/4, 543-601.
12. J. A. Cañizo, J. L. López, and J. Nieto, *Global L^1 theory and regularity for the 3D nonlinear Wigner-Poisson-Fokker-Planck system*, J. Diff. Equ. **198** (2004), 356–373.
13. P. R. Chernoff, *Perturbations of Dissipative Operators with Relative Bound One*, Proc. Am. Math. Soc. **33** (1972), no. 1, 72–74.
14. E. B. Davies, *Quantum Theory of Open Systems*, Academic Press (1976).
15. H. Dekker, *Quantization of the linearly damped harmonic oscillator*, Phys. Rev. A **16** (1977), 2126–2134.
16. L. Diósi, *On high-temperature Markovian equations for quantum Brownian motion*, Europhys. Lett. **22** (1993), 1-3.
17. A. Donarini, T. Novotný, and A.P. Jauho, *Simple models suffice for the single-dot quantum shuttle*, New J. of Physics **7** (2005), 237–262.
18. J.-P. Eckmann and M. Hairer, *Spectral Properties of Hypocoelliptic Operators*, Commun. Math. Phys. **235**, no. 2, 233–253.
19. M. Elk and P. Lambropoulos, *Connection between approximate Fokker-Planck equations and the Wigner function applied to the micromaser*, Quantum Semiclass. Opt. **8** (1996), 23–37.
20. K.J. Engel and R. Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Springer-Verlang, New York, 2000.
21. F. Fagnola and R. Rebolledo, *On the existence of stationary states for quantum dynamical semigroups*, J. Math. Phys. **42** (2001), 1296–1308.
22. F. Fagnola and R. Rebolledo, *Algebraic conditions for convergence of a quantum Markov semigroup to a steady state*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **11**(3) (2008), 467–474.
23. A. Frigerio and M. Verri, *Long-Time Asymptotic Properties of Dynamical Semigroups on W^* -algebras*, Math. Z. **180** (1982), 275–286.
24. L. Gearhart, *Spectral theory for contraction semigroups on Hilbert space*, Trans. Amer. Math. Soc. **236** (1978), 385394.
25. M.P. Gualdani, S. Mischler and C. Mouhot. *Factorization for non-symmetric operators and exponential H -theorem*. Preprint arxiv.org/pdf/1006.5523.
26. B. Helffer and J. Sjöstrand, *From resolvent bounds to semigroups bounds*. Preprint arxiv.org/pdf/1001.4171v1.
27. F. Hérau and F. Nier, *Isotropic hypoellipticity and trend to the equilibrium for the Fokker-Planck equation with high degree potential*, Arch. Rat. Mech. Anal. **171** (2004), no. 2, 151–218.
28. C. Manzini and L. Barletti, *An analysis of the Wigner-Poisson problem with inflow boundary conditions*, Nonlinear Anal. **60** (2005), no. 1, 77–100.
29. C. Mouhot, *Rate of convergence to equilibrium for the spatially homogeneous Boltzmann equation with hard potentials*, Comm. Math. Phys. **261** (2006), no. 3, 629672.
30. C. Mouhot, *Enlarging the functional space of decay estimates on semigroups*, WSPC–Proceedings, 2009.
31. A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer (1983).
32. K. Pravda-Starov, *Subelliptic estimates for quadratic differential operators*, Amer. J. Math. **133** (2011), no. 1, 39–89.
33. J. Prüss, *On the spectrum of C_0 -semigroups*, Trans. Amer. Math.Soc. **284** (1984), 847857.
34. B. Simon, *Trace ideals and their applications*, Cambridge Univ. Press (1979).
35. C. Sparber, J. A. Carrillo, J. Dolbeault, and P. Markowich, *On the long time behavior of the Quantum Fokker-Planck Equation*, Monatsh. Math. **141** (2004), 237–257.

36. B. Vacchini, *Translation-covariant Markovian master equation for a test particle in a quantum fluid*, J. Math. Phys. **42** (2001), 4291-4312.
37. B. Vacchini and K. Hornberger, *Relaxation dynamics of a quantum Brownian particle in an ideal gas*, Eur. Phys. J. Special Topics **151** (2007).
38. E. Wigner, *On the quantum correction for the thermodynamical equilibrium*, Phys. Rev. **40** (1932), 742-759.
39. S. Zhang and E. Pollak, *Quantum dynamics for dissipative systems: A numerical study of the Wigner-Fokker-Planck equation*, J. Chem. Phys. **118** (2003), 4357-4364.

INSTITUTE FOR ANALYSIS AND SCIENTIFIC COMPUTING, TECHNICAL UNIVERSITY VIENNA, WIENER HAUPTSTR. 8, A-1040 VIENNA, AUSTRIA
E-mail address: `anton.arnold@tuwien.ac.at`

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN, 1 UNIVERSITY STATION C1200, TEXAS 78712, USA
E-mail address: `gamba@math.utexas.edu`

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN, 1 UNIVERSITY STATION C1200, TEXAS 78712, USA
E-mail address: `gualdani@math.utexas.edu`

CEREMADE, UNIVERSITÉ PARIS-DAUPHINE, PLACE DU MARECHAL DE LATTRE DE TASSIGNY F-75775 PARIS CEDEX 16, FRANCE
E-mail address: `mischler@ceremade.dauphine.fr`

DPMMS, CENTRE FOR MATHEMATICAL SCIENCES, CAMBRIDGE UNIVERSITY, WILBERFORCE ROAD, CAMBRIDGE CB3 0WA, UK, *On leave from:* DMA, ÉNS, 45 RUE D'ULM F-75230 PARIS CEDEX 05, FRANCE
E-mail address: `Clement.Mouhot@ens.fr`

DEPARTMENT OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE, M/C 249, UNIVERSITY OF ILLINOIS AT CHICAGO, 851 S. MORGAN STREET, CHICAGO, IL 60607, USA
E-mail address: `sparber@math.uic.edu`